

ECCO XLII Meeting


"Microbe & Microbiome Management for a Better Planet"



European Culture
Collections' Organisation

Book of Abstracts

Editors: Antonio Moretti, Giancarlo Perrone



Bari (Italy), 18-20 September 2024



Published by

National Research Council - Institute of Sciences of Food Production (CNR-ISPA)

Editors in Chief

Dr. Antonio Moretti
Dr. Giancarlo Perrone

Editorial and Organizing Board

Pamela Anelli (CNR-ISPA, Italy)
Vittorio Capozzi (CNR-ISPA, Italy)
Giuseppe Cozzi (CNR-ISPA, Italy)
Nicola De Simone (CNR-ISPA, Italy)
Luciana De Vero (CNR-ISPA, Italy)
Massimo Ferrara (CNR-ISPA, Italy)
Katia Gialluisi (CNR-ISPA, Italy)
Donato Magistà (CNR-ISPA, Italy)
Mario Masiello (CNR-ISPA, Italy)
Giuseppe Petruzzino (CNR-ISPA, Italy)
Antonio Pugliese (CNR-ISPA, Italy)
Stefania Somma (CNR-ISPA, Italy)
Antonia Susca (CNR-ISPA, Italy)

Digital Book by

Luciana De Vero (CNR-ISPA, Italy)
Giuseppe Petruzzino (CNR-ISPA, Italy)

National Research Council

Institute of Sciences of Food Production

© CNR Edizioni, 2024

Piazzale Aldo Moro, 7 - 00185 Roma

www.edizioni.cnr.it

ISBN: 978 88 8080 652 3 (Digital Edition)

DOI: 10.5281/zenodo.13752745



Dear Colleagues,

we are very pleased to announce that the XLII European Culture Collections' Organisation (ECCO) meeting will be held in the amazing city of Bari (Apulia region, Southern Italy), from 18 to 20 September 2024.

The ECCO meeting is organised by the Institute of Sciences of Food Production of the National Research Council of Italy (CNR-ISPA), and sponsored by the CNR-ISPA Agri-food Microbial Culture Collection (ITEM), and the International Society on Mycotoxicology.

The meeting will be focused on the study, management and valorisation of Microbe & Microbiome, as biological resources and bio-systems crucial tools for a better understanding of biological systems, to support quality of life and promote the transition towards a better planet. The conference promotes a holistic and integrated approach on the value of microbial resources and Microbial Biological Research Centers (mBRCs): i) to support basic and applied transdisciplinary sciences, ii) as resilience factors to deal with global changes and sustainable development, and iii) to promote human and animal health in the framework of a OneHealth perspective.

ECCO supports cooperation and scientific exchanges concerning all the activities of mBRCs. Annual ECCO meetings represent an opportunity for the scientific community of the field, to discuss current trends and future development in the culture collections activities.

We would like to invite you to be part of the XLII ECCO meeting that will take place at "Sala Aldo Moro" (Dipartimento di Giurisprudenza), University of Bari "Aldo Moro", near the central train station and the wonderful historic city centre.

Director of CNR-ISPA

Antonio Moretti

Curator of ITEM Collection

Giancarlo Perrone

Wednesday 18

1.00 p.m. - 2.30 p.m.	Registration and Poster placing
2.30 p.m. - 3.00 p.m.	Welcome addresses: Rector of University of Bari - <i>Prof. Stefano Bronzini</i> ; President of CNR Research Area - <i>Cinzia Giannini</i> ; Major of Bari - <i>Vito Leccese</i> ; President of the European Culture Collections' Organisation (ECCO) - <i>Dr Gerard J. M. Verkley</i> ; Co-Chairs of XLII ECCO Meeting - <i>Antonio Moretti</i> and <i>Giancarlo Perrone</i>
3.00 p.m. - 4.30 p.m.	Session 5 – Microbiomes preservation and exploitation - Chairs: Ferrara Massimo (Italy), Aznar Rosa (Spain)
3.00 p.m. - 3.25 p.m.	Microbiomes for the Industry - <i>Porcar Manuel, Spain</i> (Invited Speaker)
3.25 p.m. - 3.40 p.m.	Exploring soil microbiome preservation strategies: culturable fraction, metabolic profiling and metagenomics - <i>Visca Andrea, Italy</i>
3.40 p.m. - 3.55 p.m.	Microbiome Biobanking: The missing link - <i>Kostic Tanja, Austria</i>
3.55 p.m. - 4.10 p.m.	Cryopreservation and recovery of a complex hypersaline microbial mat community - <i>Grego Michele, France</i>
4.10 p.m. - 4.25 p.m.	Direct injection Mass Spectrometry for the Real-Time volatilomics in Food System Microbiomes: the potential of providing temporal dimension in multi-omics studies - <i>Corvino Antonia, Italy</i>
	Closing remark of session (5 min)
4.30 p.m. - 5.00 p.m.	Coffee break + poster session
5.00 p.m. - 6.30 p.m.	Session 2– Microbes from farm to fork - Chairs: De Vero Luciana (Italy), Oivanen Pekka (Finland)
5.00 p.m. - 5.25 p.m.	Using microbial diversity or complex communities for valorization of side-streams for food applications - <i>Bang-Berthelsen Claus Heiner, Denmark</i> (Invited Speaker)
5.25 p.m. - 5.40 p.m.	Are isolation, identification and preservation of microbial strains still useful for food applications? - <i>Baruzzi Federico, Italy</i>
5.40 p.m. - 5.55 p.m.	Soil Health and Agri-Food System Sustainability from a microbiology perspective: A Data-Driven Approach for Agricultural Policy and Practices - <i>Bevivino Annamaria, Italy</i>
5.55 p.m. - 6.10 p.m.	Be a QPS or not to be, that is the Quirky Paradox of Safety - <i>Chessa Luigi, Italy</i>
6.10 p.m. - 6.25 p.m.	Investigation on the microbial evolution of cow milk in the passage from stable to mountain pasture, and evaluation of Bitto cheese microbial community - <i>Zago Miriam, Italy</i>
	Closing remark of session (5 min)
7.00 p.m.	Tour + Welcome Party

Thursday 19

9.00 a.m. - 10.30 a.m.	Session 3 – Pathogenic and beneficial aspects of microbes in human and animal health - Chairs: Hurtado Ortiz Raquel (France), Susca Antonia (Italy)
9.00 a.m. - 9.25 a.m.	Studying host-microbe interactions in health and disease using germ-free and gnotobiotic mouse technology - <i>Vereecke Lars, Belgium</i> (Invited Speaker)
9.25 a.m. - 9.40 a.m.	Using DNA Metabarcoding as a non-invasive tool for the conservation of the critically endangering Seychelles bat <i>Coleura seychellensis</i> - <i>Mattarelli Paola, Italy</i>
9.40 a.m. - 9.55 a.m.	Riboflavin overproducing food-grade bacteria: genomic basis, biotechnological applications and perspectives - <i>Capozzi Vittorio, Italy</i>
9.55 a.m. - 10.10 a.m.	Defining the <i>Enterobacter cloacae</i> species complex, with particular emphasis on <i>Enterobacter hormaechei</i> - <i>Rahi Praveen, France</i>
	Closing remark of session (5 min)
10.15 a.m. - 10.30 a.m.	Sponsor Session
10.30 a.m. - 11.00 a.m.	Coffee break + poster session
11.00 a.m. - 12.30 p.m.	Session 4 – Microbial life in extreme habitats: a 21st century challenge - Chairs: Varese Giovanna Cristina (Italy), Sedláček Ivo (Czech Republic)
11.00 a.m. - 11.25 a.m.	Diversity of microalgae in low pH environments: fifty years of the ACUF Collection, from strain to microbiome conservation and exploitation - <i>Pollio Antonino, Italy</i> (Invited Speaker)
11.25 a.m. - 11.40 a.m.	Life at the limits: diversity, adaptation strategies and bioprospecting of microbes living in Arctic deep-sea habitats (INDEPTH) - <i>Kaczorowska Anna-Karina, Poland</i>
11.40 a.m. - 11.55 a.m.	Cold-adapted carboxylic ester hydrolases from two Antarctic <i>Psychrobacter</i> strains: genomic analyses and in-vitro studies - <i>Cattaneo Andrea, Italy</i>
11.55 a.m. - 12.10 p.m.	Cold-adapted yeasts: a restricted club of extremophilic organisms - <i>Buzzini Pietro, Italy</i>
12.10 p.m. - 12.25 p.m.	Preliminary investigations of alkali-tolerant fungi in stromatolites from Lake Salda (Burdur province, SW Türkiye) - <i>Cecchi Grazia, Italy</i>
	Closing remark of session (5 min)

12.30 p.m. - 2.00 p.m. Lunch

2.00 p.m. - 3.30 p.m. Session 1 – Microbes for environmental sustainability, under a climate change scenario - Chairs: Turchetti Benedetta (Italy), Kermode Anthony (United Kingdom)

- 2.00 p.m. - 2.25 p.m. The importance of fungi for food security and One Health under a climate change scenario - *Lange Lene, Denmark (Invited Speaker)*
- 2.25 p.m. - 2.40 p.m. Fungi in marine plastisphere: ecological role and biotechnological potential - *Varese Giovanna Cristina, Italy*
- 2.40 p.m. - 2.55 p.m. Exploiting the agri-food waste and by-products potential for bioplastic production through *Haloferax mediterranei* fermentation - *Montemurro Marco, Italy*
- 2.55 p.m. - 2.10 p.m. Plant growth-promoting bacterial consortia isolated from halophytes to improve crop response to salinisation and climate change - *Racioppo Angela, Italy*
- 2.10 p.m. - 2.25 p.m. Genomic and metabolomic characterization of a new toxic cyanobacteria - *Rita Cordeiro, Portugal*
- Closing remark of session (5 min)

3.30 p.m. - 4.00 p.m. Coffee break + poster session

4.00 p.m. - 5.30 p.m. Session 6 – Microbes and Citizen Sciences: through the lens of public opinion - Chairs: Moretti Antonio, (Italy) Perrone Giancarlo (Italy)

- 4.00 p.m. - 4.25 p.m. The Isala project: characterizing the female microbiome through citizen science - *Wittouck Stijn, Belgium (Invited Speaker)*
- 4.25 p.m. - 5.25 p.m. Side show for citizen - interactive demonstrations
- 5.25 p.m. - 5.45 p.m. Results of the Survey on microbial biodiversity and taxonomy - *Felis Giovanna, Italy*
- 5.45 p.m. - 7.30 p.m. ECCO General Assembly
- 8.30 p.m. Social Dinner

Friday 20

9.30 a.m. - 11.45 a.m. Session 7 – Advanced approaches in taxonomy, phylogeny and functional genomic - Clermont Dominique (France), Masiello Mario (Italy)

- 9.30 a.m. - 9.55 a.m. Metabarcoding with Nanopore MinION: not only 16S or ITS - *Faino Luigi, Italy (Invited Speaker)*
- 9.55 a.m. - 10.10 a.m. Genomic overview over 100 cyanobacterial strains: taxonomy, microbiome and biosynthetic gene clusters - *Rúben Luz, Portugal*
- 10.10 a.m. - 10.25 a.m. Drivers of aquatic fungal diversity and ecological functions throughout Europe: Biodiversa+ FUNACTION and MoSTFun projects - *Marchese Pietro, Italy*
- 10.25 a.m. - 10.40 a.m. ITSoneDB v1.144 and BioMaS@ITSoneWB: two ELIXIR-IT main resources for amplicon based mycobiome investigation - *Defazio Giuseppe, Italy*
- 10.40 a.m. - 11.10 a.m. Coffee break + poster session
- 11.10 a.m. - 11.25 a.m. The new challenge in virus taxonomy: a binomial nomenclature for virus species - *Rubino Luisa, Italy*
- 11.25 a.m. - 11.40 a.m. Genomic insights into Mrakia: expanding horizons in microbial biotechnology - *Di Cesare Francesca, Italy*
- Closing remark of session (5 min)

11.45 a.m. - 1.20 p.m. Session 8 – Networking, services, quality and data management of Microbial culture collections - Chairs: Lima Nelson (Portugal), Verkley Gerard (Netherlands)

- 11.45 a.m. - 12.10 p.m. MIRRI-ERIC as a microbial, genetics and data resources hub to advance scientific discovery - *Portugal Melo Ana, Portugal (Invited Speaker)*
- 12.10 p.m. - 12.25 p.m. PLAVIT, the Italian Plant Virus Collection in 2024 - *Accotto Gian Paolo, Italy*
- 12.25 p.m. - 12.40 p.m. How the MIRRI-PT has been developed in its smart specialisation and cutting-edge technologies to offer better services - *Lima Nelson, Portugal*
- 12.40 p.m. - 12.55 p.m. The Italian network of microbial culture collections: an overview on management and sustainability of the future research infrastructure MIRRI-IT – *Moretti Marino, Italy*
- 12.55 p.m. - 1.10 p.m. Managing change within a culture collection: A case study on the challenges within quality, data management and infrastructure within CABI's collection - *Kermode Anthony, United Kingdom*
- Closing remark of session and Ecco meeting (15-20 min)
- 1.30 p.m. - 3.00 p.m. Closing lunch

Abstracts Summary

Session 1 – Microbes for environmental sustainability, under a climate change scenario.....	1
Thursday 19th September, 2:00pm.....	1
The importance of fungi for food security under climate change scenario	2
Fungi in marine plastisphere: ecological role and biotechnological potential.....	3
Exploiting the agri-food waste and by-products potential for bioplastic production through <i>Haloferax mediterranei</i> fermentation	4
Plant growth-promoting bacterial consortia isolated from halophytes to improve crop response to salinisation and climate change	5
Genomic and metabolomic characterization of a new toxic cyanobacteria.....	6
Microorganisms for the environment: Technological robustness of Plant Growth Promoting Bacteria for Mediterranean crops	7
Microbial characterization of endophytic bacteria from halophytic plants	8
Microbial collection from vegetable compost tea to develop bacterial consortia suitable for plant disease control	9
Dietary microbes and Mediterranean diet	10
Fungi as bioresources for remediation of HCH-contaminated soils: from microbial community-level physiological profile to selective isolation in enrichment	11
From culture collections to biorecovery of strategic elements: exploring fungal interactions with Germanium and Indium	12
Microbial consortia the great bio-based solution for the bioremediation of co-contaminated soils: study of microbial resources isolated from a decommissioned military site.	13
The CABI Biological Resource Collections: evolving to meet the changing needs of users.....	14
Exploring methodological aspects of microbiota transplantation in black elder (<i>Sambucus nigra</i>).....	15
Plant growth promotion endophytes from Sicilian native grapevines as plant biostimulants for climate-ready crops	16
Exploring Sicilian olive tree eco-friendly endophytes for sustainable agriculture	17
Fungal diversity in the precious Mediterranean red coral <i>Corallium rubrum</i>	18
A network-based approach to elucidate the dynamics of the native microbiota's response to a multi-kingdom microbial inoculant in raw brewer's spent grain	19
Exploring the oenological characteristics of different non- <i>Saccharomyces</i> species in grape must fermentation.	20
Harnessing Microbiomes for Drought-Resilient Agriculture: Insights from the BIOMEnext Project.....	21
Harnessing Fungi for Eco-Friendly Biosurfactant Production: joint efforts from CECT and MUT collections to unveil this potential among the preserved strains	22
Changes in bacterial, archaeal, and fungal communities from saline soils cultivated with tomatoes in Italy	23
A polycyclic aromatic hydrocarbons (PAHs) contaminated soil as source of microorganisms for bioremediation purposes	24
Impact of newly developed biofertilizer “polyazorich” on soil agrochemical properties during crop cultivation	25
Session 2 – Microbes from farm to fork.....	26

Wednesday 18th September, 5:00pm	26
Using microbial diversity or complex communities for valorization of side-streams for food applications.....	28
Are isolation, identification and preservation of microbial strains still useful for food applications?	29
Soil Health and Agri-Food System Sustainability from a microbiology perspective: A Data-Driven Approach for Agricultural Policy and Practices	30
Be a QPS or not to be, that is the Quirky Paradox of Safety	31
Investigation on the microbial evolution of cow milk in the passage from stable to mountain pasture, and evaluation of Bitto cheese microbial community	32
Influence of the cheesemaking process on the bioprotection activity of <i>Lactococcus</i> strains	33
Dietary Microbes and Case Studies from Food Processing	34
Creation of an “Egg-to-Meat” biobank of microbiota collected from broiler chickens raised with or without outdoor access.	35
<i>Lactiplantibacillus plantarum</i> diversity and plant-based sources	36
<i>Lactiplantibacillus plantarum</i> and bio-based solutions in foodbiotechnologies.....	37
Enhancing bioprotection on fresh-cut apples with microwave heating: preliminary tests for inclusion in hurdle technology	38
Biocontrol, edible packaging, and physical technologies: towards the design of integrated solutions for postharvest fruitsand vegetable management.....	39
Application of lactic acid bacteria in the postharvest of fruits and vegetables	40
Microbial Diversity from fermented Apulian ‘ <i>Gentile di Puglia</i> ’sheep milk: polyphasic characterization, dairy valorisation and cross-over cereal based applications.....	41
Characterization of the rib operon riboswitch in riboflavinoverproducing <i>Lactiplantibacillus plantarum</i> strains	42
Bioprospecting and valorization of citrus crops to obtain preharvest and post-harvest biocontrol agents.....	43
Fermentation as a driver of innovation for the development of the agri-food sector in marginal areas.....	44
Innovation in food fermentations and the cross-over fermentation concept in new product development	45
Selection of <i>Bacillus strains</i> as potential candidate against pathogenic <i>Fusarium</i> occurring on cereal.....	46
Effect of plant essential oils on <i>Botrytis cinerea</i>	47
Lactic acid bacteria as microbial starters in malting by-product fermentation to obtain innovative bread formulation.....	48
Unconventional raw plant matrices as source of pro-technologicalmicrobes for application in food fermentation.	49
Real-Time investigation of <i>Lactiplantibacillus plantarum</i> Fermentation in Experimental and Real Food Conditions Using a Metabolomics Approach.....	50
Cell surface proteome of <i>L. plantarum</i> strains isolated from vegetable foods and dairy products	51
Biodiversity in natural fungal community isolated from cheese aged in Apulian cave	52
Dried cell-free supernatant of <i>Bacillus subtilis</i> strain ET-1against <i>Penicillium digitatum</i> on clementine fruits	53
<i>Penicillium nalgioense</i> : a promising fungal starter to control <i>Aspergillus westerdijkiae</i>	

contamination in salami	54
Biodiversity in fungal community on cheese aged in Dossena mine	55
Proteolysis in soy tempeh obtained with commercial starter	56
Relationship between the chemical content of fruits and endophytic fungal communities in black elder <i>Sambucus nigra</i> L.....	57
Genomic characterization of <i>Companilactobacillus</i> spp.: assessing safety and application potential in the food chain.....	58
Screening of yeast and filamentous fungi from VTT Culture Collection for cacao butter and palm oil production.....	59
Diversity of Non-Conventional Yeasts (NCY) for innovating fermented beverages: investigating microbiomes	60
Principles for selection of strains of microorganism cultures for the creation of biopreparations of agricultural designation.....	61
Tailoring diverse microbial consortia for the improvement of plant nutritional and nutraceutical properties.....	62
Session 3 – Pathogenic and beneficial aspects of microbes in human and animal health	63
Thursday 19th September, 9:00am.....	63
Studying host-microbe interactions in health and disease using germ-free and gnotobiotic mouse technology.....	66
Using DNA Metabarcoding as a Non-Invasive Tool for the Conservation of the Critically Endangering Seychelles Bat <i>Coleura seychellensis</i>	67
Riboflavin overproducing food-grade bacteria: genomic basis, biotechnological applications and perspectives	68
Defining the <i>Enterobacter cloacae</i> species complex, with particular emphasis on <i>Enterobacter hormaechei</i>	69
Functionalization of chickpeas through <i>Saccharomyces cerevisiae</i> var. <i>boulardii</i>	70
Growth inhibition of Human Pathogenic Microorganisms (HPMOs) by endophytic and rhizosphere bacteria.....	71
Supercritical fluid extraction of torularhodin from a new red yeast strain.....	72
Interaction between probiotic lactic acid bacteria and immunosuppressive drugs	73
Factors affecting the production of γ -aminobutyric acid in <i>Levilactobacillus brevis</i> LB12.....	74
Comparison of microbial populations of peritonsillar abscess determined by molecular and bacteriological methods.....	75
Use of potential probiotic autochthonous <i>L. plantarum</i> as co-starter for the production of goat's milk cheeses	76
Session 4 Microbial life in extreme habitats: a 21st century challenge.....	77
Thursday 19th September, 11:00am.....	77
Diversity of microalgae in low pH environments: fifty years of the ACUF Collection, from strain to microbiome conservation and exploitation.....	78
Life at the limits: diversity, adaptation strategies and bioprospecting of microbes living in Arctic deep-sea habitats (INDEPTH)	79
Cold-adapted carboxylic ester hydrolases from two Antarctic <i>Psychrobacter</i> strains: genomic analyses and in-vitro studies	80
Cold-adapted yeasts: a restricted club of extremophilic organisms	81
Preliminary investigations of alkali-tolerant fungi in stromatolites from Lake Salda (Burdur	

province, SW Türkiye).....	82
From microbial diversity of Campania thermal springs of their maintenance and preservation	83
The influence of prokaryote communities on <i>Galdieria sulphuraria</i> strain maintenance and cultivation	84
Cryopreservation strategies for extremophilic microalgae at ACUF: growth kinetic and photosynthetic assessment.....	85
Fermentation Technologies for the Optimization of levan Production by <i>Pseudomonas</i> strain 2ASCA.....	86
Extremophiles Collection of Institute of Biomolecular Chemistry National Research Council (CE-ICB CNR): microbial biodiversity preservation of extreme environments within the SUS-MIRRI.IT project	87
Exopolysaccharides from psychrophilic microorganisms as biological device for plant promotion.	88
Genome Prospecting and Wet-Lab Evidence of Plant Growth Promotion Traits in the Psychrophilic Arctic Bacterium <i>Pseudomonas</i> strain 2ASCA.....	89
Biosynthesis of Nanoparticles by Extremophilic Bacteria for Potential Biotechnological Application.....	90
The Haloarchaea Morphological Atlas discloses animal-like tissue structures emergence in single-celled archaea	91
Session 5 – Microbiomes preservation and exploitation.....	92
Wednesday 18th September, 3:00pm	92
Microbiomes for the Industry.....	95
Exploring soil microbiome preservation strategies: culturable fraction, metabolic profiling and metagenomics	96
Microbiome Biobanking: The missing link.....	97
Cryopreservation and recovery of a complex hypersaline microbial mat community.....	98
Direct Injection Mass Spectrometry for the Real-Time Volatilomics in Food System	
Microbiomes: the Potential of Providing Temporal Dimension in Multi-Omics Studies.....	99
Possible “microbial bioremediators” isolated from wasted lands.....	100
Food microbiomes and packaging: past, present and future.....	101
Marine microbial community sample cryopreservation: finding the best preservation strategy	102
Fermenting metacommunities of food interest: an invaluable biodiversity asset to be maintained overtime in its wholeness and functionality Micro4ever	103
Biochemical and biophysical characterization of two recombinant enzymes obtained from termite gut metagenome	104
Exploring Microbiomes in PFAS Contaminated Environments: Preservation for Bioremediation Strategies.	105
UMCC’s investigative approach for the preservation and exploitation of fermented beverage microbiomes	106
A cryopreservation protocol of microbial consortia from Apulian table olives	107
Synthetic microbial communities as a tool to understand microbial interactions and their role in bacterial cellulose production	108
Back-slopping of dried, freeze-dried and frozen Tempeh: the effect of long-term preservation techniques on the microbiome and volatile.....	109

New frontiers for <i>Saccharomyces cerevisiae</i> flor strains.....	110
Microbiome characterisation of sourdough from the MBDS-UNISSCC microbial collection.....	111
Culturomics and microbiome-based approach for the development of NatComs as biofertilizers and biocontrol agents for sustainable agriculture: the reinforcement of ENEA microbial culture collection.....	112
<i>Trichoderma asperellum</i> and <i>Bacillus</i> spp. antifungal activity evaluation against <i>Fusarium</i> fungi.....	113
The use of microbial inoculants for the control of <i>Fusarium</i> related diseases in durum wheat	114
Session 6 – Microbes and Citizen Sciences: through the lens of public opinion.....	115
Thursday 19th September, 4:00pm.....	115
The Isala project: characterizing the female microbiome through citizen science.....	116
Mediterranean Diet and microbes: an emerging target for educational paths and citizen science.....	117
Didactic Proposals in Education for Sustainable Development in Teaching in Portugal and Brazil: Circular Relationship between Climate Change, Healthy Ecosystems and Soil Microbial Diversity.....	118
Session 7 – Advanced approaches in taxonomy, phylogeny and functional genomics	119
Friday 20th September, 9:30am.....	119
Metabarcoding with Nanopore MinION: not only 16S or ITS.....	120
Genomic overview over 100 cyanobacterial strains: taxonomy, microbiome and biosynthetic gene clusters	121
Drivers of aquatic fungal diversity and ecological functions throughout Europe: Biodiversa+ FUNACTION and MoSTFun projects	122
ITSoneDB v1.144 and BioMaS@ITSoneWB: two ELIXIR-IT main resources for amplicon based mycobiome investigation	123
The new challenge in virus taxonomy: a binomial nomenclature for virus species.....	124
Genomic insights into <i>Mrakia</i> : expanding horizons in microbial biotechnology	125
New Enterobacterales species isolated from dead seals in Antarctica	126
Taxonomic classification of Metagenome-Assembled Genomes with kMetaShot	127
Presumptive identification of two Gram-positive microaerophilic strains isolated from Weddell's seals in Antarctica.	128
New protocol development of bioinformatics analysis for species verification of the genus <i>Pseudomonas</i> from Polish Collection of Microorganisms resources.	129
Novel insights into heterocyte patterning and proheterocyte germination in <i>Kaarinaea lacus</i> PCC 9237T gen. nov. sp. nov.	130
Exploring <i>Fusarium oxysporum</i> biodiversity in Beauvericin gene-cluster	131
Exploring cyanobacteria genomes for antithrombotic secondary metabolites.....	132
GEN-ERA toolbox: reproducible bioinformatics workflows for genomics and taxonomy in culture collection.	133
Comparison study between molecular fingerprinting and FTIRBIOTYPER.....	134
Advancing research in synthetic biology through DTU Bioengineering's Fermentation and HT-automation facility	135
Identification and characterization by a polyphasic approach of the microbial agent responsible of	

spoilage in Grana Padanocheese	136
The importance of the <i>Legionella surveillance</i> approach implemented in the company's water distribution system to discover a novel <i>Legionella</i> species: <i>Legionella petroniana</i> sp. nov.....	137
Session 8 – Networking, services, quality and data management of Microbial culture collections .	138
Friday 20th September, 11.45:00am.....	138
MIRRI-ERIC as a microbial, genetics and data resources hub to advance scientific discovery	140
How the MIRRI-PT has been developed in its smart specialisation and cutting-edge technologies to offer better services.....	141
PLAVIT, the Italian Plant Virus Collection in 2024	142
The italian network of microbial culturecollections: an overview on management and sustainability of the future research infrastructure MIRRI-IT.....	143
Managing change within a culture collection: A case study on the challenges within quality, data management and infrastructure within CABI's collection.....	144
Establishment of the ITEM Culture Collection backup and development of the related disaster recovery plan	145
A cloud-based, open-access platform for comprehensive metagenomic data analysis	146
Strengthening the CoSMi culture collection: a contribution to preserving the biodiversity of microalgae	147
The SUS-MIRRI.IT project: research, services, and training for sustainable bioscience and bioeconomy by the MIRRI Italian Research Infrastructure.....	148
The Usage and Fate of Patent Deposits.....	149
Performing an upgraded and interoperable data management system for ITEM Microbial Culture Collection	150
Bank of Algae and Cyanobacteria of the Azores – BACA culture collection	151
A ranking tool for microbial biobanks.....	152
A shared quality management system for Italian Culture Collections: The challenge of SUS-MIRRI.IT project.....	153
The ENEA Microbial Culture Collection as valuable resource for developing new sustainable biotechnologies and bioeconomy.....	154
Speaking the same language: Changing the way to share microbial strain data.....	155

Session 1 – Microbes for environmental sustainability, under a climate change scenario

Thursday 19th September, 2:00pm

Chairs: Turchetti Benedetta (Italy), Kermode Anthony (United Kingdom).

The importance of fungi for food security under climate change scenario

Lange, Lene¹

¹ *LL-BioEconomy, Denmark*

Fungi can play a major role in improving food security, achieved by upgrading what is now wasted. This is of special importance, when climate change is causing serious obstacles for agricultural food production. Fungal enzymes can valorize crop residues, agro-industrial side-streams and left over food, all being food compatible resources, now underutilized or even wasted. Enzymes can open recalcitrant structures and enable the recovery of nutritious components such as proteins, dietary fibers, vitamins, lipids and antioxidants from such bioresources. The fungi instrumental for such significant improved use of the (global and local) bioresources are well studied; making it obvious that also a rather low number of fungal strains, can be key for sustainable and safe improved use of the bioresources. Significant progress could be made if many more throughout the world had access to such strains and how to use them. Culture collections can play a vital and major role in making such an ambitious plan into reality. Proposal: Each culture collection establishes and promotes a set of cultures of fungi (and bacteria) which are documented to be safe and shown to be efficient and with no strings attached, with regard to use. Such set of cultures, being fungal strains, producing blends of biomass degrading enzymes; or fungal strains, easy to use for recombinant production; or fungal hosts, for biological production (preferably efficient also when grown on residues); or fungi for fermentation of residues, resulting in tasty, nutritious and gut health promoting food. The Culture collections are hereby encouraged to organize collaborative projects and seek funding (public or philanthropy) for developing such sets of cultures for Circular and Biobased food production, including funding for providing such sets of cultures in countries or regions, where climate change is seriously challenging food security, nutrition and health.

Fungi in marine plastisphere: ecological role and biotechnological potential

Spina, Federica¹; Stilo, Giulia¹; Venice, Francesco¹; Fiorin, Anita¹; Reginato, Alessia¹; Florio Furno, Matteo¹; Bracco, Perangiola¹; Dodhia, Maya²; Zanetti, Marco³; Posth, Nicole R.²; Varese, Giovanna Cristina¹

¹ University of Torino, Department of Life Sciences and Systems Biology, Torino, Italy; ² University of Copenhagen, Department of Geosciences and Natural Resource Management Geology, Sedimentary Systems, København, Denmark; ³ University of Torino, Department of Chemistry, Torino, Italy

Plastic released into the environment represents one of the most important environmental problems with important repercussions on the functionality of ecosystems and on the health of animals, including humans. Every year, marine environment receives more than 10 million tons of plastic which is quickly colonized by microorganisms. The co-occurrence of bacteria, fungi and microalgae creates a dynamic scenario which remains largely unexplored as regard the fate of the different plastics (traditional versus biodegradable plastics), the effect of abiotic features as salinity and the microbes involved in the possible biodegradation.

In the present study, the marine plastisphere of two Danish locations (characterized by different salinities) was studied through culture dependent and independent techniques, unveiling a heterogeneous mycobiota.

The fungal community was various, with a strong role of the site, the plastic polymer (biodegradable polymer versus a non-biodegradable polymer) and environmental parameters as salinity. Hundreds of filamentous fungi and yeasts were isolated, mostly associated with Ascomycota phylum. Unveiling the hidden biotechnological potential of marine fungi is of major interest, focusing on their capability to transform plastic polymers. Some fungi were able to use different plastics as sole carbon source and capable of degrading biopolymers. The extent and the kinetics of such process are currently under investigation via respirometry analysis. Moreover, the genome annotation of some of these strains highlights the presence of putative Plastic Active Enzymes (PAZy).

Acknowledgements

Part of this work was granted by the European Commission—NextGenerationEU, Project “Strengthening the MIRRI Italian Research Infrastructure for Sustainable Bioscience and Bioeconomy”, coden. IR0000005.

Exploiting the agri-food waste and by-products potential for bioplastic production through *Haloferax mediterranei* fermentation

Montemurro, Marco¹; Longo, Angela²; Fanelli, Francesca¹; Villano, Marianna³; Rizzello, Carlo Giuseppe²

¹ Institute of Sciences of Food Production, National Research Council (ISPA-CNR), Bari, Italy;

² Department of Environmental Biology, "Sapienza" University of Rome, Italy; ³ Department of Chemistry, Sapienza University of Rome, Rome, Italy

Several microorganisms can synthesize polyhydroxyalkanoates (PHA) which represent environmentally friendly substitutes for conventional plastic. The halophilic *Haloferax mediterranei* has been largely investigated for its ability to produce the copolymer Poly(3-hydroxybutyrate-co-3-hydroxyvalerate) (PHBV) using organic waste as carbon sources and in high-salinity environments without requiring sterilization. Different authors studied genes and pathways responsible for PHBV production, thus allowing the identification and regulation of fermentation parameters for enhancing cell growth and increasing PHBV accumulation.

Different food waste has been tested for PHBV production by *Haloferax mediterranei*. Among them, the whey from cheese production and starch-based byproducts have been widely investigated. The use of whey required a-galactosidase treatment to hydrolyze the lactose and the addition of trace element solution (SL6) to obtain a final concentration of PHBV of 1.18 g/L. The production of wasted bread-derived substrate was moreover optimized including seawater instead of the SL6 to reduce the costs for media supplementation. In this case the production under optimized conditions of fermentation was 1.53 g/L. Both productions were conducted in a 3-liter bioreactor. However, the amount of biopolymer produced is not the only parameter to assess. The characterization of PHBV and the importance of hydroxyvalerate (HV) abundance in the formation of PHBV copolymer is usually evaluated to assess the properties of this biopolymer and the downstream processing options can influence this composition. Nowadays several crucial factors associated with industrial scale-up still need to be evaluated at industrial scale to obtain a sustainable and economic alternative to the other biopolymers using inexpensive food waste and *Haloferax mediterranei*.

Acknowledgements

This study was carried out at the Agritech National Research Center, and it received funding from the European Union Next-GenerationEU (PIANO NAZIONALE DI RIPRESA E RESILIENZA (PNRR)—MISSIONE 4 COMPONENTE 2, INVESTIMENTO 1.4—D.D. 1032 17/06/2022, CN000000022).

This manuscript reflects only the authors' views and opinions; neither the European Union nor the European Commission can be considered responsible for them.

Plant growth-promoting bacterial consortia isolated from halophytes to improve crop response to salinisation and climate change

Racioppo, Angela¹; Guerrieri, Angela¹; Speranza, Barbara¹; Botticella, Lucia¹; Conversa, Lucia¹; Corbo, Maria Rosaria¹

¹ Department of the Science of Agriculture, Food, Natural Resources and Engineering, University of Foggia, Foggia, Italy

Salinisation of soils is one of the main biotic stresses impacting on agricultural lands [1]. The use of beneficial bacteria adapted to high salinity conditions is a viable option for improving crop production in salinity affected areas, given that saline soil reclamation is a complex process. Halotolerant plant growth promoting bacteria (PGPB) are able to enhance plant growth by increasing soil carbon, nitrogen and mineral availability and uptake [2].

Over the years, coastal and saline regions, including the area of Margherita di Savoia saltworks, north-eastern Apulia (Italy), have proven to be a natural source of beneficial microbes adapted to high salinity. The area of Margherita di Savoia saltworks is said to have a unique, diverse and rich indigenous microbiota that is still largely unexplored.

Therefore, the aim of the present study was to isolate, characterise and select potential PGPB from *Cakile maritima* plants collected in two sites located in Margherita di Savoia. In particular, the microbiological sampling was carried out in triplicate at three phenological stages (seedling, vegetative growth, flowering) of plant life cycle, during the year 2023. A total of 180 (150 rhizobacteria and 30 endophytes) bacteria were isolated and characterised by using morphological, biochemical, and molecular approaches.

The halotolerant isolates which possessed plant growth promoting traits including phosphate, and silicon solubilization, indole acetic acid, and siderophores production, ammonia generation, drought and salt resistant were selected. Further, the effect of three halotolerant isolates which showed most prominent PGPB activities, and genotypically identified, was evaluated in a growth chamber on

C. maritima plants under high salinity conditions. The halotolerant isolates improved the plant growth especially under moderate saline conditions in comparison to non-inoculated control plants.

References

- [1] Mukhtar, S, Malik KA, Mehnaz S. (2019). Microbiome of Halophytes: Diversity and Importance for Plant Health and Productivity. Microbiology and Biotechnology Letters, 47(1), 1–10.
- [2] Racioppo A, d'Amelio A, De Santis A, Bevilacqua A, Corbo MR, Sinigaglia M. (2023). Potential Use of Plant Growth-Promoting Bacteria to Enhance Growth and Soil Fertility in Marginal Areas: Focus on the Apulia Region, Italy. Agronomy, 13(12), 2983.

Acknowledgments

This study was carried out within the: Agritech National Research Center and received funding from the European Union Next-Generation EU (PIANO NAZIONALE DI RIPRESA E RESILIENZA (PNRR)—MISSIONE 4 COMPONENTE 2, INVESTIMENTO 1.4—D.D. 1032 17/06/2022,CN

Genomic and metabolomic characterization of a new toxic cyanobacteria

Cordeiro, Rita¹; Luz, Rúben¹; Fonseca, Amélia¹; Gonçalves, Vitor¹

¹ CIBIO, Centro de Investigação em Biodiversidade e Recursos Genéticos, InBIO Laboratório Associado, BIOPOLIS Program in Genomics

Cyanobacteria genomes and genomic analysis have been increasing in recent years since genomics allows for more complete studies and a deeper understanding of several matters, such as taxonomy and secondary metabolites bioprospection. A previous work on BACA's (Bank of Algae and Cyanobacteria of the Azores) cyanobacteria diversity, based on 16S rRNA phylogenetics, revealed the first evidence of the high and undescribed diversity of cyanobacteria in the Azores islands. This work reported four toxic strains, two cylindrospermopsin producers (by LC-MS/MS), BACA0025 and BACA0031, with 16S rRNA phylogenetics indicating it as a possible new genus. Here, we aimed to produce and explore these strains genomes for a deeper characterization. Genomes were sequenced through Illumina and, assembled and annotated following Luz et al. [1]. Genomic characterization was based on phylogenomics, average nucleotide identity (ANI), average amino acid identity (AAI) and biosynthetic gene clusters (BGCs) identification. Metabolomic analysis was assessed through HPLC-MS. Phylogenomics, ANI and AAI revealed evidence that strengthened the description of a new genus within the Aphanizomenaceae family. Identification of BGCs revealed several secondary metabolites besides the previously identified cylindrospermopsin [2], such as Anabaenopeptins and heterocyst glycolipids. The comparison of BGCs between producing cyanobacteria also allowed a deeper characterization. The metabolomic analysis confirmed metabolites identified by genomics, such as Anabaenopeptins and the Cylindrospermopsin analog. This work confirms the rise of a new toxic cyanobacteria genus, isolated from several freshwater lakes in Pico island (Azores, Portugal), producer of secondary metabolites of interest, thus highlighting the uncovered cyanobacteria diversity in the Azores and its biotechnological potential.

References

- [1] Cordeiro R., et al. 2021. Cyanotoxin screening in BACA culture collection: identification of new cylindrospermopsin producing cyanobacteria. *Toxins* 13: 258.
- [2] Luz R., et al. 2023. Description of *Pseudocalidococcus azoricus* gen. sp. nov. (Thermosynechococcaceae, Cyanobacteria), a rare but widely distributed coccoid cyanobacteria. *Diversity* 2023, 15, 1157.

Acknowledgments

This work was funded by FEDER funds through the Interreg-MAC 2014-2020 Program under the projects REBECA (MAC1.1a/060) and REBECA-CCT (MAC2/1.1b/269). By Portuguese National Funds, through FCT, the European Union, QREN, FEDER, COMPETE, by funding the CIBIO/InBIO (project UID/BIA/ 50027/2013 and POCI-01-0145-FEDER-006821).

Microorganisms for the environment: Technological robustness of Plant Growth Promoting Bacteria for Mediterranean crops

d'Amelio, Annalisa¹; De Santis, Michele¹; Bevilacqua, Antonio¹; Flagella, Zina¹; Sinigaglia, Milena¹; Corbo, Maria Rosaria¹

¹ *University of Foggia, Foggia, Italy*

Soil degradation caused by climate change could threaten economically important crops, such as durum wheat, especially in the Mediterranean regions where climatic and meteorological conditions are strong contributors. The aim of modern agriculture is therefore to find sustainable solutions to optimise the biological and economic productivity of the soil resources. Plant Growth Promoting Bacteria (PGPB) are the most promising approach to achieve these goals, as they offer a wide range of benefits in agriculture, including increasing crop productivity, improving soil nutrient levels and restoring soil fertility [1]. However, to use them properly, it is necessary to understand their limits, for example, whether they can be applied in different conditions from those in which they were isolated and developed. Thus, this research aims at assessing the technological robustness of some PGPB strains, focusing on the resistance to commercial fungicides, adhesion to seeds to design a protocol for bacteria use in field, growth profiles as a function of pH and temperature, persistence in soil also under extreme conditions.

The strains were generally resistant to fungicides; in addition, the adhesion performance was >80% and for many microorganisms at 95-98%.

Concerning the persistence in soil, they show a prolonged viability, although the increase of temperatures (at 45°C) could be a challenge.

Generally, the results show a high technological robustness of the strains and their potentiality as active ingredients of bioformulates.

References

- [1] Racioppo, A.; d'Amelio, A.; De Santis, A.; Bevilacqua, A.; Corbo, M.R.; Sinigaglia, M. (2023) Potential Use of Plant Growth-Promoting Bacteria to Enhance Growth and Soil Fertility in Marginal Areas: Focus on the Apulia Region, Italy. *Agronomy*, 13, 2983.

Microbial characterization of endophytic bacteria from halophytic plants

Guerrieri, Angela¹; Speranza, Barbara¹; Racioppo, Angela¹; Giancaspro, Angelica¹; Lotti, Concetta¹; Corbo, Maria Rosaria¹

¹ Department of the Science of Agriculture, Food, Natural Resources and Engineering, University of Foggia, Foggia, Italy

Halophytic plants are adapted to harsh environmental conditions and can grow in coastal areas known for their high salinity. The physiology and metabolism of such plants are influenced by their microbiome, which includes a specific class of microorganisms: Plant Growth-Promoting Endophytic bacteria (PGPE), a heterogeneous group of microorganisms that inhabit the interior of plant tissues and are known for their beneficial effects on growth and health of the host. PGPE facilitate plant development by several mechanisms similar to the rhizosphere bacteria, including fixation of N₂, solubilization of phosphate, production of indole acetic acid and siderophores, and greater availability and uptake of carbon, nitrogen, and minerals from the soil. The identification of endophytic microorganisms appears to be an important tool for developing sustainable agriculture and improving the growth and stress tolerance of crops in marginal areas.

This study aimed to isolate and characterize endophytic bacteria from two halophytic plants: *Cakile maritima* Scop. and *Salicornia europea*, collected in the Margherita di Savoia saltworks, in the north-eastern Apulia (Italy). The isolation of 40 endophytes was carried out according to Christakis et al. [1]. All isolates were characterized by phenotypic (Gram staining, catalase, oxidase, and urease) and technological tests (phosphate solubilization, ammonium production, nitrification, siderophore production, drought and salt resistance, and silicon solubilization). DNA barcoding based on 16S rRNA sequencing was used for the identification of the isolates.

The obtained results suggest that halophytic plants harbour a variety of putative endophytic bacteria which exhibits tolerance to NaCl and displays different plant growth-promoting traits. Our findings suggest the possible implications of these PGPE in improving the quality and productivity of halophytes in saline soil and in protecting other important plants against salt stress.

References

- [1] Christakis CA, Daskalogiannis G, Chatzaki A, Markakis EA, Mermigka G, Sagia A, Rizzo GF, Catara V, Lagkouvardos I, Studholme DJ, Sarris PF. Endophytic Bacterial Isolates From Halophytes Demonstrate Phytopathogen Biocontrol and Plant Growth Promotion Under High Salinity. *Front Microbiol.* 2021, 4;12:681567.

Acknowledgments

This study was carried out within the Agritech National Research Center and received funding from the European Union Next-Generation EU (PIANO NAZIONALE DI RIPRESA E RESILIENZA (PNRR)—MISSIONE 4 COMPONENTE 2, INVESTIMENTO 1.4—D.D. 1032 17/06/2022, CN00000022).

Microbial collection from vegetable compost tea to develop bacterial consortia suitable for plant disease control

Sigillo, Loredana¹; Dell'Olmo, Eliana¹; Raio, Aida²; Senape, Veronica¹; Serratore, Giovanna¹; Napoletano, Giovanna¹; Zaccardelli, Massimo¹

¹ Council for Agricultural Research and Economics Research centre for Vegetable and Ornamental Crops, Pontecagnano, Italy; ² National Research Council – Institute for Sustainable Plant Protection Sesto Fiorentino (FI), Italy

Compost teas (CTs) are fermented aqueous extracts obtained from vegetable compost by extraction and/or fermentation processes in liquid phase. They are rich in organic and inorganic molecules active as plant biostimulants, and in microorganisms capable to counteract phytopathogenic fungi and bacteria. About 5% of the total CT microflora is cultivable in vitro: the most represented genera are *Bacillus*, *Pseudomonas*, *Serratia*, *Stenotrophomonas*, *Aspergillus*, *Flavobacterium*, *Fusarium*, *Gliocladium*, *Penicillium* and *Trichoderma*. The PLANTiA project, funded by AGER Agroalimentare e Ricerca (Ager Project – Third Edition), aims to select beneficial microorganisms from two vegetable CTs. One hundred isolates extracted from these CTs are characterized for their biocontrol activity against four phytopathogenic fungi of leguminous and grapevine crops: *Rhizoctonia solani* (RS) CREA OF 1333.1, *Macrophomina phaseolina* (MP) CREA OF 373.2, *Fusarium solani* (FS) 897.1 and *Botrytis cinerea* (BC) CREA_OF 1531. Preliminary results showed that the growth of RS appeared inhibited by 4 isolates, FS by 13 and BC by 6. MP growth was slightly affected by 10 isolates. The effective antagonistic bacteria were identified as *Pseudomonas*. They will be further characterized for species identity, motility, production of biofilm, indole acetic acid, protease, siderophores and volatile compounds, ability to fix nitrogen and to solubilize phosphate. Moreover, tolerance to extreme pH, salinity and temperature will be determined to select bacteria able to protect plants under unfavourable environmental conditions. The most promising accessions will be used for in vivo biocontrol assays. This work will allow to implement microorganism collections, to preserve microbial biodiversity and to detect microbial features related to the effectiveness of a microorganism in protecting the plants from biotic and abiotic stress factors. The design of new effective microbial consortia is the final objective of the research.

Dietary microbes and Mediterranean diet

Omri, Ghofrane¹; Fragasso, Mariagiovanna¹; Diaferio, Antonietta¹; Riso, Patrizia²; Guglielmetti, Simone³; Spano, Giuseppe¹; Capozzi, Vittorio⁴

¹ Department of Agricultural Sciences, Food, Natural Resources and Engineering, University of Foggia, Foggia, Italy; ² Department of Food, Environmental and Nutritional Sciences (DeFENS), University of Milan, Italy; ³ Department of Biotechnology and Biosciences (BtBs), University of Milano-Bicocca, Milan, Italy; ⁴ National Research Council of Italy Institute of Sciences of Food Production (ISPA), Foggia, Italy

The Mediterranean diet, characterised by high consumption of fruits, vegetables, whole grains, legumes, and olive oil, alongside a moderate intake of fish and poultry, has long been associated with numerous health benefits, including reduced risks of cardiovascular diseases, cancer, and neurodegenerative disorders. Emerging trends in scientific research highlight a contribution of dietary microbes in human health and nutrition. Our research synthesises current evidence on the microbial content of Mediterranean diet components, such as fermented dairy products, olives, vegetables and fruits, in order to define appropriate microbe-depleted and microbe-rich diets for future experimental trials. The research explores the microbiological classification of food items within the Mediterranean diet and examines the abundance and diversity of the related microorganisms. We classified Mediterranean foods based on their microbial profiles using previous information opportunely integrated with original findings using culture-dependent methods. The different food items were categorised according to microbial content: low (Lo), 10 CFU/g. This classification revealed significant variations in terms of microbial abundance, providing the basis for a critical evaluation of this aspect. Subsequently, we established two isocaloric daily diets, both compliant with the guidelines of the Mediterranean diet: one microbe-depleted and one microbe-rich diet. By starting with similar products, we highlighted how food processing and ingredient choices can modify dietary microbe load, contributing to design diets for specific experimental proposes. Our findings emphasize the need for further research to enhance health benefits from microbial intake associated with the Mediterranean diet, offering promising directions for future studies in food science and nutrition.

Acknowledgments

This work was completed as part of the bEat project (We all eat microbes: diet as reservoir of microorganisms that preserve the ecosystem services of the human gastrointestinal microbiota) supported within the call “Progetti di Ricerca di Rilevante Interesse Nazionale (PRIN) 2022” granted by the Italian Ministry for Universities and Research (MUR) (Prot. 2022TF9AHZ, CUP G53D2300574 0006). Vittorio Capozzi was partially funded by CNR project “NUTRAGE FOE2021 DBA.AD005.225”; Mariagiovanna Fragasso is supported by the European Union NextGeneration EU [PNRR, CN00000022] within the Agritech National Research Centre for Agricultural Technologies.

Fungi as bioresources for remediation of HCH-contaminated soils: from microbial community-level physiological profile to selective isolation in enrichment

Ceci, Andrea¹; Riga, Alessandra¹; Spinelli, Veronica¹; Persiani, Anna Maria¹

¹ *Department of Environmental Biology, Sapienza University of Rome, Rome, Italy*

The interaction between human activities and global change (including persistent chemicals pollution) poses severe threats for the soil microbiota thus reducing the provision of ecosystem services¹. In this context -, -, and -hexachlorocyclohexane (HCH) are highly persistent organic pollutants of global concern, and a severe risk for human health and ecosystem functioning. Soil fungi, thanks to the ability to tolerate, bioaccumulate and biodegrade HCH, are important bioresources as biobased solutions for HCH-contaminated soil remediation. The study area was selected within the National Priority Site “Bacino del Fiume Sacco” in the Metropolitan City of Rome (Italy). Soil cores, up to 1 m of depth, were collected from 2 plots and later divided in topsoil (TS: 0-10 cm) and subsoil (SS: 10-100 cm) samples. The first goal was to characterize the microbial community level physiological profile, so the soil samples were analysed by the Biolog EcoPlate™ Technique² to compare metabolic activities of the communities at different depths (TS and SS). Moving on, the project focused on the fungal fraction of the microbial community, evaluating the fungal load differences between TS and SS, through the count of the colony forming units (CFUs/dry soil weight). The CFUs results show a higher fungal load in topsoil than that in subsoil by one order of magnitude. To isolate fungal bioresources suitable for HCH degradation, a selective enrichment procedure with a high concentration HCH mixture as the only carbon source, was carried out. At the end of the procedure several species, mainly belonging to *Fusarium* and *Alternaria* genera, were isolated and are currently preserved in the Culture Collection of the Fungal Biodiversity Laboratory (FBL) of the Department of Environmental Biology of Sapienza University of Rome. The isolated fungi represent useful bioresources for further studies aimed at the development of mycoremediation application for HCH contaminated soil remediation.

References

- Bach et al., 2020 10.3390/su12072662
Decena Jr. & dela Cruz, 2024 10.1007/s12088-024-01294-73
Spini et al., 2018 10.3389/fmicb.2018.02543

Acknowledgments

Funder: Project funded under the National Recovery and Resilience Plan (NRRP), Mission 4 Component 2 Investment 1.4 Call for tender No. 3138 of 16 December 2021, rectified by Decree n.3175 of 18 December 2021 of Italian Ministry of University and Research funded by the European Union – NextGenerationEU; Award Number: Project code CN_00000033, Concession Decree No. 1034 of 17 June 2022 adopted by the Italian Ministry of University and Research, CUPB83C22002950007, Project title “National Biodiversity Future Center NBFC”

From culture collections to biorecovery of strategic elements: exploring fungal interactions with Germanium and Indium

Spinelli, Veronica¹; Pinzari, Flavia²; Muzzini, Valerio Giorgio³; Donati, Enrica²; Iori, Valentina⁴; Astolfi, Maria Luisa⁵; Persiani, Anna Maria¹; Mazzonna, Marco²; Ceci, Andrea¹

¹ Department of Environmental Biology, Sapienza University of Rome, Rome, Italy; ² Institute for Biological Systems National Research Council (CNR-ISB), Montelibretti (Rome), Italy; ³ Research Institute On Terrestrial Ecosystems National Research Council (CNR-IRET), Montelibretti, Rome, Italy; ⁴ Institute of Agricultural Biology and Biotechnology National Research Council (CNR-IBBA), Montelibretti, Rome, Italy; ⁵ Department of Chemistry, Sapienza University of Rome, Piazzale Aldo Moro 5, 00185, Rome, Italy

Fungi in nature play crucial roles in geology and ecology, and represent valuable resources for sustainable recovery processes¹. Fungi can tolerate, bioaccumulate, and transform toxic compounds and several strategic elements, e.g. germanium (Ge) and -indium (In). These are elements of fundamental interest in high-tech products, and their extraction often heavily impacts ecosystems. Their fungal-mediated recovery from electronic wastes offers an eco-friendly solution to mitigate environmental impacts and support sustainable resource management². In this study 24 fungal isolates were selected based on available literature on fungi-element interactions, known metal-fungi mechanisms, and biological and ecological features. These fungi were chosen among those preserved in the culture collections of the Fungal Biodiversity Laboratory at Sapienza University of Rome and CNR mycology laboratories. Since siderophores are crucial in metal-fungi interactions, the selected fungi were screened for their production through the ChromeAzurool-S (CAS) assay³. Moreover, considering the importance of laccase in bioleaching processes and the fact that these elements share chemical features with those playing primary roles in fungal biology, a screening to evaluate the production and release of laccase was carried out through the Guaiacol assay⁴. Finally, fungal tolerance to Ge and In was evaluated using exposure to 5 mM Ge as GeO₂ or 2.4 mM In as In₂O₃. The screenings were carried out on solid Raper Hoffman medium for 14 days at 25°C. The Rt:Rc (%) index, based on colony areas, and the T.I. (%) index, based on dry weights⁵, were calculated to evaluate fungal tolerance. The variation in medium pH was also assessed. Most of the strains were able to produce siderophores, with 7 showing high response, whereas laccase release was only detected in 8, all basidiomycetes. Finally, all fungi were able to grow in all tested conditions, even though several evident Ge/In effects were observed.

References

- Gadd G.M., Rhee Y.J., Stephenson K., Wei Z. (2012) Environ. Microbiol. Rep. 4:270–296.
 Liang X., Gadd G.M. (2017) Microb. Biotechnol. 10:1199–1205.
 Mehnert M., Retamal-Morales G., Schwabe R., Vater S., Heine T., et al. (2017) Sol. St. Phen. 262:509–512.
 Senthivelan T., Kanagaraj J., Panda R.C., Narayani T. (2019) Biotechnol. Rep. 23:e00344.
 Ceci A., Spinelli V., Massimi L., Canepari S., Persiani A.M. (2020) Appl. Sci. 10:3218.

Acknowledgments

PRIN 2022 PNRR Project funded by European Union -NextGenerationEUProject Prot. P2022ENEWL
 Title “Fungal interaction with metals (FUN METALS): transformation and mechanisms for biorecovery” CUP B53D23032130001.

Microbial consortia the great bio-based solution for the bioremediation of co-contaminated soils: study of microbial resources isolated from a decommissioned military site.

Giovannini, Roberto¹; Spinelli, Veronica¹; Ceci, Andrea¹; Bellino, Mariarosa²; Scaffidi, Simona²; Capozzi, Nunzio²; Persiani, Anna Maria¹

¹ Department of Environmental Biology, Sapienza University of Rome, Rome, Italy; ² Centro Tecnico Logistico Interforze NBC, Italian Army, Ministry of Defense, Italy

It has been estimated that around 33% of global soil is degraded¹. Considering estimations of soil loss and of contamination effects exacerbation due to future climate change scenarios, the need for sustainable remediation of contaminated soils is crucial for maintaining the ecosystem services.

This study investigated fungal and bacterial strains, isolated from soils and a *Plantago lanceolata*'s rhizosphere sampled in a decommissioned military site, for their potential in detoxification of co-contaminated soil. Globally 101 fungal taxa and 185 bacterial strains were isolated from the samples and are currently preserved in the Fungal Biodiversity Lab. culture collection. To select the most suitable bioresources, all these isolates were screened in a microwell plate enrichment experiment² for their ability to utilize PAHs as their sole C source also in co-contamination with Pb and/or Zn. 13 fungal species and 15 bacterial strains were able to grow using PAHs as C source also in co-contamination conditions. So, as microbial consortia are reported for their higher bioremediation efficacy³, in-vitro compatibility tests⁴ were performed among the selected strains isolated from the same sampling plot to define the consortia formulations. The 9 resulting consortia and an additional one formulated with 5 *Trichoderma* strains, were tested in microcosm conditions to assess their effectiveness in detoxifying soil polluted with PAHs, PAHs+Pb, PAHs+Zn, PAHs+Pb+Zn. The detoxification efficacy of the biological treatments was evaluated by an array of ecotoxicological assays, whose results revealed that, despite several consortia successfully reduced the toxicity in at least few contamination conditions, consortium D was the most effective in reducing toxicity in the microcosm experiment. Consequently, the consortia D and A are currently being studied in a pot experiment, including also a microbially assisted phytoremediation condition with *P. lanceolata*, to confirm their efficacy.

References

- FAO and ITPS. 2015. ISBN 978-92-5-108960-6
 Zeng, J., Lin, X., Zhang, J., and Li, X. (2010) J. Hazard. Mater. 183, 718–723.
 Massot, F., Bernard, N., Alvarez, L.M.M., Martorell, M.M., Mac Cormack, W.P., and Ruberto, L.A.M. (2022). Appl. Microbiol. Biotechnol. 106, 2283–2297.
 Tabacchioni, S., Passato, S., Ambrosino, P., Huang, L., Caldara, M., Cantale, C., Hett, J., DelFiore, A., Fiore, A., Schlüter, A., et al. (2021). Microorganisms 9, 426.

Funding

This study was funded by the Italian Ministry of Defense under the project “Bioremediation: caratterizzazione di comunità microbiche da suoli contaminati” C.I.G. 8744192F7F

The CABI Biological Resource Collections: evolving to meet the changing needs of users.

Kermode, Anthony¹; Cobb, Emma¹; Stewart, Helen¹; Bonnin, J. Miguel¹; Buddie, Alan G¹; Ryan, Matthew J.¹

¹ CABI, Egham, UK

Established in 1947, the CABI Genetic Resource Collection (GRC) is one of the most significant global collections of microorganisms, containing >28,000 fungi and approximately 2,000 bacteria. The collection is primarily comprised of plant-associated and environmental microorganisms. It contains several smaller collections, including the UK's National Filamentous Fungus Collection, National Collection of Wood-rotting Fungi; the UK Aquatic Phycomycetes Collection, the British Antarctic Survey Collection and others. One benefit of the broad geographic coverage of the collection and the lengthy timeframe over which strains have been deposited, is that strains in the collection may be used to study the evolution of key fungal groups, as well as enabling reappraisals of legacy strains, with new molecular tools, which may not only shed light on 'new' cryptic species. But also allow us to understand the relationships between organism groups. Recent studies have included contributing to the major reappraisal of the genus *Fusarium* (being led currently by the Westerdijk Institute) and a study of genome evolution in the coffee wilt fungus, *Fusarium xylarioides* with Imperial College, London.

Recently, the UKRI-funded UK Crop Microbiome CryoBank has established a collection of 36,000 bacteria and 4,800 environmental soil rhizosphere samples, held within the CABI CryoFacility, this extends the collection's holdings to include environmental and pure cultures from several key crops. This builds on CABI's expertise in cryopreservation science and provides links with EU infrastructure projects such as the EU Microbiome (RI) Enabler and EU Microbes4Climate and together enables both a better understanding of the rapidly developing 'microbiome' research area, but crucially supports the broader research community. Such developments are clearly demonstrating the continued relevance of the CABI Collections to cutting-edge science.

References

- Smith D, Kermode A, Cafà G, Buddie AG, Caine TS & Ryan MJ (2020) Strengthening mycology research through coordinated access to microbial culture collection strains. *CABI Agriculture and Bioscience* 1: 2 <https://doi.org/10.1186/s43170-020-00004-9>
- Ryan MJ, Mauchline TH, Malone JG et al. The UK Crop Microbiome Cryobank: a utility and model for supporting Phytobiomes research. *CABI Agriculture & Bioscience* 4, 53 (2023). <https://doi.org/10.1186/s43170-023-00190-2>
- Ryan MJ, Peck L, Smith D, Flood J & Barraclough T (2022) Culture Collections as a source of historic strains for genomic studies in plant pathology *Journal of Plant Pathology* <https://doi.org/10.1007/s42161022-01217-9>

Exploring methodological aspects of microbiota transplantation in black elder (*Sambucus nigra*)

Rimša, Antonija¹; Gudra, Dita²; Balode, Laura¹; Boroduške, Anete¹

¹ Department of Microbiology and Biotechnology, University of Latvia, Riga, Latvia; ² Latvian Biomedical Research and Study Centre, Riga, Latvia

Plant microbiome studies have previously revealed an impact of endophytic microbial communities on complex plant traits such as disease resistance, phytochemical content, or yield. However, targeted manipulation of plant microbiota to improve plant properties is yet to be achieved. Here we explore the idea of microbiota transplantation in wild black elder (*Sambucus nigra*) populations in Latvia. We attempt to develop transplantation methods by identifying methodological aspects significantly affecting the preparation and storage of microbiota inoculums. For this research, we explored three methodological aspects – we analysed the effect of fractionation of plant slurry used for microbiota transplantation on the composition of the microbial community, we conducted a long-term storage experiment to see if and how the microbiome inocula changed after storing in the freezer for longer periods, and lastly, we tested two different microbiota application techniques

– injecting with a syringe and spraying.

Wild samples of unripe black elderberries were collected and surface-sterilized in the lab. Samples were disintegrated, divided into sub-fractions based on particle size, and stored at -80°C. For further analysis, total DNA was extracted, and microbial communities were characterised using target loci amplicon next-gen sequenced. We used the sequence data of ITS and 16S (v1 – v6 regions) ribosomal RNA gene sequences to characterize the endophytic microbiome.

The preliminary results show a slightly increased bacterial diversity in plant slurry samples composed of particles ranging in size from 57 to 165 µm. The proportions of the main bacterial genera found in filtered samples also differ between larger-sized particles (57-165 µm) and smaller-sized particles (0-57 µm). Our research prospects are to optimize and implement in practice plant microbiota transplantation methods, which could potentially be applied in other economically important plant species in the future.

Acknowledgements

The research was supported by the Latvia Council of Science Project No. lzp-2022/1-0179 “Role of endophytic microbiota in the anti-viral activity of *Sambucus nigra* L. against type A influenza and SARS-CoV-2.”

Plant growth promotion endophytes from sicilian native grapevines as plant biostimulants for climate-ready crops

Crucitti, Dalila²; Navarro-Torre, Salvadora³; Carimi, Francesco¹; Zottini, Michela⁴; Pacifico, Davide¹

¹ *Institute of Biosciences and Bioresources (IBBR), CNR, Palermo, Italy;* ² *Department of Agricultural, Food and Forest Sciences (SAAF), University of Palermo, Palermo, Italy;* ³ *Department of Microbiology and Parasitology, Faculty of Pharmacy, University of Seville, Seville, Spain;* ⁴ *Department of Biology, University of Padova, Padova, Italy*

Mediterranean crops are undermining by increasingly extreme weather events in particular heat waves, drought and intense rainfall. Water scarcity has become the most limiting factor of agriculture in the Mediterranean regions and the expansion of arid areas is threatening traditional productions, such as viticulture. To improve the resilience and sustainability of vineyards, the endophytic microbiota of the Phyllosphere was studied in native grapevine cultivars adapted to arid conditions, and in wild grapevines from Sicily.

Endophytic communities were isolated and identified by sequencing of 16S rDNA region. The most representative isolates, belonging to species already known for their Plant Growth Promoting (PGP) properties were characterized by in vitro assays. The tested PGP properties included indole-3-acetic acid and siderophore production, phosphate solubilization, nitrogen fixation, biofilm formation, the presence of 1-aminocyclopropane-1-carboxylic acid deaminase gene and of enzymatic activities. Among the 13 PGP activities tested, more than 70% of the strains exhibited five to seven PGP traits concurrently. In vitro tests showed that most of the isolates were able to stimulate plant nutrition and growth, through biofilm formation, amylase and lipase activity, siderophore production and nitrogen fixation. Two bacterial consortia combining three distinct strains were selected to cover the greatest number of PGP properties. The potential role of each consortium in promoting plant growth and drought tolerance was tested on an indicator plant (*Medicago sativa*) both in vitro using drought resistance tests and in pots under controlled greenhouse conditions. Plant physiological status was determined by measuring plant growth and photosynthetic parameters. The use of native microbial consortia is becoming a promising technique to reduce the harmful impacts of chemical fertilizers and contribute to a more sustainable agricultural system, to cope with climate change.

Exploring sicilian olive tree eco-friendly endophytes for sustainable agriculture

Crucitti, Dalila¹; Navarro-Torre, Salvadora³; Carimi, Francesco²; Caruso, Tiziano¹; Pacifico, Davide²

¹ Department of Agricultural, Food and Forest Sciences (SAAF), University of Palermo, Palermo, Italy; ² Institute of Biosciences and Bioresources (IBBR), CNR, Palermo, Italy; ³ Department of Microbiology and Parasitology, Faculty of Pharmacy, University of Seville, Seville, Spain

Olea europaea L. is one of the most important crops that characterize the Mediterranean agriculture. Olive trees can withstand various environmental stresses such as heat, drought, salinity, and high levels of UVB, by implementing a variety of adaptations strategies including the morphological changes and the activation of physiological and biochemical mechanisms that make it a resilient woody crop. Olive tree microbiota comprises endophytes, microorganisms that promote plant growth and stress tolerance. To develop a microbe-assisted improvement of production, and agroecosystems resilience, the Phyllosphere bacterial communities in three olive cultivars and wild olive trees from Sicily were investigated over four phenological stages. The isolated and molecular identified microorganisms were characterized to assess the presence of plant growth promoting (PGP) traits. The tested PGP properties included indole-3-acetic acid and siderophore production, phosphate solubilization, nitrogen fixation, biofilm formation, the presence of 1-aminocyclopropane-1-carboxylic acid deaminase gene and of enzymatic activities (DNase, amylase, cellulase, lipase, chitinase, protease and pectinase). All isolates showed at least one of the screened PGP properties and more than 60 % of them exhibited from five to ten PGP traits concurrently. The most common PGP properties were siderophore production, biofilm formation and nitrogen fixation. Based on PGP activities, two different endophytic consortia were formed and tested for in vitro compatibility and drought assay. The potential role of bacterial consortia to promote the growth and the tolerance to drought of an indicator plant (*Medicago sativa*) was assessed in pots in controlled greenhouse conditions. Plant growth and photosynthetic parameters were measured to determine the plant physiology status. Applications of native microbial consortia could be thought for a sustainable agriculture capable of facing the new climate scenarios.

Fungal diversity in the precious Mediterranean red coral *Corallium rubrum*

Prioux, Camille^{1,2,3}; Carrasco-Acosta, Marina^{4,5}; Prigione, Valeria⁵; Tignat-Perrier, Romie²; Allemand, Denis³; Ferrier-Pages, Christine³; Varese, Giovanna Cristina⁵

¹ Sorbonne Université, Collège Doctoral, Paris, France; ² Unité de Recherche sur la Biologie des Coraux Précieux CSM CHANEL, Centre Scientifique de Monaco, 8 Quai Antoine 1er, MC 98000, Monaco, Principality of Monaco; ³ Centre Scientifique de Monaco, 8 Quai Antoine 1er, MC 98000, Principality of Monaco; ⁴ Department of Biology, Instituto Universitario de Investigación en Estudios Ambientales y Recursos Naturales i-UNAT, University of Las Palmas de Gran Canaria, Las Palmas, Spain; ⁵ Mycotheca Universitatis Taurinensis, Department of Life Sciences and Systems Biology, University of Torino, Turin, Italy

Corals maintain a symbiotic relationship with a variety of microorganisms, including bacteria and microeukaryotes, which inhabit their tissues and provide important nutrients. These microeukaryotes include various groups of protozoa, microalgae, and fungi. However, fungi, an often overlooked but integral component of these symbiotic communities, may play a critical role in coral health and resilience to stress. In this work, we have characterized the diversity of cultivable and non-cultivable fungi associated with the precious red coral *Corallium rubrum* that together with other gorgonians, is part of the Marine Animal Forests (MAFs) harbouring an incredible biodiversity in the Mediterranean Sea. For this purpose, we collected 6 coral nubbins at 40m depth in Villefranche-sur-mer, (Ligurian Sea). By coupling culture isolation and metabarcoding techniques, and separating the host tissue from the skeleton, we observed a large fungal biodiversity. This comparative approach showed that the combination of dependent and independent-culture methods is essential to reduce bias and obtain a more accurate representation of the fungal community. Our findings underscore the importance of understanding the diversity and ecological roles of microeukaryotes. This is critical for coral conservation and management, especially in the face of anthropogenic threats and climate change.

A network-based approach to elucidate the dynamics of the native microbiota's response to a multi-kingdom microbial inoculant in raw brewer's spent grain

Zara, Giacomo^{1,3}; Bianco, Angela^{1,3}; Assandri, Davide²; Cossu, Carla¹; Budroni, Marilena^{1,3}

¹ *Department of Agricultural Sciences, University of Sassari, Sassari, Italy;* ² *Institute of Sciences and Technologies for Sustainable Energy and Mobility (STEMS), National Research;* ³ *Associated member of the JRU MIRRI-IT, Italy*

Identifying the key factors that influence microbiota stability and resilience is of uttermost importance for effectively managing microbial resources in various environmental applications [1]. Indeed, microbial inoculants, typically chosen under controlled conditions, often struggle to survive in the field due to factors like soil type, nutrient levels, environmental stress, and competition with native microbiota [2,3]. Recently, the interactions between a multi-kingdom microbial inoculant and the native microbial population of untreated brewers' spent grains (BSG) were assessed [4,5]. Specifically, inoculated and spontaneous solid-state fermentation (SSF) processes were monitored for 90 days by analyzing BSG chemical and biochemical properties as well as the evolution of the microbiota through high-throughput sequencing of 16S and ITS rRNA. Starting on day 14, the bacterial and fungal genera present in the bioinoculant were no longer able to survive in the BSG and were outcompeted by the native microbiota. Nonetheless, the bioinoculant had a significant impact on the chemical, biochemical and fertilizing properties of BSG. Cross-kingdom co-occurrence network analysis suggested that this impact was associated with the reshaping of the native microbial population in the inoculated samples. Indeed, the network from the inoculated samples was more connected and stable due to stronger and more significant correlations. Interestingly, most of these interactions were positive, suggesting mutualistic relationships between microbial taxa. These findings suggest that comprehending the interplay between native and external microbial communities in complex ecosystems, and employing microbial inoculants tailored to specific environments or conditions, facilitates the creation of effective bio-based applications. Consequently, the microbial strains with agronomic significance, derived from the native microbiota of BSG, have been archived in the MBDS-UNISSCC culture collection.

References

- [1] French, et al. Emerging strategies for precision microbiome management in diverse agroecosystems. *Nat. Plants* 7, 256–267 (2021).
- [2] Zhang, et al. Arbuscular mycorrhizal fungi increase grain yields: a meta-analysis. *N. Phytol.* 222, 543–555 (2019).
- [3] Bianco, et al. Microbial community assembly and chemical dynamics of raw brewers' spent grain during inoculated and spontaneous solid-state fermentation. *Waste Manag.* 174, 518–527 (2024)
- [4] Bianco, et al. The potential use of brewers' spent grain-based substrates as horticultural bio-fertilizers. *Frontiers in Sustainable Food Systems*, 8, 1404914. (2024).

Exploring the oenological characteristics of different non-*Saccharomyces* species in grape must fermentation.

Costantini, Antonella¹; Pulcini, Laura¹; Tsolakis, Christos¹; Vaudano, Enrico¹

¹ CREA-Centro di Ricerca Viticoltura ed Enologia, Asti, Italy

Complex biochemical processes occur in wine fermentations, where yeasts, specifically *Saccharomyces cerevisiae*, convert sugars to ethanol, carbon dioxide, and other volatile compounds. Nowadays, the role of non-*Saccharomyces* yeasts in winemaking is gaining significant interest due to their ability to enhance the aromatic complexity and sensory profile of wines.

In this work, thirty-five yeast species from the CREA-Collection of Microorganisms of the Viticultural and oenological Environment (CMVE), conserved at CREA-VE in Asti, were tested. Yeasts were inoculated in wine must, and fermentations were monitored by weight loss. At the end of alcoholic fermentation, the following parameters were analysed: alcoholic degree, residual sugars, malic acid, lactic acid, volatile acidity, and glycerol.

The results showed that only three species completed alcoholic fermentation, with residual sugars less than 1 g/L: *Saccharomycodes ludwigii*, *Schizosaccharomyces pombe*, and *Zygosaccharomyces bisporus*. Regarding the volatile acidity, most of the tested species produced below 1 g/l of acetic acid. Some yeasts, in particular *Hanseniaspora osmophila*, *Hanseniaspora valbyensis*, *Hanseniaspora uvarum*, *Zygosaccharomyces rouxii*, *Zygosaccharomyces bailii*, *Pichia terricola*, and the already-known *Schizosaccharomyces pombe*, could reduce malic acid.

Interesting data were found for *Kazachstania exigua*, which exhibited high glycerol production and high consumption of the sugars present in the must.

This study underlines the importance of discovering the distinctive characteristics of the yeast species. This could facilitate to find interesting yeasts that are currently not employed in wine fermentation. Non-*Saccharomyces* species as starter cultures are becoming an increasingly important approach in modern winemaking; they can contribute to producing specific aromas or be used to make wines with reduced ethanol concentrations.

Harnessing Microbiomes for Drought-Resilient Agriculture: Insights from the BIOMEnext Project

Visca, Andrea¹; Bevivino, Annamaria¹; Costanzo, Manuela¹; Di Gregorio, Luciana¹; Perrotta, Gaetano¹

¹ ENEA, Italian National Agency for New Technologies, Energy and Sustainable Economic Development, Department for Sustainability, Rome, Italy

The global increase in water scarcity presents a major challenge to sustaining crop productivity. Lack of adequate water results in the degradation of the photosynthetic apparatus, interruptions in essential metabolic processes, increased production of free radicals, and compromised plant root structures. Drought is a key stress factor that directly affects the osmotic balance of plant cells. Within the BIOMEnext Project, innovative, composite, and eco-friendly farming systems have been evaluated to improve the resilience of Mediterranean fruit farming to climate change. Specifically, metagenomic analyses were conducted, characterizing the core rhizosphere and endophyte microbiomes and their predicted functions in four olive varieties (Arbequina, Chemlal, Koroneiki, and Shengeh) under wet and dry conditions during a one-year field experiment. DNA from both root endophytes and soil rhizosphere was extracted and sequenced using Oxford Nanopore technology, producing long reads of the 16S rRNA gene. These long reads enabled species-level identification of the microbial composition using the Emu tool. Additionally, a novel tool was developed for functional annotation using PICRUSt2 with long reads. A focus was placed on functions related to cold-heat stress, general stress and heat-shock, which are all functions often associated with drought resistance. Finally, PGP functions were also included in the research. This research allowed the identification of genera associated with these functions which, although not differing much at the functional level between the rhizosphere and roots, are completely different at the genus level. In fact, two core microbiomes have been defined, one for rhizospheric soils and one for roots, which are potentially involved in drought resistance.

Acknowledgements

This project has received funding from the European Union (project BIOMEnext) under the Grant Agreement no. 101102316

Harnessing Fungi for Eco-Friendly Biosurfactant Production: joint efforts from CECT and MUT collections to unveil this potential among the preserved strains

Florio Furno, Matteo¹; Gumiel, Marta²; Spina, Federica¹; Laera, Angelica¹; Perugini, Iolanda¹; LopezOcaña, Laura²; Varese, Giovanna Cristina¹; Aznar, Rosa²

¹ University of Turin, Turin, Italy; ² University of Valencia, Valencia, Spain

Biosurfactants are gaining significant attention in the global market as natural additives for commercial products and environmental applications due to their ability to reduce surface and interfacial tension between fluid phases. Compared to chemical surfactants, biosurfactants offer enhanced stability across a range of physico-chemical conditions and are environmentally friendly. The biotechnological production of biosurfactants is rapidly evolving, with fungi emerging as highly promising producers of these valuable biomolecules.

In this context, the aim of this work is to explore the potential of fungal biosurfactant production through various screening tests on 20 yeast strains. These strains were selected from the most representative genera deposited in the Spanish Type Culture Collection (CECT) and Mycotheca Universitatis Taurinensis (MUT) mainly isolated from wastewater sediments of olive oil mills.

The screening involved: the oil displacement test and the drop collapsing assay. The fungi were pre-grown in 100 mL flasks containing 40 mL of modified mineral salt medium (MSM) with olive oil (50 mL/L) as the sole carbon source to stimulate surfactant production. The flasks were incubated at 24 °C with agitation at 120 rpm for 21 days. The capability to produce biosurfactants was tested at four time points (T3, T7, T14, and T21).

The results indicate that four strains belonging to the species *Wickerhamomyces anomalus* were the best performers in both screening tests revealing their capability to produce strong biosurfactants and powerful emulsification activities. Ongoing analyses are trying to elucidate the metabolites and the metabolic pathways involved. However, preliminary results based on Phenotype Microarray (PM, Omnilog) highlight the activation of similar metabolic pathways between the four strains of *W. anomalus*, which are distinct from those of the other strains tested.

Biotechnological production of biosurfactants from fungi represents a promising, simple, and cost-effective method for large-scale production, potentially replacing synthetic surfactants in various industries. Further works are necessary to identify the secreted and to scale-up their production.

Acknowledgements

Part of this work was granted by the European Commission—NextGenerationEU, Project “Strengthening the MIRRI Italian Research Infrastructure for Sustainable Bioscience and Bioeconomy”, code n. IRO000005. Marta Gumiel is recipient of INVESTIGO and ERASMUS+ grants.

Changes in bacterial, archaeal, and fungal communities from saline soils cultivated with tomatoes in Italy

Prigigallo, Maria Isabella¹; Pirra, Sara¹; Picciotti, Ugo¹; Bubici, Giovanni¹

¹ CNR, Istituto per la Protezione Sostenibile delle Piante, Bari, Italy

Soil salinization is a major concern in modern agriculture. In southern Italy, the climate change scenario is challenging tomato crops, especially because of increased average temperature, irregular precipitation patterns, and scarcity of irrigation fresh water. Apulia ranks among the top Italian regions producing processing tomatoes, but certain areas are being subjected to salinization, especially near the coast. A metagenomic study was conducted to compare the microbiome from saline and non-saline soils in the Apulia region. Three areas, each including four fields with three replicated samples, were sampled at three time points during the tomato cropping cycle, viz. post transplanting, mid-season, and near-harvest time, for a total of 108 soil samples. Physical-chemical analysis was carried out to characterize the soils, and DNA was extracted for Illumina sequencing of amplicons specific to bacteria, Archaea, and fungi.

Saline and non-saline soils differed for exchangeable sodium, limestone, cation exchange capacity, and some macro and micro-nutrients, whereas no significant difference occurred in the texture. Electrical conductivity increased during the cropping cycle in fields irrigated with saline water, suggesting that soil salinization depended on irrigation water quality rather than pedoclimatic conditions.

The amount of DNA extracted from soil samples, as well as the number of sequencing reads, were unexpectedly low at post-transplanting compared to the other two time points, indicating that the establishment of microbiomes in the rhizosphere took place as the crop had developed. In the three considered kingdoms, i.e., Bacteria, Archaea, and Fungi, differential operational taxonomic units (OTUs) were identified in saline and non-saline soils, thus taxa with a better fitness under high salinity were identified. This paves the way for searching for new plant-beneficial microbes sustaining tomato crops cultivated in salinized soils.

Acknowledgements

This work was funded by the research project “Deciphering microbiomes from extreme environments for the improvement of crop resilience to climate change, food security and safety (MiCroResi)” (framework “4th EU-LAC Multi-thematic Joint Call for collaborative projects from Europe, Latin-America and the Caribbean Countries”) and by “Agritech National Research Center and received funding from the European Union Next-GenerationEU” (PIANO NAZIONALE DI RIPRESA E RESILIENZA (PNRR), MISSIONE 4 COMPONENTE 2 INVESTIMENTO 1.4, D.D. 1032 17/6/2022, CN00000022). This manuscript reflects only the authors' views and opinions, neither the EU nor the European Commission can be considered responsible for them.

A polycyclic aromatic hydrocarbons (PAHs) contaminated soil as source of microorganisms for bioremediation purposes

Poli, Anna^{1,2}; Crespi, Matteo¹; Marchitelli, Andrea Lara¹; Perugini, Iolanda¹; Varese, Giovanna Cristina^{1,2}; Prigione, Valeria^{1,2}

¹ *Department of Life Sciences and Systems Biology, University of Torino, Torino, Italy;* ² *NBFC, National Biodiversity Future Center, Italy*

The soil ecosystem is variable and complex and hosts a huge variety of microorganisms, including fungi that provide several ecosystem services.

Due to the increase of industrial activities and to the combustion of fossil fuels and organic matter, the release of PAHs threatens the health of natural ecosystems. Soil contamination by PAHs is an ever more severe issue that requires a sustainable solution. A wide range of fungi is capable of degrading these recalcitrant compounds through their enzymatic arsenal. Thus, investigating and identifying the cultivable mycobiota inhabiting PAH-contaminated soils is crucial for designing appropriate remediation approaches.

In this work, we investigated the fungal diversity of a formerly cultivated urban garden in Torino (Italy), mainly contaminated by benzo(a)pyrene, benzo(g,h,i)perylene, fluoranthene, and phenanthrene.

Overall, 181 fungal isolates, mostly belonging to Ascomycota were retrieved and were affiliated to 38 genera and 66 species, including putative novel species affiliated to genus *Coniochaeta*. In parallel, a culture independent approach based on the sequencing of ITS2 region, applied to investigate the fungal diversity, confirmed the dominance of Ascomycota (99%), while Basidiomycota and Mortierellomycota accounted for less than 1%.

Following, to select those strains capable of degrading PAHs, enrichment assays were conducted with the target pollutants as sole carbon source. More than 99% of the 104 isolates retrieved belonged to Ascomycota, while only one Basidiomycota was isolated in axenic culture. The selection applied allowed the development of a number of species (61%) that were not detected in the bulk soil.

The fungi derived from the enriched cultures are currently objects of deep investigation whose outcome will serve to evaluate their degradative potential and to develop remediation strategies for in situ applications.

Acknowledgments

Bando BRiC – edizione 2022 – ID:65 Tecniche avanzate di micorimedio: una soluzione sostenibile per la bonifica di suoli contaminati e per la sicurezza per gli operatori CUP: D13C22003330005

Impact of newly developed biofertilizer “polyazorich” on soil agrochemical properties during crop cultivation

Harutyunyan, Seda¹; Stepanyan, Tamara¹; Khachatryan, Gayane¹; Karganyan, Arthur²; Bagiyani, Valeri¹; Goginyan, Vigen¹

¹ Scientific and Production Center “Armbiotechnology” NAS RA; ² «BIONA» Limited Liability Company, Russia

The development of methods for restoring the natural balance of soil microflora, the use of different preparative complexes separately and in various compositions are of great importance for agricultural farming in order to revive degraded soils. The use of preparations on the basis of microorganisms is the most effective way to increase the productivity of plants, which makes it possible to preserve the soil fertility and the ecological balance of soils.

The new biofertilizer contains association of most promising domestic free-living (*Azotobacter chroococcum*), symbiotic nitrogen-fixing (*Rhizobium* spp., *Mesorhizobium ciceri*, *Bradyrhizobium japonicum*, *B. arachidis*) and phosphate-solubilizing (*Paenibacillus polymyxa*) bacteria, which are able to convert soil nutrients into a soluble form available for absorption by plants, increase yields, have fungicidal properties.

Agricultural crops were grown in light loamy soil of brown semi-desert type, characterized by a relatively low humus content (2.73%), salinity (0.092%), concentration of carbonates (0.67 mgEq/100g) and sulfates (0.07 mg-Eq/100g), indicators of basic soil nutrients: mobile form of nitrogen (NO_3^-) 30.2, immobile and mobile forms of phosphorus (P_2O_5 , PO_4^{3-}) – 10.4, 14.0, respectively and immobile and mobile forms of potassium (K^+) 14.7 mg-Eq/100g.

After biopreparation impact, the organic nutrients content in the soil reached 5.79% within 60 days vegetation. At the same time, there is an increase in NO_3^- to 79.1 mg-eq/100g, as well as saturation of the soil with N_2 up to 16.2 mg/100g. Meanwhile, a decrease is observed in the content of phosphorus salts P_2O_5 , PO_4^{3-} and K^+ to 2.86, 3.82 and 5.86 mg-Eq/100g, respectively.

Acknowledgements

This work was supported by the Higher Education and Science Committee of RA (Research project № 23EDP-4D003).

Session 2 – Microbes from farm to fork

Wednesday 18th September, 5:00pm

Chairs: De Vero Luciana (Italy), Oivanen Pekka (Finland).

Using microbial diversity or complex communities for valorization of side-streams for food applications

Xiao, Hang ¹; Todorov, Sanne Kjærulf ¹; Bang-Berthelsen, Claus Heiner ¹

¹ Technical University of Denmark, Denmark

The global food system significantly impacts the biodiversity and climate, with bovine dairy and meat production being the primary contributors. Today, many European consumers, especially the younger generation, are keen to reduce greenhouse gas emissions by adopting healthier and more sustainable dietary choices, including reduced meat consumption. Reflecting this global trend towards lower animal protein intake, a majority of consumers, now identifying themselves as flexitarians, are open to decreasing their consumption of animal-based products. Currently, market-available starter cultures are optimized for bovine dairy, highlighting a pressing need for improved starter cultures that can enhance the nutritional value of plant-based dairy alternatives. In this context, we present a novel droplet microfluidics community-level high-throughput screening method to isolate lactic acid bacteria (LAB) strains capable of producing riboflavin in soy-based substrates. This technique has the potential to fortify fermented dairy alternatives with riboflavin. Additionally, recent studies have shown the effects of co-culturing LAB with edible basidiomycetes. This investigation explores the interactions between two LAB strains and their influence on the mycelium of two edible fungal species, *Wolfiporia cocos* and *Laetiporus sulphureus*. Notably, changes in mycelial biomass, color, and morphology were observed, suggesting that co-culturing can functionally modify mycelium for potential food applications. These findings underscore the potential impact of novel microbial diversity and co-culturing on various plant-based food industry applications.

References

- Sanne Kjærulf Todorov et al 2024. Interactions between two strains of lactic acid bacteria and *Laetiporus sulphureus* strain FH24 and FH319, and *Wolfiporia cocos* strain FH9 mycelium, LWT, 2024.
- Hang Xiao et al. 2024 submitted. Isolation, adaptation and screening for riboflavin-producing lactic acid bacteria in plant-based fermentation using droplet microfluidics, Microbiological Research, 2024.

Acknowledgements

This work was funded by Innomission 3 partnership AgriFoodTure (grant No: 1152-00001B, Innovation Fund Denmark) and by the GUDP grant mycoPROTEIN. Additional funding for Hang Xiao was received from Novonesis. The biodiversity included in the abstract is from DTU National Food Institute Culture Collection (NFICC) (dtu.dk).

Are isolation, identification and preservation of microbial strains still useful for food applications?

Baruzzi, Federico¹

¹ *Institute of Sciences of Food Production, National Research Council of Italy (CNR-ISPA), Bari, Italy.*

In the last 20 years, knowledge and understanding of complex microbial populations have boosted up thanks to culture-independent high-throughput sequencing (HTS) techniques. The microbiota of a huge number of foods, related environments and processes have been analyzed improving the definition of microbial complexity of samples that is not possible to achieve with traditional microbiological investigation approaches.

For example, in milk and cheeses, several studies confirm that their microbiomes are usually dominated by lactic acid bacteria. However, supplementary information and data report the presence of unexpected, or new, microbial species.

Thus, the question in the title is the daily challenge for food microbiologists.

In the last three years, over than 140 isolates, mainly belonging to the species *Streptococcus thermophilus* and *Limosilactobacillus fermentum*, have been collected from raw milk, natural whey cultures, fresh curd, brined cheese and different cheeses during ripening or cold storage. In particular, 8 potentially probiotic *Lactococcus lactis* strains and 10 *Lactocaseibacillus paracasei* strains, were isolated by specific protocols from raw milk and Caciocavallo cheese samples, respectively.

In addition to lactic acid bacteria, responsible for milk fermentation and cheese maturation, many other non-dairy species, such as *Klebsiella oxytoca* or *Raoultella terrigena*, were identified.

Thanks to these efforts, 98 bacterial strains were isolated, identified and characterized for their technological properties allowing us to define 4 new dairy processing.

Since these cheeses were realized at dairy plant, at pilot scale level, and will be on the market soon, we could conclude that the improvement of fermented foods can't help but avoid to carry out isolation and characterization of microorganisms from foods and food environments. Overturning the question, are HTS studies really useful for food applications?

Acknowledgements

This study was supported by Italian Ministry of Economic Development within the project "L'evoluzione delle produzioni lattiero-casearie: le biotecnologie valorizzano la tradizione" – ELEVATO n. F/200112/03/X45.

As Scientific Responsible of the project, the Author thanks the efforts made by all CNR-ISPA colleagues allowing to achieve project results here briefly reported.

Soil Health and Agri-Food System Sustainability from a microbiology perspective: A Data-Driven Approach for Agricultural Policy and Practices

Nolfi, Lorenzo¹; Bindo, Arianna²; Di Gregorio, Luciana³; Costanzo, Manuela³; Bernini, Roberta¹; Varese, Giovanna Cristina⁵; Tabacchioni, Silvia³; Palojärvi, Ansa⁶; Manikas, Ioannis⁷; Bevivino, Annamaria³

¹ ENEA, Dept. Sustainability and University of Tuscia, Viterbo, Italy; ² ENEA, Dept. Sustainability, and University of Turin, Turin, Italy; ³ ENEA, Department for Sustainability, Rome, Italy; ⁴ Dept. Agriculture and Forest Sciences, University of Tuscia, Viterbo, Italy; ⁵ Dept. Agricultural, Forest and Food Sciences, University of Turin, Italy; ⁶ Natural Resources Institute Finland(Luke), Turku, Finland; ⁷ Czech University of Life Sciences, Prague, Czech Republic

To guarantee the production of healthy and high-quality food, it is crucial to maintain healthy soil, which serves as the foundational bedrock for robust and sustainable agricultural and food production systems. This research explores the vital role of microorganisms in supporting soil ecosystem functions and the broader agri-food chain. Utilizing advanced bibliometric analysis, the study investigates how microbial biomass, enzymatic activity, and community diversity contribute to agricultural sustainability. It emphasizes the importance of microbiological indicators in managing soil health, that is crucial for enhancing the quality and safety of food production from farm to fork. The analysis aims to bridge scientific knowledge on soil microorganisms with practical agricultural applications, assessing how microbial data can inform and improve agricultural policies and practices to ensure sustainable farming and food safety. The study identifies key research trends, gaps, and emerging themes within the scientific community, providing insights into the integration of microbial research into agricultural policy development. Furthermore, it discusses the connections between scientific findings and their practical applications in policy and practice. In conclusion, this study underscores the strategic importance of scientifically validated microbiological indicators in soil management and their key role in the entire food chain, supporting a solid and sustainable 'farm to fork' framework.

Acknowledgements

This research was supported by ECO-READY (GA N°101084201 <https://www.eco-ready.eu>) and DELISOIL (GA N°101112855 <https://delisoil.eu>) projects, funded by the European Union under the Horizon Europe Program. The authors gratefully acknowledge funding from the Italian project Creazione di un HUB italiano a supporto della partecipazione dell'Italia alla Global Soil Partnership ed alla rete di eccellenza europea sulla ricerca sul suolo-SOIL-HUB, granted by the Italian Ministry of Agricultural, Food and Forestry Policies MIPAAF (DM 37072 28/12/2018) CUP C52F18000200006, and the European Union's Horizon 2020 research and innovation programme under GA N° . 652615 (European Joint Programme SOIL).

Be a QPS or not to be, that is the Quirky Paradox of Safety

Chessa, Luigi¹; Daga, Elisabetta¹; Paba, Antonio¹; Comunian, Roberta¹

¹ *Agris Sardegna, Italy*

Currently, only microorganisms' species included in the Qualified Presumption of Safety (QPS) list, issued by the European Food Safety Authority (EFSA), based on a generic pre-assessment of security on the available literature and updated every six months, can be intentionally added to food or feed. For lactic acid bacteria (LAB) species not recommended for QPS status due to safety concerns, tests ensuring the absence of antibiotic resistance genes (ARGs) and virulence factors are required before their use in food production (e.g. *Enterococcus faecium*), while for other LAB no specific guidelines are available (e.g. *Streptococcus oralis* and *Streptococcus gallolyticus*). Due to the biodiversity loss in raw matrices and production environments, caused by increasingly stringent hygiene practices applied in manufacturing, starters, mainly LAB, are added to the raw material to be processed to easily and safely carry out the fermentation. Especially natural starters, complex microbial communities having a strain composition mostly undefined and obtained from raw matrices without any microbial selection, are used in artisanal and PDO production of the most typical and high-quality agri-food products, also preserving the microbial diversity of the production environment. In a recent study, a natural starter obtained from raw ewe milk, without heat treatment application or pro-technological microorganisms selection, included some strains belonging to non-QPS species. Following the current EFSA guidelines, though the strains resulted free from ARGs and virulence determinants, the culture should not be used for food production. Similarly, artisanal manufacturers may not be allowed to produce/use their own natural starter cultures, likely including non-QPS species. An answer to the paradox of non-QPS microbial consortia that are inherently present in raw matrices and natural starter cultures, but do not comply with the EFSA's rules, is needed to use them for food production.

References

- Chessa L. et al., 2023. A Strategy for the Recovery of Raw Ewe's Milk Microbiodiversity to Develop Natural Starter Cultures for Traditional Foods. doi:10.3390/microorganisms11040823.
- EFSA BIOHAZ Panel, 2023. Update of the list of qualified presumption of safety (QPS) recommended microorganisms intentionally added to food or feed as notified to EFSA. doi:10.2903/j.efsa.2023.7747.
- Hill D. et al., 2020. Extensive bacteriocin gene shuffling in the *Streptococcus bovis*/*Streptococcus equinus* complex reveals gallocin D with activity against vancomycin resistant enterococci. doi:10.1038/s41598020-70328-z.

Investigation on the microbial evolution of cow milk in the passage from stable to mountain pasture, and evaluation of Bitto cheese microbial community

Zago, Miriam¹; Bonvini, Barbara¹; Rossetti, Lia¹; Povoletto, Milena¹; Cabassi, Giovanni¹

¹ CREA-Research Centre for Animal Production and Aquaculture, Italy

Summer transhumance of dairy cows is a seasonal pastoral system practiced in many European countries from ancient times. The microbial composition of milk from three groups of cows, moving from stable in plain farms (PF) to alpine pasture in upper Valchiavenna region (Italy) during the summer period (AF) was studied. The AF milk is used to obtain Bitto, a PDO raw milk cheese produced only in summer, when the cows graze on high alpine meadows. Microbiological and metataxonomic analyses were performed to investigate the microbial evolution of 23 milk samples to determine whether the alpine pasture could influence the milk microbiota. Moreover, the microbiota of six Bitto PDO cheese was also studied. The enumeration of lactic acid bacteria (LAB) showed an overall prevalence of mesophilic cocci and of thermophilic cocci and lactobacilli. Within these groups, *Lactococcus lactis* and *Streptococcus thermophilus* were the most abundant, as shown by metataxonomic analysis. A higher total bacterial count in milk from cows in PF compared to those in AF was also detected. Besides, a higher richness of species in AF milk samples compared to PF was observed, showing differences between the microbiota of PF and AF milk samples. A high relative abundance of *Rhodococcus fascians*, a species linked to vegetations, was observed in AF cow milk, while a high abundance of *Acinetobacter johnsonii* was detected in PF milk. The microbiota of Bitto cheese showed a prevalence of thermophilic LAB species and enterococci. Specifically, a high abundance of *S. thermophilus*, *Enterococcus faecium* and *Lactobacillus paracasei* was highlighted. This study showed that alpine pasture has a strong impact on the microbial composition of cow milk, significantly increasing the abundance of some bacterial groups, specifically thermophilic LAB, recognized as natural starter culture, as also detected in Bitto cheese microbiota.

Influence of the cheesemaking process on the bioprotection activity of *Lactococcus* strains

Morandi, Stefano¹; Silvetti, Tiziana¹; De Noni, Ivano²; Stuknytė, Milda³; Brasca, Milena¹

¹ Istituto di Scienze delle Produzioni Alimentari, Consiglio Nazionale delle Ricerche, Italy;

²Dipartimento di Scienza per gli Alimenti, la Nutrizione e l'Ambiente, Università degli Studi di Milano; ³ Università degli Studi di Milano, Unitech COSPECT

Nisin is a small peptide produced by *Lactococcus* strains that exerts an antimicrobial activity against pathogenic (i.e., *Listeria monocytogenes*) and spoilage bacteria (i.e., *Clostridium* responsible of blowing defects in cheese).

Five *Lactococcus* were selected for their ability to produce nisin at the optimal growth temperature (30 °C); one strain (FT27) was able to produce Nisin A, while the other 4 were Nisin Z producers. Moreover, *Lc. lactis* VC106 harboured, in addition to Nisin Z, the *lct481* and *lcnB* genes coding for the Lacticin 481 and Lactococcin B synthesis.

At optimal growth condition, the bacteriocin production resulted to be strain dependent and a great variability was observed among the five biotypes (from 239 to 1995 IU/mL).

The strains were then used in preliminary trials to verify that they expressed their bioprotective activity in milk during the cheesemaking and ripening process that is typical of soft and semi cooked cheeses.

In this case, only one strain (VC106) was able to synthesize bacteriocins at cheesemaking temperatures, with maximum Nisin production of 729 IU/mL and persistence up to 170 days under refrigerated conditions.

The strain was then used in starter cultures in the Valtellina Casera PDO cheese productions. Twenty-one cheesemakings were performed in 4 different dairies plants, and in 13 out of 21 the antimicrobial activity was detected at different levels throughout ripening period, achieving the maximum after 110 days of storage.

This study provided important evidence on the bioprotection activity of *Lactococcus* strain VC106 under real application conditions of semi-cooked cheese production.

For a fruitful use as bioprotective starter culture, further knowledge must be gained 1) about the interaction of the bioprotective strain with the starter lactic acid bacteria used in cheesemaking and 2) regarding process conditions that maximize bacteriocin production, as well as the kinetics of their degradation in the cheese during ripening.

Acknowledgements

Funder: Lombardy Region, Project SIMCA PSR 2014-2020 Operazione 16.2.01 "Progetti pilota e sviluppo di innovazione"; Project funded under the National Recovery and Resilience Plan (NRRP), Mission 4 Component 2 Investment 1.3 Call for tender No. 341 of 15 March 2022 of Italian Ministry of University and Research funded by the European Union – NextGenerationEU; Award Number: Project code PE00000003, Concession Decree No. 1550 of 11 October 2022 adopted by the Italian Ministry of University and Research, CUP B83C22004790001, Project title "ON Foods Research and innovation network on food and nutrition Sustainability, Safety and Security – Working ON Foods".

Dietary Microbes and Case Studies from Food Processing

Omri, Ghofrane¹; Fragasso, Mariagiovanna¹; Riso, Patrizia²; Guglielmetti, Simone³; Spano, Giuseppe¹; Capozzi, Vittorio⁴

¹ Department of Agricultural Sciences, Food, Natural Resources and Engineering, University of Foggia, Foggia, Italy; ² Department of Food, Environmental and Nutritional Sciences (DeFENS), University of Milan, Italy; ³ Department of Biotechnology and Biosciences (BtBs), University of Milano-Bicocca, Milan, Italy; ⁴ National Research Council of Italy Institute of Sciences of Food Production (ISPA-CNR), Foggia, Italy

Dietary microbes can contribute in maintaining gut health and overall well-being, influencing digestion, immunity, and metabolic processes. The microbial content in food matrices is significantly affected by processing methods such as pasteurisation, cooking, and handling techniques, which can alter microbial diversity and load. Understanding and optimising factors that influence microbial content in foods can enhance the potential beneficial effects of dietary microbes, supporting the design of tailored investigations. This study explores a selection of case studies, interesting for the definition of microbe-depleted and microbe-rich diets. The impact of food processing on the microbial content of various food matrices, presenting case studies on dietary microbes in i) fermented food processing (i.e. pasteurisation of fermented table olives), ii) within relevant food categories (i.e. fruits, vegetables, cheeses), iii) associated with the fate of a given ingredient during different culinary reparations) (i.e. pecorino cheese). Using culture-dependent methods, we analysed interesting components of the microbial diversity/loads associated with the different matrices experimental matrices. These case studies underscore the critical role of food processing techniques in shaping the microbial loads associated to dietary components. Understanding these effects is crucial for optimizing food processing methods to enhance or preserve microbial content, improving food quality and potential health benefits. Our findings emphasize the need for further research to explore these dynamics across a broader range of foods and processing conditions. By doing so, we can develop more effective strategies in food science and nutrition to maintain or boost the live microbes in our diets.

Acknowledgements

This work was completed as part of the bEat project (We all eat microbes: diet as reservoir of microorganisms that preserve the ecosystem services of the human gastrointestinal microbiota) supported within the call "Progetti di Ricerca di Rilevante Interesse Nazionale (PRIN) 2022" granted by the Italian Ministry for Universities and Research (MUR) (Prot. 2022TF9AHZ, CUP G53D2300574 0006). Vittorio Capozzi was partially funded by CNR project "NUTRAGE FOE2021 DBA.AD005.225". Mariagiovanna Fragasso is supported by the European Union NextGeneration EU [PNRR, CN00000022] within the Agritech National Research Centre for Agricultural Technologies.

Creation of an “Egg-to-Meat” biobank of microbiota collected from broiler chickens raised with or without outdoor access.

Helloin, Emmanuelle¹; Berri, Cécile²; Bordeau, Thierry²; Briandet, Romain³; Chaillou, Stéphane³; Chambellon, Émilie¹; Duclos, Michel²; Germain, Karine⁴; Hondelatte, Alice⁴; Ravon, Laure⁴; Rué, Olivier⁵; Zagorec, Monique⁶

¹ CIRM-BP, INRAE, Nouzilly, France; ² BOA, INRAE, Nouzilly, France; ³ MICALIS, INRAE, Jouy-en-Josas, France; ⁴ EASM, INRAE, Surgères, France; ⁵ MaIAGE, INRAE, Jouy-en-Josas, France; ⁶ SECALIM, INRAE, Nantes, France

Chicken is the most consumed meat worldwide, with a variety of farming schemes from intensive indoor to free-range systems which raises different concerns in terms of animal health and welfare. Recent studies suggest that microbial fluxes in chicken are influenced by the aviary environments, the diet, the broiler's genotype and that various and successive microbial contamination or dispersion events occur in the production chain. A better understanding of these fluxes could help to identify potential levers for controlling microbiota and guaranteeing the health and robustness of animals, as well as the safety of the meat.

The objective of our project was to assess the effect of outdoor access of broiler chickens on bacterial fluxes along the production chain. Microbiota samples were collected from eggshells to carcasses as well as from aviary environment: 1) to analyse microbiota compositions using 16S rDNA amplicon sequencing and search impact of housing (indoors vs outdoor access) on chick colonisation, on the dynamics and shaping of the microbiota during breeding and on the composition of carcasses bacterial contamination; 2) to build a biobank of microbiota. Several zootechnical parameters have also been monitored.

This challenging and extensive study was made possible by the gathering of the varied expertise of partners in a INRAE consortium (France) involving two Poultry experimental units and five joint research units as well as a microbial Biological Resource Center dedicated to animal and human health (CIRM-BP).

A biobank of 620 samples was dispatched in a total of 1981 aliquots preserved at -80°C. The sampling points encompassed eggshells just after hatching, chick droppings and environmental samples during the whole breeding, ceca and carcasses of individualized chickens at slaughterhouse.

The biobank of samples is now available for further studies allowing reanalysis of certain samples if necessary or new analyses such as a culturomics approach.

The work was done by the authors and the whole consortium Egg to Meat.

Lactiplantibacillus plantarum diversity and plant-based sources

Presutto, Ester¹; Capozzi, Vittorio²; Spano, Giuseppe¹; Fragasso, Mariagiovanna¹

¹ Department of Agricultural Sciences, Food, Natural Resources and Engineering, University of Foggia, Foggia, Italy; ² Institute of Sciences of Food Production of National Research Council (ISPA-CNR), Foggia, Italy

Lactic acid bacteria (LAB) play a crucial role in food systems due to their high potential to ensure food quality, safety and shelf life. In particular, the species *Lactiplantibacillus* (*Lpb.*) *plantarum* emerges as a key model for the metabolic versatility, potential in the stress response and ecological diffusion, with a positive impact in food production where it is used as a starter culture, protechnological microbe, probiotic culture and biocontrol agent. Numerous strains of *Lpb. plantarum* have their genome sequenced and are available in specialized databases. There are also many studies on new selections of *Lpb. plantarum* from specific agro-food productions and from different ecological niches. Recently, the scientific community has turned its attention to plant-based matrices as unconventional reservoirs of new strains of *Lpb. plantarum* with distinct genetic and phenotypic traits, thus offering new opportunities to design bio-based innovation in the food industry. Here, we present an overview of the ecological origin of different *Lpb. plantarum* strains, both completely sequenced strains and strains from new selection. In mapping the different origins associated with the intraspecific diversity of *Lpb. plantarum*, we report some new sources of isolation we recently performed, linked to agro-food and the world of plants, highlighting the future efforts that are necessary to increase the ecological representativeness of this species in dedicated bio-banks.

Acknowledgements

We are exploring the potential of this driver of innovation in the framework of the ongoing projects i) the European Union Next-Generation EU [Piano Nazionale di Ripresa e Resilienza (PNRR)—Missione 4 Componente 2, Investimento 1.4—D.D. 1032 17/06/2022, CN00000022] within the Agritech National Research Centre for Agricultural Technologies and ii) the NextGeneration EU [PNRR], in the framework of the Mission 4 Component 2 Investment 1.3-Award Number: Project code PE00000003, Project title: “ON Foods-Research and innovation network on food and nutrition Sustainability, Safety and Security—Working ON Foods”. Vittorio Capozzi was partially funded by CNR project “NUTRAGE FOE-2021 DBA.AD005.225”.

Lactiplantibacillus plantarum and bio-based solutions in food biotechnologies

Presutto, Ester¹; Capozzi, Vittorio²; Spano, Giuseppe¹; Fragasso, Mariagiovanna¹

¹ Department of Agricultural Sciences, Food, Natural Resources and Engineering, University of Foggia, Foggia, Italy; ² Institute of Sciences of Food Production of National Research Council (ISPA-CNR), Foggia, Italy

Lactic acid bacteria (LAB) play a crucial role in food processing, as they can positively affect safety and all the aspects of food quality (e.g. sensory, nutritional, functional), and they also have interesting applications in agriculture. LAB represent bio-resource to design solutions inspired by nature to support the green transition of agro-food systems. Among LAB, lactobacilli are a heterogeneous group of different genera (recently re-classified from the taxonomic point of view) with relevant applications from the farm to the fork. Here, we propose the species *Lactiplantibacillus (Lpb.) plantarum* as a model species for lactobacilli because of the nomadic lifestyle and the huge versatility in ecological diffusion. In contrast to many other lactobacilli species, which have a spectrum of adaptation to a limited number of ecological niches, the ubiquity of *Lpb. plantarum* emphasizes the wide range of metabolic targets and the potential in the stress response. *Lpb. plantarum* strains are used as starter cultures, probiotic cultures, and biopreservation agents, with several properties that are used to pursue economic, social, and environmental sustainability. Through a selection of case studies, we want to provide an overview of the broad-spectrum potential of *Lpb. plantarum* in agro-food productions, with a particular focus on the protechnological, probiotic and biocontrol traits that configure this species as an example for bio-based innovations in the field of food biotechnology.

Acknowledgements

We are exploring the potential of this driver of innovation in the framework of the ongoing projects i) the European Union Next-Generation EU [Piano Nazionale di Ripresa e Resilienza (PNRR)—Missione 4 Componente 2, Investimento 1.4—D.D. 1032 17/06/2022, CN00000022] within the Agritech National Research Centre for Agricultural Technologies and ii) the NextGeneration EU [PNRR], in the framework of the Mission 4 Component 2 Investment 1.3-Award Number: Project code PE00000003, Project title: “ON Foods-Research and innovation network on food and nutrition Sustainability, Safety and Security—Working ON Foods”. Vittorio Capozzi was partially funded by CNR project “NUTRAGE FOE-2021 DBA.AD005.225”.

Enhancing bioprotection on fresh-cut apples with microwave heating: preliminary tests for inclusion in hurdle technology

de Chiara, Maria Lucia Valeria¹; De Simone, Nicola¹; Palumbo, Michela¹; Russo, Pasquale²; Capozzi, Vittorio¹

¹ Institute of Sciences of Food Production of National Research Council (ISPA-CNR), Foggia, Italy; ² Department of Food, Environmental and Nutritional Sciences, University of Milan, Milan, Italy

The inclusion of microwave (MW) and biocontrol agents in a 'hurdle' approach represents a novel method for preserving the quality and safety of fresh-cut produce. This study explores the application of MW heating to enhance the bioprotective efficacy of *Lactiplantibacillus plantarum*, employing a Central Composite Design (CCD) and response surface methodology to determine the optimal treatment conditions for minimally processed apples. Microwave treatment parameters, including power level and treatment time, were varied according to the CCD matrix. The effect of MW applied before or after *L. plantarum* (UFG 121) inoculation, aiming to identify the conditions that can enhance the biocontrol efficacy of this strain without compromising the apples quality was studied. Preliminary results demonstrated that the power level and exposure time influenced the extent of the MW impact. Optimal conditions identified through CCD analysis indicated a synergistic effect between moderate MW power and shorter exposure times, which enhanced the survival of the biocontrol agent. Quality parameters, including colour, visual quality, and chemical aspects, were assessed post-treatment and after 7 days of storage. The optimal MW treatment should maintain the apples colour while minimizing quality losses. The sensory evaluation confirmed that fresh-like attributes were significantly affected by power and time of treatment. In conclusion, the study established that a carefully calibrated microwave treatment, in conjunction with the application of lactic acid bacteria, could be a suitable technology to extend the shelf life and safety of fresh-cut apples. The CCD approach was exploited to preliminarily define the optimal conditions, balancing microbial survival and product quality. This dual treatment strategy holds promise for the fresh produce industry, offering a feasible method to deliver high-quality, fresh, and safe products to consumers.

Acknowledgements

Project funded under the National Recovery and Resilience Plan (NRRP), Mission 4 Component 2 Investment 1.3 Call for tender No. 341 of 15 March 2022 of Italian Ministry of University and Research funded by the European Union – NextGenerationEU; Award Number: Project code PE00000003, Concession Decree No. 1550 of 11 October 2022 adopted by the Italian Ministry of University and Research, Project title "ON Foods Research and innovation network on food and nutrition Sustainability, Safety and Security – Working ON Foods". Vittorio Capozzi was partially funded by CNR project "NUTRAGE FOE-2021 DBA.AD005.225".

Biocontrol, edible packaging, and physical technologies: towards the design of integrated solutions for postharvest fruits and vegetable management

de Chiara, Maria Lucia Valeria¹; De Simone, Nicola¹; Gökşen, Gülден²; Capozzi, Vittorio¹

¹ *Institute of Sciences of Food Production of National Research Council (ISPA-CNR), Foggia, Italy;* ² *Department of Food Technology, Vocational School of Technical Sciences at Mersin Tarsus Organized Industrial Zone, Tarsus University*

The application of hurdle technology and integrated solutions in postharvest handling integrates multiple conventional and emerging preservation methods to ensure microbial safety and maintain the quality of minimally processed horticultural products. This approach, tailored to specific matrices and challenges, exploits the synergistic effects of various agents, which are not individually decisive, to create a series of preservative factors that microorganisms cannot overcome. The primary hurdles in food preservation include high or low temperatures, natural or synthetic preservatives, biocontrol microorganisms, packaging solutions and emerging mild physical treatments. In the context of fresh fruit and vegetable minimal processing, the main objective is to meet safety standards without compromising food quality and consumer health. Innovative combinations of hurdles, their intensity, and their sequence offer a promising future for minimally processed fruits and vegetables, reducing the intensity of single techniques while preserving freshness and quality. This work focuses on the main hurdles/integrated approaches that include at least one emerging physical treatment and/or one edible packaging solution and/or one biocontrol application, assessing their impact on the quality and safety of fresh-cut fruits and vegetables. Combining emerging physical approaches or edible packaging solutions with microbial biocontrol agents represents a promising area of study. Applying physical treatments in combination with biocontrol inoculation could enhance the effectiveness and dominance of biocontrol agents, offering a dual mechanism for improving the safety and quality of fresh produce. The inclusion of biocontrol agents in edible packaging presents new opportunities for the application of lactic bacteria on the surface of products, allowing their contribution to be enhanced with respect to the overall quality of the products.

Acknowledgements

We are exploring the potential of this driver of innovation in the framework of the ongoing projects i) INTACTBioPack INTElligent, ACTive MicroBIOMebased, biodegradable PACKaging for Mediterranean food – founded by the PRIMA Section 2 2023 multi-topic and ii) the NextGeneration EU [PNRR], in the framework of the Mission 4 Component 2 Investment 1.3-Award Number: Project code PE00000003, Project title: “ON Foods-Research and innovation network on food and nutrition Sustainability, Safety and Security–Working ON Foods”.

Application of lactic acid bacteria in the postharvest of fruits and vegetables

Palumbo, Michela¹; De Simone, Nicola¹; Scauro, Angela²; Cefola, Maria¹; Spano, Giuseppe²; Fragasso, Mariagiovanna²; Capozzi, Vittorio¹; De Chiara, Maria Lucia¹

¹ *Institute of Sciences of Food Production of National Research Council (ISPA-CNR), Foggia, Italy;* ² *Department of Agriculture Food Natural Science Engineering (DAFNE), University of Foggia, Foggia, Italy*

The heterogeneous group of lactic acid bacteria (LAB) provides the basis of several bio-based solutions in postharvest applications. The complex microbial ecosystems associated with fruit and vegetables include a wide variety of naturally associated bacteria, yeasts, and moulds. This microbiota may be connected with undesired pathogens, microbes displaying spoilage activities, but can also represent the source of new LAB isolations. One of the main sustainable applications of LAB on fruits and vegetables concerns the potential for biocontrol. After harvest, the handlings may increase these undesired microorganisms, negatively impacting the quality of these products and causing a reduction in yield, shelf-life, safety and marketability, with consequent production losses. Biocontrol, considered one of the most sustainable postharvest approaches, consists of applying selected microorganisms that can limit the development of undesired ones. Selected LAB have been successfully applied on several types of fresh products (leafy green, mixed salads, lettuce, potato, mushroom, tomato, melon, apple, table grape, strawberry, kiwifruit, banana) having as target microorganisms mainly pathogenic bacteria (e.g., *Listeria monocytogenes*, *Escherichia coli*) and/or moulds (e.g., *Botrytis cinerea*, *Penicillium expansum*). LAB not only improve shelf-life and safety, but can also contribute to enhancing the nutritional and functional quality of foods, enriching products with probiotics and dietary microbes, and supporting the beneficial aspects associated with this type of matrix. The combination of LAB with other physical, chemical and biological strategies to obtain synergistic effects is a much-discussed aspect, highlighting the importance of studying the stress response in bacteria as a driver for the design of innovative solutions.

Acknowledgements

We are exploring the potential of this driver of innovation in the framework of the ongoing European Union Next-Generation EU-funded projects i) "ON Foods-Research and innovation network on food and nutrition Sustainability, Safety and Security-Working ON Foods" [Project code PE00000003] and ii) "AgriTech National Research Centre for Agricultural Technologies" [N00000022]. Vittorio Capozzi was partially funded by CNR project "NUTRAGE FOE-2021DBA.AD005.225". Giuseppe Spano was supported by PON project "Conservabilità, qualità e sicurezza dei prodotti ortofrutticoli ad alto contenuto di servizio"-POFACS-CUP B74I20000120005, while Angela Scauro by "the bEat project" PRIN 2022".

Microbial Diversity from fermented Apulian ‘*Gentile di Puglia*’ sheep milk: polyphasic characterization, dairy valorisation and cross-over cereal based applications

Cunedioğlu, Hülya¹; Capozzi, Vittorio²; Molinaro, Marcella³; De Santis, Letizia³; Gagliardi, Roberto⁴; Spano Giuseppe¹; Fragasso, Mariagiovanna¹

¹ Department of Agricultural Sciences, Food, Natural Resources and Engineering, University of Foggia, Foggia, Italy; ² Institute of Sciences of Food Production of National Research Council (ISPA-CNR), Foggia, Italy; ³ Scienzanova srl, Termoli, Italy; ⁴ Scienzanova srl, Termoli, Italy; ⁴ Mediterranea Biotechnologie srl, Termoli, Italy

Traditional fermented dairy products are important for preserving Geographical Indications and gastronomic heritage. These foods and beverages are also source of nutrients and of functional properties and a reservoir of lactic acid bacteria interesting for the design of new starter cultures and biotechnologies. This study aimed to perform a polyphasic characterisation of the microbial diversity of fermented ‘*Gentile di Puglia*’ sheep milk from Subappennino Dauno (Apulia, Italy). Sheep milk samples were fermented under different conditions to selectively isolate microorganisms. Gram staining, catalase, CO₂ production analysis, and microscopic evaluation were performed to perform a preliminary characterisation of the isolates. Additionally, the antimicrobial and antifungal activities of all the isolates against several targets representative of spoilage and pathogens in the food industry were also assessed. The study identified a diverse microbial population in the fermented sheep milk, with the heterogeneous group of lactic acid bacteria (LAB) predominating. Protechnological and genetic characterisations have been performed on a panel of strains. Based on the results, some selected LAB strains were used as a multi-strain starter culture to produce Pecorino cheese from ‘*Gentile di Puglia*’ sheep milk. The application of selected strains in Pecorino cheese production has been performed to obtain a different sensory profile and to improve shelf-life compared to commercial starter culture, highlighting the importance of valorising local dairy products by exploiting autochthonous microbiota. Finally, a selection of LAB strains have been tested in cereal matrices to evaluate the design of plant-based yoghurt-like products. The cereal tested in the trial were also original of Daunia region, promoting an experimental plan that valorises animal, plant and microbial biodiversity associated with a specific marginal area.

Acknowledgements

We are exploring the potential of this driver of innovation in the framework of the ongoing projects i) the European Union Next-Generation EU [Piano Nazionale di Ripresa e Resilienza (PNRR)—Missione 4 Componente 2, Investimento 1.4—D.D. 1032 17/06/2022, CN00000022] within the Agritech National Research Centre for Agricultural Technologies and ii) the NextGeneration EU [PNRR], in the framework of the Mission 4 Component 2 Investment 1.3-Award Number: Project code PE00000003, Project title: “ON Foods-Research and innovation network on food and nutrition Sustainability, Safety and Security—Working ON Foods”. Vittorio Capozzi was partially funded by CNR project “NUTRAGE FOE-2021 DBA.AD005.225”.

Characterization of the rib operon riboswitch in riboflavin overproducing *Lactiplantibacillus plantarum* strains

Scauro, Angela¹; Ruiz-Masó, José Ángel²; Hernández-Alcántara, Annel²; De Simone, Nicola³; Spano, Giuseppe¹; del Solar, Gloria²; Russo, Pasquale⁴

1 Department Agriculture Food Natural Science Engineering, University of Foggia, Foggia Italy; 2 Centro de Investigaciones Biológicas Margarita Salas (CIB Margarita Salas-CSIC), Madrid, Spain; 3 Institute of Sciences of Food Production, National Research Council (ISPA-CNR), Foggia, Italy; 4 Department of Food, Environmental and Nutritional Sciences, University of Milan, Milan, Italy

Lactic acid bacteria (LAB) are able to produce functional and bioactive compounds for the human health, such as B vitamins. In LAB, the expression of the rib operon for the synthesis of riboflavin (vitamin B₂) is regulated by an FMN-riboswitch consisting of a sensitive domain (aptamer) that binds the effector (FMN) and induces a conformational change in the regulatory domain. In this work, LAB strains isolated from wild fruits of the Mediterranean area were tested for their prototrophy for riboflavin in a chemically defined riboflavin-free medium (CDMRF). The strains able to survive in the absence of riboflavin were then exposed to roseoflavin (a toxic analog of riboflavin). This treatment allowed the selection of spontaneous mutants that had mutations in the riboswitch of the rib operon and were capable of overproducing riboflavin. The ability to produce vitamin B₂ in the CDMRF medium was evaluated by fluorescence. *L. plantarum* strain 187 was found to produce the highest amount of riboflavin (between 5 and 6 mg/L). Molecular characterization of the riboswitch variants was performed for all previously selected mutants by aligning the sequence of the regulatory element of the mutants with the parental. In particular, we observed that the highest vitamin B₂ producer *L. plantarum* 187 had a G to T mutation at position 91 (relative to the transcription initiation site), while the lowest producer had a C to A mutation at position 78. Work is underway with the best vitamin B₂-overproducing *L. plantarum* strain and its ability to produce functional foods, such as fermented plant-based beverages, will be evaluated using different substrates and growth conditions.

Bioprospecting and valorization of citrus crops to obtain preharvest and post-harvest biocontrol agents.

Calpe, Jorge¹; Moreno, Ana¹; Vazquez, Rosa¹; Quiles, Juan Manuel¹; Luz, Carlos¹; Escriva, Laura¹; Meca, Giuseppe¹

¹ *Department of Preventive Medicine and Public Health, Food Sciences, Toxicology and Forensic Medicine, University of Valencia, Valencia, Spain*

Introduction: Citrus fruits are among the most widely grown fruits globally. In 2022, the Food and Agriculture Organization estimated their production at 158.5 million tons, with about one-third of the produce lost to fungal rot. In addition, 20-30% of the inedible parts of these fruits generates significant food waste.

Objectives: To investigate the microbiome of citrus fruits and crops and select and characterize microbes for developing natural alternatives through microbial fermentations using citrus residues. The aim is to prevent fungal contamination and reduce losses during field and storage.

Methods and Results: To isolate strains of lactic acid bacteria (LAB) and fungi from different citrus crops. An initial test determined the antifungal activity of the LAB strains. Selected strains were used to ferment a medium with citrus residues. Antifungal metabolites were studied, and *in vitro* tests were performed against pathogenic fungi isolated. The most effective treatment was applied as a biocoating on fruits to test its biopreservation capacity. The metabolome of the fermented medium was studied to determine its biostimulant and biocontrol capacity, aiming to develop dual purpose products that enhance citrus crop productivity and prevent pathogenic fungal infections. Strains N3B1 showed the best antifungal activity, with MIC/MFC values between 1.8-250g/L. These culture media had a presence of antifungal metabolites, such as lactic, acetic, DL-3-phenyllactic, 3-4-dihydroxyhydrocinnamic, salicylic, and vanillic acids. Tests on oranges showed that treatment reduced the proportion of contaminated oranges by 90% after 10 days of storage and decreased the presence of fungi by 4Log₁₀ units of spores/g of citrus.

Conclusions: The results demonstrate the potential of the fermented medium with citrus residues as a biostimulant and biocontrol agent for the development of natural products in the field and storage, exemplifying a circular economy model.

Fermentation as a driver of innovation for the development of the agri-food sector in marginal areas

Fragasso, Mariagiovanna¹; Cunedioğlu, Hülya^{1,2}; Presutto, Ester¹; Spano, Giuseppe¹; Capozzi, Vittorio³

¹ Department of Agriculture Food Natural Science Engineering, University of Foggia, Foggia, Italy; ² Scienzanova srl; ³ Institute of Sciences of Food Production, National Research Council (ISPA-CNR), Foggia, Italy

The social, economic, and environmental management of marginal areas is crucial for sustainable development and resource valorisation in Italy. Preserving and enhancing biodiversity is a key strategy to promote the innovation of agro-food systems in these regions. Improving the quality and safety of fermented products, selecting indigenous microorganisms, studying fermentative microbiomes, and promoting microbial biotechnologies are important factors supporting green transition in the food sector of marginal areas. In this context, we present an overview of significant advances at the international scale in the field of microbial management to generate added value and support new product development in the food sector in marginal areas. In addition, we propose the Apulian region (Italy) as a model territory for studying this type of valorisation. By mapping products and resources that represent good targets for regional development, we aim to create added value through fermentation and the exploitation of microbial diversity. The mapping includes a review of geographical indications, traditional, typical and artisanal products, and raw materials that can be processed through fermentation or using microbial-based solutions.

Acknowledgements

We are exploring the potential of this driver of innovation in the framework of the ongoing projects i) the European Union Next-Generation EU [Piano Nazionale di Ripresa e Resilienza (PNRR)—Missione 4 Componente 2, Investimento 1.4—D.D. 1032 17/06/2022, CN00000022] within the Agritech National Research Centre for Agricultural Technologies and ii) the NextGeneration EU [PNRR], in the framework of the Mission 4 Component 2 Investment 1.3-Award Number: Project code PE00000003, Project title: “ON Foods-Research and innovation network on food and nutrition Sustainability, Safety and Security—Working ON Foods”. Vittorio Capozzi was partially funded by CNR project “NUTRAGE FOE-2021 DBA.AD005.225”.

Innovation in food fermentations and the cross-over fermentation concept in new product development

Cunedioğlu, Hülya^{1,2}; Fragasso, Mariagiovanna¹; De Santis, Letizia²; Gagliardi, Roberto^{2,3}; Spano, Giuseppe¹; Capozzi, Vittorio⁴

¹ Department of Agriculture Food Natural Science Engineering, University of Foggia, Foggia, Italy; ² Scienzanova srl, Termoli, Italy; ³ Mediterranea Biotecnologie srl, Termoli, Italy; ⁴ Institute of Sciences of Food Production of National Research Council (ISPA-CNR), Foggia, Italy

Fermented products represent a dynamic and constantly evolving sector, that can offer solutions to improve the well-being and nutrition of different population targets. Fermentation also makes it possible to promote sustainable innovation and enhance the quality of foods and beverages. Among the various dynamics of innovation in this sector, the concept of 'cross-over fermentation' in the development of new products should be underlined. Cross-over fermentation has been defined as "processes in which a microorganism from one traditional fermentation process is introduced onto a new substrate and/or to a new partner" [1]. Here, we report a series of case studies to highlight the potential of cross-over strategies in valorising the microbial biodiversity associated with traditional fermentations to design novel fermented foods and beverages. It is a trend that allows the exploitation of the latent biotechnological potential related to studies of isolation and characterisation of autochthonous bacteria, yeasts and filamentous fungi from spontaneous fermentations. In addition, this approach makes it possible to improve the added value of raw materials and stimulate the development of new products, creating a link between traditional fermented products and innovative fermented products. All these aspects are particularly interesting to promote the sustainable potential associated with fermentation processes and to support the development of a plant-based sector designed with solutions inspired by nature and not with ultra-processed foods.

References

[1] doi: 10.1016/j.lwt.2021.111041.

Acknowledgments

We are exploring the potential of this driver of innovation in the framework of the ongoing projects i) the European Union Next-Generation EU [Piano Nazionale di Ripresa e Resilienza (PNRR)—Missione 4 Componente 2, Investimento 1.4—D.D. 1032 17/06/2022, CN00000022] within the Agritech National Research Centre for Agricultural Technologies and ii) the NextGeneration EU [PNRR], in the framework of the Mission 4 Component 2 Investment 1.3-Award Number: Project code PE00000003, Project title: "ON Foods-Research and innovation network on food and nutrition Sustainability, Safety and Security—Working ON Foods".

Selection of *Bacillus* strains as potential candidate against pathogenic *Fusarium* occurring on cereal

Masiello, Mario¹; Touati, Rim¹; Somma, Stefania¹; Haidukowski, Miriam¹; Moretti, Antonio¹; De Bellis, Palmira¹

¹ Institute of Sciences of Food Production, National Research Council (ISPA-CNR), Bari, Italy

Fusarium species are among the most important pathogens of cereals able to synthesize harmful mycotoxins. The most important cereal diseases are caused by co-occurrence of multiple *Fusarium* species, mainly *F. graminearum* and *F. culmorum*, producers of deoxynivalenol, and *F. proliferatum*, producer of fumonisins. Nowadays, the increasing interest to reduce chemicals in agriculture, prompts researchers to select eco-friendly strategies against fungal diseases.

This study aimed to select potential bacterial bio-control agents to control *Fusarium* species. Forty-eight *Bacillus* strains belonging to *Bacillus velezensis*, *B. amyloliquefaciens*, *B. subtilis*, *B. licheniformis*, *B. mojavensis*, *B. simplex*, *B. megaterium*, *B. oleronius*, *B. pumilus* and *B. safensis*, isolated from wheat and maize kernels, were tested. The antagonistic activity against *F. graminearum*, *F. culmorum* and *F. proliferatum*, by co-culture assay, and the antimicrobial effect of bacteria filtrates were evaluated. Twenty-three strains, including all *B. velezensis* and *B. amyloliquefaciens* strains, showed a good antagonistic activity, with mycelial growth inhibition values up to 70%. Moreover, 14 strains produced active compounds inhibiting mycelial growth up to 60%. To elucidate the molecular mechanisms associated to the activity of these strains against *Fusarium* species, all selected strains were screened for the presence of genes involved in the synthesis of active biomolecules, including surfactin, fengycin, iturin, bacillomycin, bacilysin, difficidin and mycosubtilin. The gene fragments were also sequenced and analysed, to identify, among and within *Bacillus* species, possible polymorphisms associated to the variable capability to inhibit *Fusarium* growth. In addition, the possible capability of *Bacillus* strains to influence deoxynivalenol and fumonisin production was studied. Although no activity was observed against fumonisin production, half of *Bacillus* strains inhibited completely DON production.

Acknowledgements

This work was supported by Progetti@CNR "Cold Plasma Processes and Technologies for Sustainable Agriculture and Food Processing"

Effect of plant essential oils on *Botrytis cinerea*

Novotný, David¹; Němcová, Karolína²

¹ Crop Research Institute (Výzkumný ústav rostlinné výroby, v.v.i.), Praha, Czech Republic; ² Czech University of Life Sciences Prague, Faculty of Agrobiological Sciences, Food and Natural Resources, Praha, Czech Republic

Botrytis cinerea is a fungus that causes pre-harvest and post-harvest diseases of various plants and cabbage (*Brassica oleracea* convar. *capitata*) is one of them. *B. cinerea* causes grey mould of cabbage and this disease is little significant in the pre-harvest phase, but is a serious problem for stored cabbage heads. Protecting stored cabbage is difficult because the use of synthetic chemical plant protection products (pesticides) is very limited to protect the health of consumers during storage. Some essential oils have the ability to reduce the growth of certain fungi and are environmentally friendly. The ability of thyme, cinnamon, cloves, oregano and lemongrass essential oils to suppress the growth of *B. cinerea* strains from Culture collection of microorganisms of Crop Research Institute (VURV) was investigated in *in vitro* experiments. Essential oils at five concentrations were added to agar culture medium in Petri dishes and their effect on the growth of *B. cinerea* colonies was evaluated. Differences were found in the effect of individual essential oils and between different concentrations, as well as between different strains of *B. cinerea*. Oregano essential oil inhibited growth the most, not allowing growth even at a concentration of 83.33 l/l. Another essential oil that significantly inhibited the growth of *B. cinerea* colonies (completely at the highest concentration evaluated, 250 l/l.) was thyme essential oil. Cinnamon and clove essential oils reduced the growth of *B. cinerea* colonies. The effect on colony growth in all strains examined was evident at all concentrations for these two essential oils, especially at the highest concentration evaluated, 250 l/l. Lemongrass essential oil reduced the growth of *B. cinerea* strainsless, but still suppressed colony growth at the highest concentration evaluated (250 l/l). At lower concentrations, growth was less restricted.

Acknowledgements

The research was supported by the Czech National Programme on Conservation and Utilization of Microbial Genetic Resources Important for Agriculture, Ministry of Agriculture of the Czech Republic and Mze-RO0423 (Ministry of Agriculture of the Czech Republic).

Lactic acid bacteria as microbial starters in malting by-product fermentation to obtain innovative bread formulation

Palumbo, Michela¹; De Simone, Nicola¹; De Chiara, Maria Lucia¹; Capozzi, Vittorio¹; Montemurro, Marco²

¹ Institute of Sciences of Food Production of National Research Council (ISPA-CNR), Foggia, Italy; ² Institute of Sciences of Food Production of National Research Council (ISPA-CNR), Bari, Italy

Sourdough-based preparations represent attractive models for fortifying food product with microbially treated by-products. The malting by-products represent a significant waste and they have not been exploited in details. Malting rootlets and malt glumellae were selected to test fermentation by LAB previously characterized for other food applications as microbial starters, with the final aim to include the fermented by-product in one innovative bread formulation to increase shelf-life, digestibility and nutritional value.

Malting by-products have been characterised to study the presence of microbial contaminants before and after the fermentation process. In detail, *Lactiplantibacillus plantarum* (PLA) and *Lacticaseibacillus rhamnosus* (RHM), isolated from milk and cheese respectively, were individually inoculated at initial cell density of ca. Log 7.0 cfu/g in malting rootlets and mixture of rootlets and glumellae dough (dough yield of 400). The dough was fermented at 30 °C for 24 h, and the growth and acidification of dough fermented by PLA and RHM were monitored every 2 hours. At each sampling hour, LAB count, pH and titratable acidity of the dough were measured, while dough colour, antioxidant activity and total phenols were detected at the beginning and the end of fermentation.

The enterobacteria in malting rootlets and malting rootlets and glumellae decreased at a density lower than 100 CFU/g. The total free amino acids of doughs fermented increased up to 140%. The growth and acidification of dough fermented by PLA and RHM showed a lag phase lower than 3 hours for both parameters. High growth speed was found in rootlets, while higher acidification speed in mixed matrix.

Finally, preliminary results on the inclusion of fermented and unfermented malting rootlets (5% w/w of flour total weight) in fortified bread formulations showed good sensorial acceptability.

The inclusion in the sourdough of two different strains of yeast will be tested.

Acknowledgements

The project "ON Foods—Research and innovation network on food and nutrition Sustainability, Safety and Security—Working ON Foods" (Project code PE00000003, Concession Decree No.1550 of 11 October 2022) founded under the National Recovery and Resilience Plans (NRRP) Mission 4 Component 2, Investment 1.3 – Call for tender No. 341 of 15 March 2022 of Italian Ministry of University and Research founded by the European Union – NextGenerationEU is acknowledged

Unconventional raw plant matrices as source of pro-technological microbes for application in food fermentation.

De Bellis, Palmira¹; Di Biase, Mariaelena¹; Cifarelli, Valentina¹; Lonigro, Stella Lisa¹; Valerio, Francesca¹

¹ *Institute of Sciences of Food Production, National Research Council (ISPA-CNR), Bari, Italy*

The use of unconventional raw materials within food processing is becoming a widely used sustainable and innovative approach for developing food products with enhanced nutritional, organoleptic, technological, functional and preservative characteristics to satisfy the increasing request for a healthy diet. Alternative matrices can also be a source of microbial strains with enzymatic features, which can be useful for enriching food ingredients in bioactive compounds. Biotechnological processes of fermentation have been used for centuries to produce food often using selected starters, such as lactic acid bacteria (LAB) and yeasts with enhanced nutritional, organoleptic, technological, functional, and preservative molecules. However, the selection of a good candidate as starter strain needs molecular characterization and evaluation of safety and potential pro-technological activities.

In this work different cereal by-products (flours of wheat bran and wheat germ, defatted or not), vegetable by-product (fresh substandard peas) and finally flours obtained from Okra (*Abelmoschus esculentus* L.), a native African plant, were collected.

In order to isolate beneficial microorganisms, these by-product matrices were plated onto MRS agar. A total of 20% of the colonies randomly picked from countable plates, were purified and stored at -80°C. The bacterial isolates were characterized by repetitive extragenic palindromic PCR (REP-PCR) and isolates representative of each REP-PCR profile (19) were identified by sequencing of the 16S rRNA gene. Among strains isolated, only those belonging to the Qualified Presumption of Safety (QPS) list were characterized for antibiotic resistance, biogenic amine production, EPS production and proteolytic activity.

Bacterial strains were deposited in the CNR-ISPA Culture Collection (ITEM Culture Collection) in cooperation with SUS-MIRRI.IT (an Infrastructure Research Project of the National Recovery and Resilience Plan (NRPP)).

Acknowledgements

The work was supported by Project SUS-MIRRI.IT Strengthening the MIRRI Italian Research Infrastructure for Sustainable Bioscience and Bioeconomy, D.D. n. 3264 del 28 Dicembre 2021 –Mission 4 Component 2 Investment 3.1; Project ExpOkra, CNR/MHSR 2023-2024 – AD002-186; Project Agritech National Research Center for Agricultural Technologies – Project PNRR M4C2, Investimento 1.4; Project “ON Foods –Research and innovation network on food and nutrition Sustainability, Safety and Security – Working ON Foods”, funded under the National Recovery and Resilience Plan (NRRP), Mission 4 Component 2 Investment 1.3 – NextGenerationEU Project code PE00000003.

Real-Time investigation of *Lactiplantibacillus plantarum* Fermentation in Experimental and Real Food Conditions Using a Metabolomics Approach.

Corvino, Antonia¹; Khomenko, Iulia¹; Betta, Emanuela¹; Capozzi, Vittorio²; Biasioli, Franco¹

¹ Research and Innovation Centre, Fondazione Edmund Mach, San Michele all'Adige (TN), Italy;

² Institute of Sciences of Food Production of National Research Council (ISPA-CNR), Foggia, Italy

Many food industries use *Lactiplantibacillus plantarum*, a ubiquitous facultative heterofermentative lactic acid bacterium (LAB), as a starter culture due to its technological potential and versatility in adapting to different food environments. To understand and optimize fermentation, 'omics' technologies, especially metabolomics, are employed, offering insights into the metabolic state of microorganisms and the dynamics of metabolic pathways. Microbial volatile organic compounds (VOCs), which influence odour and flavour, serve as biomarkers to track microbial metabolism during fermentation, providing information on the quality of the matrices. Among the sensors used to monitor VOCs in the food and beverage sector, Direct Injection Mass Spectrometric (DIMS) techniques have gained increasing interest due to their versatility, real-time analysis, and strong analytical performance. Proton-Transfer-Reaction coupled with Time-of-Flight Mass Spectrometer (PTR-ToF-MS) is a prime example of DIMS technology, known for its sensitivity, accuracy, time efficiency, non-invasiveness, and eco-friendly analysis. PTR-ToF-MS measurements allow for the assessment and monitoring of VOCs i) during fermentation processes to track their evolution and potential reaction kinetics, and ii) in final products to provide information on possible consumer sensory experiences and product quality. This technique offers a green alternative for profiling volatiles during food fermentations [1]. In this study, PTR-ToF-MS was used for the online monitoring of the fermentation of i) six *L. plantarum* strains in MRS for 32 hours, and ii) one selected *L. plantarum* strain from the previous experiment in three plant-based (soy, almond, and oat) beverages and one animal-origin beverage (milk) as a control for 72 hours. This preliminary investigation also aims to develop a PTR-based strategy to build a reference framework for the development and design of fermented beverages.

References

- [1] V. Capozzi, S. Yener, I. Khomenko, B. Farneti, L. Cappellin, F. Gasperi, M. Scampicchio, F. Biasioli, PTR-ToF-MS coupled with an automated sampling system and tailored data analysis for food studies: bioprocess monitoring, screening and nose-space analysis. doi:10.3791/54075
- L. Cappellin, F. Biasioli, P.M. Granitto, E. Schuhfried, C. Soukoulis, F. Costa, T.D. Märk, F. Gasperi, On data analysis in PTR-TOF-MS: From raw spectra to data mining, Sens. Actuators B Chem. 155 (2011) 183–190.

Acknowledgements

We developed this overview in the framework of the projects i) "Interconnected Nord-Est Innovation Ecosystem (iNEST) funded by the European Union Next-Generation EU (PIANO NAZIONALE DI RIPRESA E RESILIENZA (PNRR) – MISSIONE 4 COMPONENTE 2, INVESTIMENTO 1.5 D.D. 1058 23/06/2022, ECS00000043) and ii) "ON Foods-Research and innovation network on food and nutrition Sustainability, Safety and Security – Working ON Foods" funded by NextGeneration EU [PNRR], in the framework of the Component 2 Investment 1.3 Award Number: Project code PE00000003, Vittorio Capozzi is also supported by NUTRAGE CNR project FOE2021 DBA.ADO05.225.

Cell surface proteome of *L. plantarum* strains isolated from vegetable foods and dairy products

Mazzeo, Maria Fiorella¹; Sorrentino, Alida¹; Cufino, Gaia¹; Morandi, Stefano²; Abouloifa, Houssam³; Asehraou, Abdeslam³; Brasca, Milena²; Siciliano, Rosa Anna¹

¹ Institute of Food Sciences, National Research Council (CNR-ISA), Avellino, Italy; ² Institute of Sciences of Food Production, National Research Council (CNR-ISPA), Milan, Italy; ³ Laboratory of Bioresources, Biotechnology, Ethnopharmacology and health, Faculty of Sciences, Mohammed Premier University, Oujda, Morocco

Lactiplantibacillus plantarum (formerly *Lactobacillus plantarum*) exhibits relevant probiotic and technological features and is widely used in agrifood sector as plant growth-promoting bacteria and to improve quality properties of food products. Surface proteins are directly involved in functional and technological traits of bacteria and their dynamic cross-talk with the host. Many genomic studies were carried out to predict the exoproteome of this species revealing the presence of several classes of proteins differently linked to cell surface [1]. Although proteomics deeply contributed to investigate surface architecture of bacteria cells leading to define protein location and topology [2], a limited number of proteomic studies have investigated the repertoire of *L. plantarum* surface proteins. In this context, we performed a shotgun proteomic study to identify the proteins present on the surface of four different *L. plantarum* strains (two isolated from vegetable foods, V strains, and two from dairy products, D strains) aimed at highlighting differences in protein profiles potentially related to habitat of origin and specific properties of the analyzed strains. Proteins predicted to contain a signal peptide for extracellular translocation, transmembrane proteins and moonlighting proteins known to be involved in adhesion processes were identified. Results showed a more complex pattern of surface proteins in V strains compared to that from D strains. Cell surface proteomes of V strains were characterized by a higher number of proteins displaying glycosyltransferase activity and peptidoglycan hydrolases compared to those of D strains. Peculiar molecular functions of proteins identified in V strains suggested a probiotic and biotechnological potential of these bacteria.

References

- [1] Albarracín L et al. 2022. Genomic characterization of *L. plantarum* strains possessing differential antiviral immunomodulatory activities. *Bacteria* 1, 136.
- [2] Siciliano RA et al. 2019. Proteomics for the investigation of surface exposed proteins in probiotics. *Front Nutr* 6, 52.

Acknowledgements

This study was funded by the European Commission—NextGenerationEU, Project AGRITECH National Research Center (code n. CN00000022) and Project SUS-MIRRI.IT “Strengthening the MIRRI Italian Research Infrastructure for Sustainable Bioscience and Bioeconomy” (code n. IR0000005).

Biodiversity in natural fungal community isolated from cheese aged in Apulian cave

Anelli, Pamela¹; Ferrara, Massimo¹; Cozzi, Giuseppe¹; Carella, Daria¹; Haidukowski, Miriam¹; Mulè, Giuseppina¹; Baruzzi, Federico¹; Moretti, Antonio¹; Susca, Antonia¹

¹ *Institute of Sciences of Food Production, National Research Council of Italy (CNR-ISPA), Bari, Italy*

Filamentous fungi contribute to the organoleptic characteristics, texture and color of long-ripened cheese, with their metabolism, acting as secondary starters.

Traditionally worldwide cheesemakers gathered empirically that cheese production is affected by environment, technological procedures, and especially by microorganisms. Cheesemakers have naturally selected through time strains able to ensure the correct seasoning of cheeses, from which later some strains became commercial cultures, intentionally inoculated for driven-maturations of cheese.

Currently, there is a reversal trend compared the past that appreciates traditional protocols for food manufacturing in light of an increased awareness of the importance of food safety. Therefore, the interest in typical local foods and traditional protocols have led to branch out the way of aging, exploiting sustainable ripening in natural environments like caves, similarly to the past. In these natural environments, cheeses come into contact with non-dairy microorganisms, potentially able to affect positively or negatively cheese aging.

The mycobiota of caciocavallo cheese, aged in cave of an Apulian dairy factory, was analyzed to select pro-technological molds compatible with cheese aging, according to quality and safety parameters. Fungal community was isolated, taxonomically identified by DNA-based analysis and characterized for their relevant technological properties. Two strains belonging to non-toxicogenic species, *Penicillium flavigenum* and *P. biforme*, were selected as secondary starters for caciocavallo pilot aging in cave. Main results related to their isolation, characterization, inoculation of cheese and the preliminary sensory analysis are here reported.

Acknowledgements

This study was supported by Italian Ministry of Economic Development within the project “L’evoluzione delle produzioni lattiero-casearie: le biotecnologie valorizzano la tradizione” – ELEVATO n. F/200112/03/X45.

The authors thank the staff of “Capurso Azienda Casearia srl” for the cooperation during cheese manufacture and ripening.

Dried cell-free supernatant of *Bacillus subtilis* strain ET-1 against *Penicillium digitatum* on clementine fruits

Ambrico, Alfredo¹; Magarelli, Rosaria Alessandra¹; Trupo, Mario¹; Larocca, Vincenzo¹; Martino, Maria¹; Palazzo, Salvatore¹; Della Mura, Brigida²

¹ ENEA, Department for Sustainability, Trisaia Research Center, Rotondella (MT), Italy; ² University of Basilicata, Department of Science, Potenza, Italy

In this work a low medium for *Bacillus subtilis* growth was designed with the aim to obtain a cell-free supernatant (CFS) rich in secondary metabolites for post-harvest control of green mould. For medium development, culture growth trials with *B. subtilis* ET-1 strain were carried out in flasks using different agro-industrial wastes as carbon and nitrogen sources and Iturin A concentration was determined. The medium with higher level of Iturin A (2,45 g L⁻¹) was choice and used for scaling up process. In specific, the bacterium was grown in a 21 L stirred-tank bioreactor with working volume of 10 L at temperature of 25 °C and without pH controlling.

At the end of culture, the supernatant was clarified from the cells by centrifugation and five litres of CFS was stabilized by drying processes with skim milk addition.

Then, the effectiveness of both liquid and stabilized CFS against *Penicillium digitatum* on citrus fruits (*C. × clementina*) in vivo was evaluated. For the experiments, sanitized fruits were injured and treated before the artificial inoculation with 20 µL of *P. digitatum* suspension at concentration of 2,3x10⁴ conidia mL⁻¹.

The measurements were carried out when the first symptoms of the disease appeared and concerned the presence of infection on the individual wounds.

The results of the artificial inoculations show the excellent disease control effect on the fruits protected with CFS of the ET-1 isolate.

Furthermore, no significant difference in the protection effect was observed between the theses treated with liquid and stabilized CFS. Therefore, the stabilization process conducted with drying technology did not reduce the antifungal activity of the secondary metabolites present in CFS.

In conclusion, the ET-1 isolate could be usefully used to produce alternative formulations to conventional chemical fungicide, thus achieving the objective of the EU "farm to fork" strategy aimed to produce healthier and more sustainable food.

Acknowledgements

This work was supported by The National Recovery and Resilience Plan (NRRP) – Next Generation EU – Mission 4, Component 2, Investment 1.4, under the “National Research Centre for Agricultural Technologies”, Agritech, [grant number I63C22000350007].

Penicillium nalgiovense: a promising fungal starter to control *Aspergillus westerdijkiae* contamination in salami

Copetti V., Marina¹; Gialluisi, Katia²; Ferrara, Massimo²; Carriero, Francesco³; Perrone, Giancarlo²; Magistà, Donato²

¹ Departamento de Tecnologia e Ciência de Alimentos, Universidade Federal de Santa Maria (UFSM), Santa Maria, Brazil; ² Institute of Sciences of Food Production, National Research Council of Italy (CNR-ISPA), Bari, Italy; ³ Salumi Martina Franca srl, Martina Franca, Taranto, Italy

Aspergillus westerdijkiae is a cause of concern in salami production in warmer climates since it can grow on the product surface and synthesize ochratoxin A. This study aimed to evaluate the capability of *Penicillium nalgiovense*, a common mould starter used for salami production, to control the growth of *A. westerdijkiae* during salami production. Salami were co-inoculated with *P. nalgiovense* and *A. westerdijkiae* with contamination rates up to 32% of the spoiler, besides a control with each single species, and a respective salami only inoculated with *A. westerdijkiae*. Salami were seasoned in maturation chambers with controlled temperatures and relative humidities. Mycological analysis has been conducted at the end of seasoning by spread plating on 18% Dichloran Glycerol Agar. Generally, *A. westerdijkiae* colonies were visually not observed in the salami co-inoculated with *P. nalgiovense*, independently of the inoculation rate accessed. On the other hand, *A. westerdijkiae* was able to colonise the salami when no *P. nalgiovense* was co-inoculated. Results demonstrated that the biological control activity of *P. nalgiovense* as starter can limit the *A. westerdijkiae* growth even under high rates of contamination, preventing the human exposure to ochratoxin A.

Acknowledgements

This study was financed in part by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior – Brasil (CAPES) – Finance Code 001.

Biodiversity in fungal community on cheese aged in Dossena mine

Anelli, Pamela¹; Dall'Asta, Chiara²; Cozzi, Giuseppe¹; Epifani, Filomena¹; Carella, Daria¹; Scarpetta, Davide³; Brasca, Milena³; Moretti, Antonio¹; Susca, Antonia¹

¹ Institute of Sciences of Food Production, National Research Council (CNR-ISPA), Bari, Italy; ² Department of Food and Drug, University of Parma, Parma, Italy; ³ Institute of Sciences of Food Production, National Research Council (CNR-ISPA), Milan, Italy

Accurate identification of the fungal community spontaneously colonizing food products, aged in natural and not controlled environments, provides information about potential mycotoxin risk associated with its consumption.

Autochthonous mycobiota colonizing cheese aging in Dossena mines, was investigated and characterized by two approaches: microbial isolations and metabarcoding. Microbial isolations and metabarcoding analysis were conducted on cheese samples, obtained by four batches, produced in four different seasons of the year, aged for 90 and 180 days, by five dairy farms. The two approaches, with different taxonomical resolution power, highlighted *Penicillium biforme* among filamentous fungi, collected from 58 out of 68 cheeses, and *Debaryomyces hansenii* among yeasts, as the most abundant species (31÷65%), none representing a health risk for human cheese consumption. Shannon index showed that the richness of mycobiota increases after 180 days of maturation. Beta diversity analysis highlighted significant differences in composition of mycobiota of cheese produced by different dairy farms and aged for different durations. Weak negative growth interaction between *P. biforme* and *Aspergillus westerdijkiae* by in vitro analysis was observed leading to hypothesize that a reciprocal control is possible, also affected by natural environmental conditions, possibly disadvantageous for the last species.

Acknowledgments

This work was supported by CHEESEMINE project "Percorso di sperimentazione della stagionatura dei formaggi nelle miniere di Dossena", FEASR – Programma di Sviluppo Rurale 2014-2020.

Proteolysis in soy tempeh obtained with commercial starter

Reale, Anna¹; Di Renzo, Tiziana¹; De Caro, Salvatore¹; Ferrara, Massimo²; Spada, Valentina¹; Verrone, Laura²; Gialluisi, Katia²; Nazzaro, Stefania¹; Pasquale, Marena¹; Mamone, Gianfranco¹

¹ *Institute of Food Sciences, National Research Council (CNR-ISA), Avellino, Italy;* ² *Institute of Sciences of Food Production, National Research Council (CNR-ISPA), Bari, Italy*

Tempeh is produced from soya or other legumes fermented by beneficial microorganisms such as moulds and lactic acid bacteria (LAB) that contribute to modify the nutritional composition of the food matrix, by improving the bioavailability of nutrients and phytochemicals [1]. The microbial fermentation process also produces biopeptides with biological effects, which are of great interest in the context of functional foods [2, 3]. The aim of the present research was to study the proteolytic activity of a commercial mold starter during the production of soy tempeh and the microbiota composition of the commercial starter and the final fermented tempeh product.

The biological soybean was soaked for 12 h, boiled for 1.5 h, drained, inoculated with commercial starter mould and incubated in perforated plastic bags at 30°C for 48 h. At time zero, after boiling and after 48 h fermentation, pH, microbial count, degree of protein hydrolysis, bioactive peptides were evaluated. Furthermore, microbiota identification of the commercial starter culture as well as the final tempeh products was evaluated by metabarcoding analysis of 16S rRNA for the identification of bacteria and ITS2 for that of fungi. The proteolytic events occurring during the fermentation process were evaluated by gel-based and gel-free proteomics analysis.

Tempeh starter was composed mainly of a pure culture of *Rhizopus microsporus*. The microbiota of the final tempeh products was dominated by the most abundant genera also detected in the commercial starter, although other bacterial and fungal genera were identified, probably related to environmental contamination.

Proteomics profiling were characterized, identifying several hundreds of peptides released mainly from glycinin and conglycinin soybean proteins.

References

- [1] Teoh S.Q. et al. (2024) A review on health benefits and processing of tempeh with outlines on its functional microbes, *Future Foods*, 9, 100330.
- [2] Rizzo G. (2024) Soy-Based Tempeh as a Functional Food: Evidence for Human Health and Future Perspective. *Front Biosci (Elite Ed)*, 16, 3.
- [3] Tamam B. et al. (2019) Proteomic study of bioactive peptides from tempe, *Journal of Bioscience and Bioengineering*, 128, 2, 241-248.

Acknowledgements

This research was funded by the project SUS-MIRRI.IT “Strengthening the MIRRI Italian Research Infrastructure for Sustainable Bioscience and Bioeconomy”, code n. IR0000005.

Relationship between the chemical content of fruits and endophytic fungal communities in black elder *Sambucus nigra* L.

Boroduske, Anete¹; Rimsa, Antonija¹; Balode, Laura¹; Gudra, Dita²; Nakurte, Ilva³

¹ Department of Microbiology and Biotechnology, Faculty of Medicine and Life Sciences, University of Latvia; ² Latvian Biomedical Research and Study Centre; ³ Institute for Environmental Solutions

It has previously been shown that endophytic microbial communities can affect the phytochemical content of medicinal plants. Here we investigate whether the phytochemical content of wild elderberries is related to endophytic fungal communities inhabiting wild elder plants *Sambucus nigra* L.

Fruit samples were collected at three different stages of maturity from wild *S.nigra* populations located in Latvia. Total DNA was isolated from surface-sterilized fruits and endophytic fungal communities were characterized using ITS1 amplicon NGS sequencing. The phytochemical content of ripe fruits was characterized by assessing total flavonoid content and the content of five dominant compounds was determined using HPLC.

Endophytic microbial communities of *S.nigra* fruits were significantly affected by the geographical location of the sampled population and by the content of one of the dominant chemical compounds of fruits. The diversity and richness of endophytic fungal communities decreased with the progression of fruit ripening. Although the chemical composition and fungal communities significantly varied among studied populations we did not observe the correlation between fruit chemical content and endophytic fungal communities (Mantel $r=0.01$; $p=0.407$). Further studies assessing the effect of individual fungal taxa on fruit phytochemistry might reveal more intricate relationships between endophytic fungal communities and the phytochemical content of *S.nigra* fruits in the future.

Acknowledgment

This research was done under project No lzp-2022/1-0179 "Role of endophytic microbiota in the anti-viral activity of *Sambucus nigra* L. against type A influenza and SARSCoV-2."

Genomic characterization of *Companilactobacillus* spp.: assessing safety and application potential in the food chain

Jaberi, Mehrdad¹; Felis, Giovanna²; Salvetti, Elisa²

¹ Dept. Biotechnology, University of Verona, Verona, Italy; ² Dept. Biotechnology, University of Verona & VUCC-DBT, University of Verona, Italy

Companilactobacillus crustorum is a lactic acid bacterium (LAB) isolated from Belgian wheat sourdough [1], but also found in dairy products and forages [2]. Members of *Companilactobacillus* genus, formed by 34 species, occur consistently in food fermentations of plant and animal origin [2].

Despite the possible applications in food and feed, only two *Companilactobacillus* species hold the Qualified Presumption of Safety (QPS) status, as defined by the European Food Safety Authority (EFSA), therefore a thorough safety assessment is crucial for determining applicability. In this perspective, the availability of over 100 *Companilactobacillus* spp. genome sequences allows the investigation of possible interesting traits related to safety and application.

This study investigates the genomic landscape of *C. crustorum* and related species, focusing on antibiotic resistance genes (ARGs), sanitizing agents resistance genes (SARGs), antimicrobial peptide sequences (AMPs), and biosynthetic gene clusters (BGCs), with an emphasis on their potential implications for food safety throughout the farm-to-fork continuum.

High-quality genomes of *Companilactobacillus* spp. were downloaded from the NCBI database. The nf-core/funcscan pipeline [3] was utilized to predict and annotate ARGs, SARGs, AMPs, and BGCs. In silico analysis offers insights into the distribution of ARGs and SARGs across the genus, aiding in the differentiation between acquired and intrinsic resistance.

The presence of AMPs and BGCs highlights the potential of these bacteria for various applications, while also raising concerns about their impact on food safety and microbial ecosystems along the food chain. The production of AMPs and BGCs could influence microbial communities in food products, affecting spoilage, fermentation processes and, potentially, consumer health. This study contributes to safety assessment of *C. crustorum* and related species within the context of the EFSA QPS approach.

References

- [1] Scheirlinck et al., 2007 <https://doi.org/10.1099/ijjs.o.64836-0>
- [2] Zheng et al., 2020 <https://doi.org/10.1099/ijsem.o.00>
- [3] <https://github.com/nf-core/funcscan>

Acknowledgments

This study was carried out within the Interconnected Nord-Est Innovation Ecosystem (iNEST) and received funding from the European Union Next-GenerationEU (PIANONAZIONALE DI RIPRESA E RESILIENZA (PNRR) – MISSIONE 4 COMPONENTE 2, INVESTIMENTO 1.5 – D.D. 1058 23/06/2022, ECS00000043). This manuscript reflects only the authors' views and opinions, neither the European Union nor the European Commission can be considered responsible for them.

Screening of yeast and filamentous fungi from VTT Culture Collection for cacao butter and palm oil production

Vidgren, Virve¹; Hölttä, Matti¹; Nygren, Heli¹; Barth, Dorothee¹; Koivuranta, Kari¹

¹ *VTT Technical Research Centre of Finland Ltd, Finland*

Yeast and filamentous fungi strains obtained from VTT Culture Collection were screened for their lipid profiles. The objective was to find strains with natural ability to produce fatty acid profiles similar to palm oil or cacao butter. VTT's culture collection contains more than 3000 yeast and filamentous fungi strains, of which 27 yeast strains belonging to 18 different species and 16 filamentous fungi strains belonging to 7 different species were selected based on a literature search revealing the most potential candidates.

Candidates were grown on C/N 75 and PD media and their growth, total lipid content and fatty acid profiles analyzed. The total lipid content varied from 5% to 42% (yeasts) and from 8 to 29% (filamentous fungi). Fatty acid profiles varied significantly among yeasts, especially share of palmitic acid, oleic acid and stearic acid, which are the main constituents of cocoa butter and palm oil.

Three different industrial side streams, obtained as by-products of different food processes, were used as substrates to cultivate the best candidates from the preliminary screen. Two of the side streams were lactose rich whereas one contained mainly glucose and galactose.

For cacao butter and palm oil production, the best candidates were found among yeast strains. Two of the yeast strains produced fatty acid profile close to cacao butter and one yeast strain close to palm oil. None of the fatty acid profiles of filamentous fungi screened resembled cocoa butter or palm oil. Instead, fatty acid profiles of some filamentous fungi were very close to meat fatty acid profile.

Diversity of Non-Conventional Yeasts (NCY) for innovating fermented beverages: investigating microbiomes

Parente, Eugenio¹; Ricciardi, Annamaria¹; Turchetti, Benedetta²; Di Cesare, Francesca³; Vitulo Nicola⁴; Felis, Giovanna³

¹ Department of Agricultural, Forestry, Food and Environmental Sciences, University of Basilicata, Potenza, Italy; ² Dept. Agricultural, Food and Environmental Sciences, Industrial Yeasts Collection DBVPG, University of Perugia, Perugia, Italy; ³ Dept. Biotechnology, University of Verona & VUCC-DBT, University of Verona, Italy; ⁴ Dept. Biotechnology, University of Verona, Italy

Fermented food and beverages represent an important part of the planetary cultural heritage, where microbial consortia contribute to the broad range of sensory and functional properties. Yeast biodiversity plays a crucial role for product diversification in fermented beverages (FB) [1]. Interestingly, fermenting abilities are present not only in Ascomycota, that includes the fermentation workhorse *Saccharomyces cerevisiae*, but also in the much less explored phylum Basidiomycota, with the interesting example of *Mrakia*, with no history of use in the food chain, but with a high innovation potential [2].

In the framework of a research project focused on nonconventional yeasts for FB, an updated version of FoodMicrobionet (FMBN) was realized: FMBN is a food-dedicated and curated repository of metabarcoding studies [3] and it now includes fungal sequences [4], from 1114 samples described in 21 published studies, including, at the date of writing of this abstract (June 2024) 3 on wine, 1 on beer and 6 on other alcoholic beverages.

FMBN 5.0 was used with the two-fold aim of evaluating possible presence of *Mrakia*-related sequences in analyzed datasets, and to define the most important genera that could deserve attention for future studies.

As for *Mrakia*-related sequences, as expected, they were detected at very low levels (relative abundance < 0.25%) in FMBN FB datasets. For this reason, an in-depth characterization of the species is necessary to assess their safety for application in fermentation, before any intentional use in the food chain can be authorized.

As for other genera, the most important ones in terms of prevalence and median abundance in alcoholic beverages were *Saccharomyces*, *Hanseniaspora*, *Pichia*, *Cladosporium*, *Candida* and *Aureobasidium*.

Data presented here on fermented beverages shows how digitalized data on microbiomes are powerful tools to generate new knowledge and insights for developing bioeconomy, supporting responsible innovation in food systems.

References

- [1] Miguel et al. 2022 10.1016/j.ijfoodmicro.2022.109951
- [2] Turchetti et al. 2023. 10.1016/j.foodres.2023.113004
- [3] Parente et al. 2022 10.1016/j.ijfoodmicro.2022.109696
- [4] Parente Ricciardi 2024 doi.org/10.3390/foods13111689

Acknowledgments

Study carried out within the PRIN 2022 PNRR Project “NCYdiversity” P20229JMMH, funded by European Union Next-GenerationEU, CUP C53D23007560001 (PIANO NAZIONALE DI RIPRESA E RESILIENZA (PNRR)—MISSIONE 4 COMPONENTE 2, INVESTIMENTO 1.4—D.D. 1364/2023). This manuscript reflects only the authors' views and opinions, neither the European Union nor the European Commission can be considered responsible for them.

Principles for selection of strains of microorganism cultures for the creation of biopreparations of agricultural designation

Goginyan, Vigen¹; Harutyunyan, Seda¹; Stepanyan, Tamara¹; Khachatryan, Gayane¹; Bagiyanyan, Valeri¹

¹ Scientific and Production Center "Armbiotechnology" NAS RA

Nowadays it is definite that the global growth in the production of biological/microbiological fertilizers is due to the partial, and in some cases, complete replacement of mineral fertilizers in agricultural technology.

The main reason for changes in approaches to intensive agriculture is the ineffective absorption by plants of nitrogen, phosphorus and potassium in the composition of agrochemicals, leading to the gradual salinization of cultivated lands. At the same time, the increase in consumption of agricultural products of plant origin necessitates expanding the range of biological preparations with improved properties for more efficient farming, including organic agriculture. In this regard, the crucial importance lies in the targeted search for new types of microorganisms endowed with increased stress resistance and adaptive capabilities in various types of soils; valuable metabolic characteristics; a powerful enzymatic system; high nitrogen-fixing and phosphate solubilizing ability; antimicrobial, antifungal, selective insecticidal activity; and resistance to various pesticides (pesticides, herbicides, fungicides) when used together; long shelf life and viability.

When selecting promising cultures of microorganisms, special attention should be paid to the reduction of cost of production of biopreparations, the safety of their use in relation to the environment, humans, animals, birds and, especially beneficial insects.

Thus, the goal of our research is to develop the technological basis for the production of biopreparations of agricultural designation using active strains of bacteria *Azotobacter chroococcum*, *Paenibacillus polymyxa*, *Paenibacillus mucilaginosus*, *Priestia megaterium*, *Rhizobium*, *Pseudomonas chlororaphis*, *Bacillus thuringiensis*, *Trichoderma viride*, *Metarhizium anisopliae*, *Beauveria bassiana*, etc.

Acknowledgment

This work was supported by the Higher Education and Science Committee of RA (Research project № 23EDP-4D003).

Tailoring diverse microbial consortia for the improvement of plant nutritional and nutraceutical properties

Sbrana, Cristiana¹; Giorgetti, Lucia¹; Fiorentino, Sofia¹; Chiellini, Carolina¹

¹ CNR-Institute of Agricultural Biology and Biotechnology, Pisa, Italy

The positive impact of microbial consortia on plant growth and health support their use as bioenhancers. The improvement of chicory (*Cichorium intybus* L.) and lettuce (*Lactuca sativa* L.) nutritional value, in terms of nutrient uptake and accumulation of health-promoting compounds, was studied using an in vivo whole-plant system, in different plant genotypes and microbial consortia (MC). The MC were selected on the basis of the occurrence of bacterial strains showing plant growth promoting (PGP) traits in vitro, and of arbuscular mycorrhizal fungi (AMF) capable of enhancing chicory nutrition. The experimental system used allowed the collection of both plant and fungal tissues to determine biomasses and concentrations of nutrients. Moreover, polyphenols, photosynthetic pigments and fructooligosaccharides content and antioxidant activity were assessed in plant tissues. The analysis of nutrients distribution showed that MC often significantly increased iron and zinc contents in plant leaves, and that some micronutrients, namely copper and iron, were accumulated by extraradical mycelium of AMF. The biosynthesis of polyphenols and flavonoids, as well as the antioxidant activity, were enhanced by MC in lettuce and chicory leaves, although differential responses were obtained depending on plant genotypes. The content of fructooligosaccharides, which varied with plant genotypes and MC used, was significantly increased in the chicory var. "Variegata di Castelfranco", resulting in the accumulation of health-promoting molecules in roots. Overall, data support the use of MC for plant biofortification and for the enhancement of nutritional and nutraceutical value of plant-derived food.

Session 3 – Pathogenic and beneficial aspects of microbes in human and animal health

Thursday 19th September, 9:00am

Chairs: Hurtado Ortiz Raquel (France), Susca Antonia (Italy).

Studying host-microbe interactions in health and disease using germ-free and gnotobiotic mouse technology

Jans, Maude^{1,5}; Thiran, Alexandra^{1,3,5}; Kolata, Magdalena^{6,7}; Andries, Vanessa^{1,3,5}; Blancke, Gillian^{1,3,5}; Ciers, Maarten^{1,5}; Thorp, Marie^{1,3,5}; Petta, Ioanna^{1,3,5}; Diependaele, Louise^{1,3,5}; Berx, Geert^{2,4,5}; Elewaut, Dirk^{1,3}; van Loo, Geert^{1,2,4,5}; Remaut, Han^{6,7}; Vereecke, Lars^{1,3,5}

¹ VIB Center for Inflammation Research, B-9052 Ghent, Belgium; ² Department of Biomedical Molecular Biology, Ghent University, B-9052 Ghent, Belgium; ³ Department of Internal Medicine and Pediatrics, Ghent University, Ghent, Belgium; ⁴ Ghent Gut Inflammation Group, Ghent University, Ghent, Belgium; ⁵ Cancer Research Institute Ghent (CRIG), 9000 Ghent, Belgium; ⁶ Structural Biology Brussels, Vrije Universiteit Brussel, Brussels, Belgium 12; ⁷ Structural & Molecular Microbiology, VIB-VUB Centre for Structural biology, Brussels, Belgium

The intricate relationship between hosts and their resident microbiota is pivotal to understanding health and disease. Germ-free (GF) and gnotobiotic mouse models are indispensable for elucidating these complex interactions. The Host-Microbiota-Interaction lab (HMI) headed by Prof. Lars Vereecke employs these technologies to uncover mechanisms underlying inflammatory diseases and cancer. We explored the presumed mechanistic link between intestinal inflammation and arthritis, known as the 'gut-joint axis', using transgenic mouse models of gut and joint inflammation [1] Remarkably, germ-free conditions completely prevented intestinal inflammation while joint inflammation persisted, highlighting the microbiota's critical role in gut pathology but not in joint inflammation. These findings suggest that while microbiota targeted therapies could be effective for intestinal inflammatory diseases, joint inflammation may require alternative therapeutic approaches. Further, using unique transgenic models of microbiota dependent colorectal cancer (CRC), we discovered that genotoxic pks+ *E. coli* strains exacerbate colorectal cancer through adhesion-mediated epithelial binding [2]. Our ongoing research employs these transgenic mouse models to identify and characterize other CRC-associated 'oncobacteria'. Understanding the mechanisms by which these microbes drive disease could lead to targeted therapies that do not disturb the resident microbiota. Collectively, these studies underscore the profound impact of host-microbe interactions on disease progression and health. GF and gnotobiotic mouse technologies are essential for disentangling these interactions, enabling precise studies of microbial influences on various diseases. They pave the way for innovative therapeutic and preventative strategies, leveraging microbiota to improve health outcomes globally.

References

- [1] Thiran, A. et al. Sterile triggers drive joint inflammation in TNF and IL -1-dependent mouse arthritis models . EMBO Mol. Med. (2023) doi:10.15252/emmm.202317691
- [2] Maude Jans, Magdalena Kolata, Gillian Blancke, Maarten Ciers, Mozes Sze, Alexandra Thiran, Geert Berx, Sabine Tejpar, Geert van Loo, Iliyan D. Iliev, Han. Remaut & Lars Vereecke. Colibactin-induced genotoxicity and colorectal cancer exacerbation critically depends on adhesion-mediated epithelial binding. bioRxiv (2023). doi:https://doi.org/10.1101/2023.08.16.553526

Using DNA Metabarcoding as a Non-Invasive Tool for the Conservation of the Critically Endangering Seychelles Bat *Coleura seychellensis*

Sandri, Camillo¹; Borruso, Luigimaria²; Mimmo, Tanja²; Modesto, Monica³; Scarafile, Donatella³; Francati, Santolo³; Mattarelli, Paola³; Joubert, Flavien⁴; Ashley, Dias⁵; Dindo, Maria Luisa³; Spiezio, Caterina¹

¹ Department of Animal Health Care and Management, Parco Natura Viva Garda Zoological Park, Bussolengo, Verona, Italy; ² Faculty of Science and Technology, Free University of Bolzano-Bozen, Bolzano-Bozen, Italy; ³ Department of Agricultural and Food Sciences, University of Bologna, Bologna, Italy; ⁴ Department of Biodiversity Conservation at Ministry of Agriculture Climate Change and Environment; ⁵ Seychelles Parks and Gardens Authority, Mahé, Seychelles

Bats constitute a quarter of all mammal species, with many globally threatened or near threatened. *Coleura seychellensis*, an insectivorous bat from the Microchiroptera suborder, is endemic to the Seychelles, restricted to Silhouette and Mahé islands. Once widely distributed, this species is now classified as Critically Endangered. Key factors contributing to its decline include human population growth leading to the loss of lowland forests, clearing of coconut plantations, and habitat modifications due to invasive plants, which have reduced insect populations. Additionally, shifts in the insect community may have disrupted the bat's dietary habits. Accurate and comprehensive knowledge of the feeding habits and ecology of *C. seychellensis* is crucial for understanding its ecological requirements, assessing how food availability affects its population status, and identifying key resources for designing effective management strategies. It has to be considered that diet plays a significant role in shaping gut microbiota, which in turn impacts host health. The present work was aimed at studying the fecal samples of bats in two different seasons with the purpose of (a) describing the bacterial and fungal groups of gut microbiota and (b) investigating the insectivorous diet of *C. seychellensis*. A plastic film was placed under the bat colony in the cave and left for 1 day. Fecal pellets were sampled on the top of the plastic film and placed in DNA/RNA Shield reagent (Zymo). DNA from six pooled samples was extracted using QIAamp PowerFecal Pro DNA Kit (Qiagen). The *C. seychellensis* microbiome and diet were investigated via DNA metabarcoding of the bacterial 16S rRNA gene, fungal ITS2 region and arthropod COI gene. When correlated with knowledge of gut microbial composition, the results obtained from insect-based diet reconstruction could assist in utilizing and addressing microbiome research to conserve endangered species.

Riboflavin overproducing food-grade bacteria: genomic basis, biotechnological applications and perspectives

Capozzi, Vittorio¹; Scauro, Angela²; Spano, Giuseppe²; Fragasso, Mariagiovanna²; Russo, Pasquale³

¹ National Research Council of Italy Institute of Sciences of Food Production (ISPA), Foggia, Italy; ² Department of Agricultural Sciences, Food, Natural Resources and Engineering, University of Foggia, Foggia, Italy; ³ Department of Food, Environmental and Nutritional Sciences, University of Milan, Milan, Italy

Food microbiota is an important reservoir of microorganisms capable of increasing the content of essential micronutrients, including, among the vitamins of the B-group, riboflavin (vitamin B₂), the precursor of the flavin mononucleotide (FMN) and flavin adenine dinucleotide (FAD). In Gram-positive bacteria the expression of the rib operon for riboflavin biosynthesis is regulated by an FMN-riboswitch consisting of a sensitive domain (aptamer) which, binding to the effector (FMN), induces a conformational change in the regulatory domain. Exposure of vitamin B₂-prototrophic strains to roseoflavin (a toxic analogue of riboflavin) allows the selection of spontaneous mutants able to overproduce riboflavin and harbour mutations in the rib operon riboswitch. In the last twenty years, this strategy allowed the selection of food-grade strains belonging to different species well-known for their technological and/or functional potential. These bacteria are associated with different food matrices and, due to their high metabolic versatility, have been suggested for the sustainable in situ riboflavin bio-fortification of a wide range of fermented foods. Moreover, several riboflavin-overproducing strains have been comprehensively characterised for their probiotic potential, and their administration has been reported to revert ariboflavinosis, as well as to mitigate intestinal inflammatory diseases and some cancers and provide neuroprotective effects. Finally, we discuss preliminary results on future efforts, such as i) to select new strains from spontaneous fermentation and/or unconventional food matrices, ii) to optimise adequate food processing, iii) to elucidate the regulation of the rib operon, iv) to analyse the microbial tolerance against oxidative stress, v) to better understand the beneficial effect of the consumption of B₂-fortified foods and/or B₂-overproducing probiotics, vi) to exploit strategies to alleviate micronutrient malnutrition in developing countries.

Acknowledgments

We are exploring the potential of this driver of innovation in the framework of the ongoing projects i) the European Union Next-Generation EU [Piano Nazionale di Ripresa e Resilienza (PNRR)—Missione 4 Componente 2, Investimento 1.4—D.D. 1032 17/06/2022, CN00000022] within the Agritech National Research Centre for Agricultural Technologies and ii) the NextGeneration EU [PNRR], in the framework of the Mission 4 Component 2 Investment 1.3-Award Number: Project code PE00000003, Project title: “ON Foods-Research and innovation network on food and nutrition Sustainability, Safety and Security—Working ON Foods”. Vittorio Capozzi was partially funded by CNR project “NUTRAGE FOE-2021 DBA.ADO05.225”.

Defining the *Enterobacter cloacae* species complex, with particular emphasis on *Enterobacter hormaechei*.

RAHI, Praveen¹; Muhle, Estelle¹; Criscuolo, Alexis²; Clermont, Dominique¹

¹ Collection of Institut Pasteur (CIP), CRBIP, Département de Microbiologie, Institut Pasteur, Paris, France; ² Institut Pasteur, Bioinformatics and Biostatistics Hub, Université de Paris, Paris, France

Accurate species-level identification within the *Enterobacter cloacae* complex (ECC) is crucial, as this heterogeneous group is often implicated in nosocomial outbreaks. This precision is essential for microbial bioresource centers (mBRCs), which provide authentic microbial strains and associated information to researchers. Various genomic features, including core-gene phylogenies and average nucleotide identity (ANI), have been used as reliable metrics for defining prokaryotic species. However, the lack of clarity in species definitions often blurs the boundaries, particularly among closely related taxa like ECC members. Recently, *Enterobacter hormaechei* has undergone several taxonomic revisions, revealing a diversity too broad for a single species classification. We sequenced genomes of 12 ECC strains from the Collection of Institut Pasteur (CIP). Phylogenetic analysis based on concatenated sequences of 120 universal (core) genes and pairwise ANI comparisons identified strains corresponding to *E. asburiae*, *E. cloacae*, *E. kobei*, and *E. ludwigii*, alongside distinct clusters of *E. hormaechei*, "*E. hoffmannii*," and "*E. xiangfangensis*." Notably, the latter two are not formally recognized species but are considered subspecies of *E. hormaechei*. Our comprehensive analysis, which included genome sequences of type strains from all *Enterobacter* species and 667 representative strains of *E. hormaechei*, facilitated the circumscription of *E. hormaechei*, "*E. hoffmannii*," and "*E. xiangfangensis*." Additionally, we identified various genes associated with antimicrobial resistance, including ACT type -lactamase, and virulence factors among the ECC strains. In summary, this study establishes accurate taxonomy for ECC strains and provides strong evidence to delineate species boundaries for *E. hormaechei*, "*E. hoffmannii*," and "*E. xiangfangensis*". It also highlights the utility of population-based approaches, such as genetic discontinuity within the populations in substantiating these.

Functionalization of chickpeas through *Saccharomyces cerevisiae* var. *boulardii*

Accettulli, Alessandra¹; Corbo, Maria Rosaria¹; Sinigaglia, Milena¹; Altieri, Clelia¹; Bevilacqua, Antonio¹

¹ University of Foggia, Foggia, Italy

Legumes occupy an important nutritional place due to their wide range of proven beneficial properties; in fact, they provide essential nutrients to our bodies by participating in the eubiotic state of the intestinal microbiota [1]. Probiotic microorganisms are key factors in modulating the microbiota toward eubiosis, and counteract the dysbiosis, that is the disruption of the state of equilibrium that results in the development of various diseases [2]. Therefore, the aim of this study is to design a new combination probiotic/matrix. Chickpeas, due to a greater surface area available, were chosen for the project, while *Saccharomyces cerevisiae* var. *boulardii* was used as probiotic. In a first step, adhesion experiments were performed to point out the conditions for a better attachment of yeasts to chickpea surface; while in a second step, the product was thermally treated and then used as a prebiotic-like matrix to assess a potential bifidogenic effect. Finally, a panel test was done. Yeasts were able to adhere at high levels to legume surface, while the addition of thermal treated chickpeas with yeasts exerted a protective effect on *Bifidobacterium* spp. during the death phase; the organoleptic scores appeared improved.

References

- [1] Maphosa, Y., & Jideani, V. A. (2017). The role of legumes in human nutrition. In: Functional food-improve health through adequate food, Intech Open, DOI: 10.5772/intechopen.69127.
- [2] Al-Rashidi, H. E. (2022). Gut microbiota and immunity relevance in eubiosis and dysbiosis. Saudi Journal of Biological Sciences, 29(3), 1628-1643.

Growth inhibition of Human Pathogenic Microorganisms (HPMOs) by endophytic and rhizosphere bacteria

Marangi, Matteo¹; Szymańska, Sonia¹; Kęsy, Jacek¹; Hryniewicz, Katarzyna¹

¹ Nicolaus Copernicus University

In the last decade, the consumption of raw vegetables contaminated with human pathogenic microorganisms (HPMOs) has increased due to poor agricultural processes [1]. The most commonly identified HPMOs were *E. coli*, *Klebsiella spp.* and *Salmonella spp.* and the main vegetable where they were detected was lettuce [1]. The microbial Volatile Organic Compounds (mVOCs) can inhibit the growth of HPMOs. *Bacillus spp.* [2] and *Pseudoalteromonas sp.* [3] were observed to synthesize VOCs against respectively *B. cereus* and *S. aureus*. According to our hypothesis, plant endophytic and rhizosphere soil bacteria can synthesize VOCs that have inhibitory effect on the growth of HPMOs. The main objective of our work was to select the most efficient bacterial strains to inhibit HPMOs growth. For this purpose, we tested 58 endophytic and rhizosphere bacteria from classes Bacilli, Gammaproteobacteria, Actinomycetes and Alphaproteobacteria (previously isolated from *Salicornia europaea L.*) growing in co-culture with *Listeria monocytogenes*, *Salmonella enterica* and *Escherichia coli* on bipartite Petri plates. The inhibitory effect of the bacteria was calculated based on the growth of HPMOs compared to the control (bacterial growth in monocultures) after 7 days of culture. A statistically significant inhibitory effect on the growth of HPMOs was shown for 8 rhizosphere soil and 8 endophytic bacteria. Strains that showed high growth inhibition of HPMOs, were selected for solid-phase microextraction mass spectrometry analysis in the headspace (HS-SPME-GC-MS). The HS-SPME-GC-MS revealed the synthesis of some mVOCs with potential anti-microbial activity. Concluding, our study revealed that 16 endophytic and rhizosphere bacteria inhibited significantly the growth of HPMOs and that some of them synthesized mVOCs with potential anti-microbial activity.

References

- [1] Rahman M., Alam M. U., Luies S. K., Kamal A., Ferdous S., Lin A., Sharior F., Khan R., Rahman Z., Parvez S. M., Amin N., Hasan R., Tadesse B. T., Taneja N., Islam M. A., Ercumen A. Intern. Jour. of Environ. Res. and Pub. Health. 2021.
- [2] Ajilogba C. F., Babalola O. O. World Jour. of Micro. & Biotech. 2019.
- [3] Garrido A., Atencio L. A., Bethancourt R., Bethancourt A., Guzmán H., Gutiérrez M., Durant Archibold A. A. Antib. 2020.

Acknowledgments

This research was conducted as a part of the SaliFood project funded by the National Science Centre (NCN, Poland) (UMO-2019/33/B/NZ9/02803).

Supercritical fluid extraction of torularhodin from a new red yeast strain

Larocca, Vincenzo¹; Ambrico, Alfredo¹; Martino, Maria¹; Trupo, Mario¹; Magarelli, Rosaria Alessandra¹; Spagnoletta, Anna¹

¹ ENEA, Department for Sustainability, Trisaia Research Center, Rotondella (MT), Italy

This study provides valuable informations for recovery of microbial pigments from microorganisms for human and animal health purposes. Some red yeasts, especially by *Rhodotorula* and *Sporobolomyces* genera are potential sources from which to obtain carotenoids. In this work a new strain of *Rhodotorula* (ELP2022), currently present in the ENEA Microbial Culture Collection (EMCC), was isolated from fresh cheese and characterized for its metabolic pattern (Biolog, Inc) and carotenoids profile. The main carotenoids identified were: -carotene, -carotene, torulene and torularhodin. The latter is a dark pink coloured carotenoid belonging to the xanthophylls group and the growing interest in this molecule is due to its biological activities such as antioxidant, anticholesterolemic, anti-inflammatory, antimicrobial and anticancer. In this context the strain ELP2022 was used to develop a new efficient method for selective extraction of torularhodin in two sequential steps by applying the extraction technique with supercritical CO₂ (CO₂-SFE) and the method has been the subject of an Italian patent deposit (n. 102023000018729). In particular, it was grown in bench-scale fermenters using glucose-based mineral media. The pre-treated and dried biomass was subjected to a CO₂ extraction process in supercritical conditions (CO₂-SC) by a bench scale system (Applied Separations Spe-ed SFE-2). After extraction the torularhodin represented no less than 95.2% of the total carotenoids in the red extracts obtained from the second step. Therefore, this method allows to separate torularhodin from other carotenoids without the use of chromatographic techniques and doesn't involve the use of solvents potentially dangerous for health and environment. So, this approach avoids the presence of residual solvents of class 1 and 2 in the extracts, which should be respectively not employed or limited in pharmaceuticals according to ICH guidelines Q3C(R8).

Acknowledgments

This work was granted by the European Commission NextGenerationEU within the framework of the National Recovery and Resilience Plan (Mission 4 "Education and Research", Component 2, Investment 3.1), Project "Strengthening the MIRRI Italian Research Infrastructure for Sustainable Bioscience and Bioeconomy" (SUS-MIRRI.IT), code IR0000005.

Interaction between probiotic lactic acid bacteria and immunosuppressive drugs

Di Renzo, Tiziana¹; Reale, Anna¹; Cataldi, Mauro²

¹ Institute of Food Sciences, National Research Council (CNR-ISA), Avellino, Italy; ² Section of Pharmacology, Department of Neuroscience, Reproductive Sciences and Dentistry, Federico II University of Naples

The human microbiota consists of trillions of microorganisms living in a commensal state in different niches of the body, of which the gut is the most studied. In some specific diseases, important changes in the composition of the microbiota occur as a consequence of the disease itself or the therapy taken by the patient. A prototypical example of such a clinical condition is the kidney transplant. After kidney transplantation, important changes in the composition of the gut microbiota have been observed, with an increase in Firmicutes, Proteobacteria and Bacteroidetes. These changes can lead to infectious complications and influence the synthesis of uremic toxins, as well as alter the response to immunosuppressive drugs commonly used during kidney transplantation therapy. Studies in the literature show that the use of probiotics can modify and/or improve the state of intestinal dysbiosis. Therefore, the aim of this study was to evaluate the interactions between different probiotic lactic acid bacteria and immunosuppressive drugs commonly administered in post-transplant therapy. In detail, different strains of probiotic lactic acid bacteria belonging to the species *Lactiplantibacillus plantarum*, *Lacticaseibacillus rhamnosus*, *Lactobacillus acidophilus*, *Lacticaseibacillus casei*, *Lacticaseibacillus paracasei*, *Limosilactobacillus reuteri*, *Streptococcus thermophilus* and immunosuppressive drugs Mycophenolate Mofetil (MMF), Mycophenolic Acid (MPA) and Mycophenolic Acid Glucuronide (MPAG) were investigated. The effect of the bacterial-drug interaction was assessed by analysing the viability of the microbial species during an incubation time of 24h at 37°C under anaerobic conditions. Furthermore, the ability of probiotic lactic acid bacteria to metabolise and/or accumulate immunosuppressive drugs in vitro was assessed by HPLC analysis.

Acknowledgments

This research was funded by the Italian Ministero della Salute, Piano Nazionale di Ripresa e Re-silienza (PNRR) Missione 6—Componente 2 Investimento 2.1 Valorizzazione e Potenziamento della Ricerca Biomedica del SSN, grant number PNRR-MAD-2022-12376812 “Presystemic metabolism of immunosuppressant drugs by gut microbiota as a risk factor for kidney transplant rejection: mitigation strategies”.

Factors affecting the production of γ -aminobutyric acid in *Levilactobacillus brevis* LB12

Lavanga, Emanuela¹; Giavalisco, Marilisa¹; Aliano, Anita¹; Ricciardi, Annamaria¹; Parente, Eugenio¹; Zotta, Teresa¹

¹ Department of Agricultural, Forestry, Food and Environmental Sciences, University of Basilicata, Potenza, Italy

γ -aminobutyric acid (GABA) provides several physiological benefits to human health [1,2]. Lactic acid bacteria (LAB) are recognized as potential GABA producers and several strains, mainly belonging to the species *Levilactobacillus brevis*, have been investigated and characterized for their ability to produce GABA in synthetic media and/or food matrices [1,2].

In this study, the effect of different pH (3.5, 4.0, 4.5, 5.0, 5.5, 6.0; buffer system), glutamate concentration (0, 1, 2, 5, 10 g/L of monosodium glutamate, MSG; buffer system) and atmosphere of incubation (cultivation and adaptation to anaerobic and aerobic conditions; AN vs AE) was evaluated on the GABA production in *Lvb. brevis* LB12. Glutamate (Glu) consumption and GABA accumulation was estimated with Thin Layer Chromatography and densitometric spot analysis [3]. The expression of GAD operon genes (*gadB/gadA*; glutamate decarboxylase, *gadC*, glutamate: gamma-aminobutyrate antiporter; *gadR*, transcriptional regulator, *gltX*, glutaminyl-tRNA synthetase), in response to the atmosphere of incubation and Glu supplementation, was evaluated by using RT-qPCR (SYBR Green protocol). The optimal pH for Glu uptake and GABA production was 4.5, while the greatest inductive effect of Glu supplementation was observed with 10 g/L MSG. AE conditions increased biomass yield, but impaired the efficiency of Glu to GABA biotransformation, resulting in a lower GABA accumulation, in both culture supernatant and reaction buffer. The expression of genes belonging to GAD operon (*gadB*, *gadC*, *gadR*) was also impaired in AE growing cells. *gadA* gene, instead, was affected to a lesser extent by atmosphere of incubation and Glu supplementation, confirming that *gadB* was mainly involved in Glu-GABA conversion. This work provides further insights on the factor affecting GABA production and regulation mechanisms of GAD operon, although additional investigations are needed to maximize the efficiency of Glu-GABA conversion in *Lvb. brevis* LB12.

References

- [1] Diana M, Quílez J, Rafecas M. 2014. doi: 10.1016/j.jff.2014.07.004
- [2] Xu N, Wei L, Liu J. 2017. doi:10.1007/s11274-017-2234-5
- [3] Zotta T, Faraone I, Giavalisco M, Parente E, Lela L, Storti L. V, Ricciardi A. 2022. doi: 10.3390/microorganisms10112184

Acknowledgments

This work was partially funded by the European Commission – NextGenerationEU, Project “Strengthening the MIRRI Italian Research Infrastructure for Sustainable Bioscience and Bioeconomy”, code n. IR0000005.

Comparison of microbial populations of peritonsillar abscess determined by molecular and bacteriological methods

Rööp, Tiiu¹; Štšepetova, Jelena¹; Vaikjärv, Risto²; Saar, Merili¹; Kasenõmm, Priit³; Sepp, Epp¹; Mändar, Reet¹

¹ Department of Microbiology, University of Tartu, Tartu, Estonia; ² Ear-Nose-Throat Clinic, Tallinn, Estonia; ³ Department of Microbiology, University of Tartu, Tartu, Estonia; ³ Ear Clinic, Tartu University Hospital, Tartu, Estonia

Peritonsillar abscess (PTA) is a severe deep neck space infection with an insufficiently characterized bacterial etiology. We aimed to reveal bacteria associated with PTA applying next generation sequencing (NGS) and traditional bacteriology.

The study group consisted 113 consecutive patients with PTA. Pus and tonsil biopsies were investigated. Semi-quantitative bacteriological methodology was used for the examination of all samples. Colonies were identified by MALDI-TOF mass spectrometry. NGS method was additionally applied to samples of 91 PTA patients. Over 400 genera and 800 species belonging to 34 phyla were revealed by NGS. The most abundant species in both sample types were *Streptococcus pyogenes*, *Fusobacterium necrophorum* and *Fusobacterium nucleatum*. Other species displaying a high mean relative abundance were *Prevotella oris* in all samples; *Porphyromonas endodontalis* in pus and *Streptococcus mitis* in tonsils. *S. pyogenes* and *F. necrophorum* were the predominant species (> 10% in a community) in 31% pus samples, while *F. nucleatum* in 23% and *S. anginosus* in 9% pus samples. In total, 174 bacterial isolates were identified, which belonged to 5 phyla, 25 genera and 62 species. In the tonsil biopsies, the most frequent isolates were *Streptococcus parasanguinis* and *Streptococcus pneumoniae* (both 10.5%), followed by *Neisseria spp* (16.6%) and *Actinomyces spp* (7%). In pus samples, the most frequently isolates were *Streptococcus spp* (48.3%). *S. pyogenes* and *F. necrophorum* was isolated from both type samples while *F. nucleatum* only from pus. 156 bacterial strains were deposited to the Human Microbiota Biobank (HUMB).

The most probable causative agents of PTA are *S. pyogenes*, *F. necrophorum* and *F. nucleatum*. Some streptococci (*S. anginosus*, *S. parasanguinis*) and anaerobes (*Prevotella*, *Porphyromonas*) may contribute to the infection as well. Pus of the abscess is more representative specimen for examination. All deposited strains are available for future research purposes.

Acknowledgments

We thank the staff of Tartu University Hospital for technical assistance. This study was supported by the Estonian Ministry of Education and Research (target financing No. SF0180132S08, institutional research funding IUT 34-19, and funding of scientific collections KOGU-HUMB) and the University of Tartu (grant No. SARMBARENG).

Use of potential probiotic autochthonous *L. plantarum* as co-starter for the production of goat's milk cheeses

Pisano, Maria Barbara¹; Viale, Silvia¹; Deplano, Maura¹; Civolani, Eleonora¹; Cosentino, Sofia¹

¹ University of Cagliari, Cagliari, Italy

Raw milk and artisanal cheeses are characterized by a microbiota mostly constituted of starter and non-starter lactic acid bacteria (NSLAB), the latter playing a key role in the ripening and flavour development of the cheeses. Mesophilic lactobacilli are the predominant group in the microbiota of NSLAB and are endowed with probiotic potential. Cheese has been proven to be an optimal carrier product to deliver living probiotic bacteria, and autochthonous strains with potential probiotic features would be the best choice for use as adjunct cultures since they should be well-adapted to this food environment. In this study three autochthonous *Lactiplantibacillus plantarum* strains possessing several probiotic and technological properties were used as co-starter cultures in the manufacturing of goat's milk cheeses. The aim of the study was to evaluate the presence and viability of these strains during maturation and ripening of goat's milk cheeses produced at industrial scale-level, using both conventional culturing and PCR-DGGE culture-independent technique. Compositional, microbiological, and sensory analyses were also carried out to evaluate the impact of the *L. plantarum* strains on the cheese's quality. The results showed that lactobacilli count evolved similarly in the cheeses produced with the three autochthonous strains, maintaining high viability until the end of ripening (>9 log₁₀ cfu/g in 30-days-cheeses). DGGE profiles showed a band corresponding to *L. plantarum* in the cheese samples throughout the whole ripening period. No significant differences in the physico-chemical parameters were found between the cheeses produced with the three different autochthonous cultures. Sensory evaluation indicated that the use of the autochthonous potential probiotic co-starter did not adversely affect the overall acceptability of the goat cheeses, however, those produced with the strain *L. plantarum* 11/20966 received significantly higher scores for odour, aroma, and texture.

Acknowledgments

This work has been funded by European Agricultural Fund for Rural Development Program 2014- 2020 and partially supported by "MIRRI Italian Research Infrastructure for Sustainable Bioscience and Bioeconomy", granted by NextGenerationEU, National Recovery and Resilience Plan (NRRP), Mission 4 "Education and Research", Component 2, Investment 3.1 – Project Code IR0000005.

Session 4 – Microbial life in extreme habitats: a 21st century challenge

Thursday 19th September, 11:00am

Chairs: Varese Giovanna Cristina (Italy), Sedláček Ivo (Czech Republic).

Diversity of microalgae in low pH environments: fifty years of the ACUF Collection, from strain to microbiome conservation and exploitation.

Pollio, Antonino¹

¹ Department of Biology, University of Naples Federico II, Italy

ACUF (Algae Collection of the University of Naples Federico II) was founded in 1974 as a group of strains collected by Professor Taddei and his collaborators to study the ecophysiology of the thermoacidophilic unicellular red alga *Cyanidium caldarium*. The strains were initially collected from the Italian peninsula and Sicily, in low pH environments (hot springs, fumaroles, sulphur mines) and have been used in numerous physiological, ecological and morphological research projects. Since 1978, the ACUF has held more than 300 *C. caldarium* strains and was established as a facility of the Istituto di Botanica of the University of Naples. In the following decade, new accessions were deposited by other researchers from the Istituto di Botanica, who explored low pH environments in America and Asia in search of Cyanidiales populations. Since then, ACUF members have visited other countries on different continents, extending the sampling to different types of extreme environments, such as dry rocks and caves. ACUF is a member of the World Federation for Culture Collections and an associate member of the Italian Microbial Resource Research Infrastructure (MIRRI-IT). The functions of ACUF are to collect and maintain in culture isolates of algae and cyanobacteria, mainly from extreme environments, to serve as a repository and to make these isolates and related information available for research and biotechnology applications. The ACUF collection contains 112 genera and about 1000 isolates, including cyanobacteria. Stocks of most cultures are maintained as live strains, but part of the isolates are also maintained in nongrowing conditions, frozen with or without cryoprotectants. Under the framework of the PNRR project SUS-MIRRI, management practices to maintain and cultivate microbiomes from low pH environments are in progress, as well as co-culture mass cultivations, in which bacteria and microalgae collected from the same site are grown together to improve productivity and biosynthesis of valuable compounds.

Life at the limits: diversity, adaptation strategies and bioprospecting of microbes living in Arctic deep-sea habitats (INDEPTH)

Kaczorowska, Anna-Karina¹; Ida Helene Steen²; Runar, Stokke²; Przemysław, Decewicz³; Wojciech, Rypniewski⁴; Łukasz, Dziewit³; Kaczorowski, Tadeusz⁵

¹ *Collection of Plasmids and Microorganisms / KPD, Faculty of Biology, University of Gdansk, Gdansk, Poland;* ² *Department of Biological Sciences, Center for Deep Sea Research, University of Bergen, Bergen, Norway;* ³ *Department of Environmental Microbiology and Biotechnology, Faculty of Biology, University of Warsaw, Warsaw, Poland;* ⁴ *Institute of Bioorganic Chemistry, Polish Academy of Sciences, Poznań, Poland;* ⁵ *Laboratory of Extremophiles Biology, Faculty of Biology, University of Gdansk, Gdansk, Poland*

The deep-sea biosphere represents the largest living system on Earth, distinctive in its unparalleled complexity. This vast biosphere, often referred to as life's last frontier, is characterized by darkness, high pressure, limited nutrients, and a wide range of temperatures from low (2-4°C) in most habitats to extremely high (above 100°C) in the seabed hot hydrothermal vents. The INDEPTH project aimed to decipher the hidden reservoir of metabolic traits of the unique biodiversity at the Arctic Mid-Ocean Ridge (AMOR) vents. By reconstructing and analyzing AMOR microbial genomes, the potential for homoacetogenesis via the Wood-Ljungdahl pathway (WLP) was elucidated in the archaeal Korarchaeia, supporting previous hypotheses that the WLP has evolved independently from the methanogenic metabolism in Archaea [1]. In parallel, a large metagenomic dataset, comprising individual assemblies from 14 samples from Loki's Castle, Perle and Bruse, and Soria Moria vent fields, formed the foundation for developing a novel biodiscovery pipeline to identify novel enzymes with unique activities [2]. The project targeted DNA processing enzymes, proteases, cellulases, betagalactosidases, laccases, alkaline phosphatases, and chitinases. As a result, more than 30 novel enzymes have been thoroughly characterized at both structural and functional levels [3]. The INDEPTH project expanded our understanding of biodiversity, evolution and the functioning of complex microbial communities, including adaptation strategies required for survival in the extreme conditions of seafloor hot vents. It also led to the development of new tools for metagenomics that enhance bioprospecting and deepen our understanding of microbial life in extreme Arctic habitats.

References

- [1] Vulcano F et al.. 2023. Potential for homoacetogenesis via the Wood-Ljungdahl pathway in Korarchaeia lineages from marine hydrothermal vents. *Environ Microbiol Rep.* 15:698-707.
- [2] Ciuchcinski K et al.. 2024. Rapid discovery of novel enzymes across diverse environments using a hybrid (Meta)genomic approach. In preparation.
- [3] Røyseth V et al.. 2023. Activation mechanism and activity of globupain, a thermostable C11 protease from the Arctic Mid-Ocean Ridge hydrothermal system. *Frontiers in Microbiology* 14:1199085.

Acknowledgments

INDEPTH was funded by The Norway Financial Mechanism through the NCN (Poland) GRIEG1 grant: UMO-2019/34/H/NZ2/00584.

Cold-adapted carboxylic ester hydrolases from two Antarctic *Psychrobacter* strains: genomic analyses and in-vitro studies

Cattaneo, Andrea^{1,2}; Papikyan, Razmik³; Leone, Luigi²; Vittoria, Maria²; Romano, Ida²; Di Donato, Paola⁴; Poli, Annarita²; Finore, Ilaria²

¹ Ca' Foscari University of Venice, Department of Environmental Sciences, Informatics and Statistics (DAIS), Mestre (VE), Italy; ² National Research Council of Italy (CNR), Institute of Biomolecular Chemistry (ICB), Pozzuoli (NA), Italy; ³ A.L. Takhtajan Institute of Botany, Acharyan 1, 0060, Yerevan, Armenia; ⁴ Parthenope University of Naples, Department of Science and Technology (DiST), Napoli (NA), Italy

Pollution from polyester plastics is increasingly affecting biotic and abiotic components of the ecosystems, making it necessary to find effective solutions. The application of bacterial enzymes-mediated degradation is a promising strategy to treat plastic waste. Therefore, extremophiles and extremozymes could be applied in bioremediation due to their peculiar features of adaptation to these ecological niches.

Psychrobacter strain ASPA161_6 and strain ASPA161_9 were isolated from marine sediment samples collected in Terra Nova Bay (74°44'854"S, 164°05'223"E; Antarctica), from the Extremophiles Research Group of Biomolecular Chemistry Institute of the National Research Council (CNR), Pozzuoli (IT), and are stored in the Extremophiles Collection (CE-ICB) (www.susmirri-catalog.di.unito.it/collections) part of the SUS-MIRRI project, which collects, characterizes, and distributes microbial strains for research and bio-industrial purposes preserving microbial biodiversity. Their genomes were sequenced and deposited at DDBJ/ENA/GenBank with accession numbers JBEFNI000000000 and JBEFNI000000000, respectively. The digital DNA-DNA hybridization and Average Nucleotide Identity were calculated.

After the functional annotation of the genomes with the RAST Server, *in silico* screenings were performed, highlighting coding sequences related to low-temperature adaption mechanisms and to shock response. Carboxylic ester hydrolases activities and enzymes potentially involved in polyester material degradation were researched. Wet-lab experiments were set up to recover esterase (EC 3.1.1.1) and lipase (EC 3.1.1.3) enzymes from the supernatants of the two strains. The detection of non-specific carboxylic ester hydrolases was done through qualitative agar plate-based assays, while quantitative in-vitro spectrophotometric assays defined the specificity of these enzymes onto classes of substrates, with the outlook to catalyze the depolymerization of plastic under competitive conditions.

Acknowledgment

This research was supported by PNRA19_00073 (TENORE), and PNRA18_00232 (AMICI).

Cold-adapted yeasts: a restricted club of extremophilic organisms

Buzzini, Pietro¹; Ciro, Sannino¹; Gianmarco, Mugnai¹; Luigimaria, Borruso²; Alessandro, Bernetti¹; Daniele, Andreani¹; Benedetta, Turchetti¹

¹ University of Perugia, Perugia, Italy; ² Free University of Bozen-Bolzano, Italy

Cold habitats (average temperature < 5°C) represent over 80% of the Earth's environments. These include the Arctic and Antarctica, high mountains in Asia, Europe, and America, cold deserts, and the deep sea. Although their harsh environmental conditions, cold habitats support both psychrophilic and psychrotolerant fungi (including yeasts), which have developed specific adaptations to cope with low temperatures, e.g. the decrease in growth rates, the synthesis of cold-active enzymes and of cryoprotectant molecules (e.g. cold-shock proteins, sugars, polyols), and an increased fluidity of cell membranes at low temperatures due to the presence of unsaturated fatty acids in their lipid fraction.

The impact of global warming on terrestrial cold habitats is amplifying every year by increasing the duration of ice-free periods. Therefore, studying the diversity of fungal communities (both yeasts and filamentous fungi) occurring in polar and Alpine cold areas may be considered strategic for enhancing our knowledge of the fungal ecology of these understudied niches.

DNA metabarcoding studies (i.e. NGS targeting the ITS fungal region) on fungal communities of soil (including permafrost), brines, ice and debris were conducted to evaluate the taxonomic structure of fungal communities and their ecological interaction with abiotic parameters. Overall, taxa (classified at the genus level) ascribed to Ascomycota dominated among filamentous fungi, while Basidiomycota dominated among yeasts. At the phylotype level, yeasts were predominant in the fungal communities. Alpha and beta-diversity analyses revealed high fungal richness and taxonomic differentiation among samples, even at the small-scale level. Salinity, conductivity, pH and the concentration of organic carbon had a significant ($p < 0.05$) impact on the abundance of some fungal and yeast taxa. Yeasts culturable diversity reflected the results of metabarcoding analysis: an overall dominance of Basidiomycota was confirmed.

Preliminary investigations of alkali-tolerant fungi in stromatolites from Lake Salda (Burdur province, SW Türkiye)

Cecchi, Grazia¹; Tagliasacchi, Ezher²; Di Piazza, Simone¹; Canonica, Laura¹; Akyildiz, Gürçay Kivanç³; Słowakiewicz, Mirosław⁴; Zotti, Mirca¹

¹ Department of Earth, Environment and Life Sciences (DISTAV), University of Genoa, Genoa, Italy; ² Department of Geology, Faculty of Engineering, Pamukkale University, Denizli, Türkiye; ³ Department of Biology, Faculty of Science, Pamukkale University, Denizli, Türkiye; ⁴ Faculty of Geology, University of Warsaw, Warsaw, Poland;

Despite that fungi prefer to colonise neutral or slightly acidic environment; it has been demonstrated that some of them can survive and thrive at high pH. This is due to fungal production of alkaline enzymes like proteases and amylases which protect fungi [1]. These enzymes can be exploited in various industries and used for biotechnological purposes. Therefore, alkaliphilic and alkali-tolerant fungi seem to be of particular interest here.

Lake Salda is a highly alkaline (pH>9) and magnesium-rich lake, located in the Burdur province (SW Türkiye). The lake is one of the deepest enclosed lakes in Türkiye and surrounded by ultramafic and karstified carbonate rocks. Large stromatolitic bodies, fossil and living, built of hydromagnesite, controlled by biological processes, are abundant here. In November 2023, samples from living stromatolites were collected in the littoral zone of Lake Salda. Samples were analysed to obtain their physico-chemical and mycological characteristics. Chemical analysis of lake waters showed presence of 0.21 mg/L PO₄, 0.02 mg/L Fe, 2.4 mg/L NO₃-N, 0.009 mg/L NO₂-N, 84.9 mg/L Cl, 0.33 mg/L NH₃-N, 0.65 mg/L Mg, and 0.18 mg/L Ca, whereas mycological analysis revealed a total of 20 strains, which were isolated from the stromatolite samples and conserved in the culture collection (Cold UNIGE JRU MIRRI IT). Many alkali-tolerant fungal genera and species have been identified such as *Acrostalagmus luteoalbus*, *Alternaria* sp., *Aspergillus templicola*, *Curvularia* sp., *Emericellopsis* sp., *Fusarium equiseti*, and *Stachybotrys cartarum*. Further investigations will be carried out not only to study the potential production of alkaline enzymes and interesting metabolites from these fungi, but also to deepen our understanding of the fungal role in hydromagnesite precipitation and stromatolite-building mats.

References

[1] Fernández-López, M. G., Batista-García, R. A., & Aréchiga-Carvajal, E. T. (2023). Alkaliphilic/AlkaliTolerant Fungi: Molecular, Biochemical, and Biotechnological Aspects. *Journal of Fungi*, 9(6), 652.

From microbial diversity of Campania thermal springs of their maintenance and preservation

Oliva, Felician¹; Selci, Matteo¹; Corso, Davide¹; Licata, Marzia¹; Migliaccio, Flavia¹; Nappi, Nunzia¹; Tonietti, Luca²; Caliro, Stefano³; Cordone, Angelina¹; De Natale, Antonino¹; Pollio, Antonino¹; Giovannelli, Donato¹

¹ Department of Biology, University of Naples "Federico II", Naples, Italy; ² Department of Science and Technology, Parthenope University of Naples, Naples, Italy; ³ Osservatorio Vesuviano, Istituto Nazionale di Geofisica e Vulcanologia, Naples, Italy

The lithosphere is enriched by significant amounts of volatile elements: carbon, oxygen, hydrogen. These elements are often released through subduction zones, where geosphere and biosphere interact with another. The recycling of these volatile substances plays a fundamental role in fueling subsurface microbial communities [1]. Much remains to be understood about the relationship between volatile delivery and microbial composition. The main Italian geothermal activity originates from the subduction of the African and Adriatic plates beneath the Eurasian plate makes the peninsula an excellent area to investigate these relationships. This process generated several secondary geothermal emissions, generating unique, in some cases extreme, geochemical compositions. For this purpose, we analyzed fluid samples collected from 13 geothermal springs present in the Campania region [2]. By combining geochemical and biological data we are able to provide an overview of the physicochemical extremes imposed by the tectonic setting on the subsurface biosphere. Knowing the role of major, minor, and trace elements in influencing the deep biosphere is necessary to the further understanding of the processes linked between the coevolution of life and geology. In turn, this will be helpful to search for the presence of such processes in other small rocky planets with Earth-like tectonic activity. In addition, due to the current technical and physiological limitations of non-cultivating 90% of microorganisms, enhancing preservation and culture methods to maintain and investigate complex communities alive become crucial.

To this end, we are currently engaged in the development of SOPs [3] aimed at microbiome cultivation through the use of the Winogradsky column technique and preservation strategies using different cryoprotective agents coupled with different freezing strategies. This work will lay the bases to constitute the first Italian collection dedicated to microbiomes from extreme environments

References

- [1] Fullerton, Katherine M., *et al.*, "Effect of tectonic processes on biosphere–geosphere feedbacks across a convergent margin." *Nature Geoscience* 14.5 (2021): 301-306.
- [2] Giovannelli, Donato, *et al.*, "Sampling across large-scale geological gradients to study geosphere–biosphere interactions." *Frontiers in Microbiology* 13 (2022): 998133.
- [3] Cocolin, Luca, *et al.*, "STANDARD OPERATING PROCEDURES (SOPs) FOR SAMPLING OF MICROBIOME IN DIFFERENT ECOSYSTEMS." (2023).

The influence of prokaryote communities on *Galdieria sulphuraria* strain maintenance and cultivation

Nappi, Nunzia¹; Cascone, Martina¹; Petraretti, Mariagioia¹; Licata, Marzia¹; Oliva, Feliciano¹; Marra, Luigi²; Cordone, Angelina¹; De Natale, Antonino¹; Giovannelli, Donato¹; Pollio, Antonino¹

1 Department of Biology, University of Naples "Federico II", Naples, Italy; 2 Department of Chemical, Materials and Industrial Production Engineering, University of Naples Federico II, P.le V. Tecchio, Naples, Italy

The Algal Collection at the University of Naples Federico II (ACUF) is a biological resource center that maintains microalgal strains, primarily from Cyanobacteria, Chlorophyta, Rhodophyta, and Bacillariophyceae. A resource of ACUF collection are extremophilic algae mostly belonging to the Cyanidiophytinae isolated from European and non-European sites. The Cyanidiophytinae are a subdivision of the Rhodophyta including species that can survive in thermal and hyperacidic conditions up to 56 °C and a pH lower than 2.0 [1]. They represent the only phototrophic and eukaryotic organisms found in geothermal sites under the above mentioned conditions, in close association with extremophilic bacteria and archaea [2]. Cyanidiophytina members, such as *Galdieria sulphuraria*, are considered the most promising microalgal species for future biotechnological applications (food, health and cosmetics). Interactions between Cyanidiophytinae and prokaryotes cover the full range of symbiotic relationships considered possible. However, most of the interactions between these microalgae and their prokaryotic partner are still poorly studied because separation is a challenging task [3]. This study presents preliminary results from sequencing and culture approaches aimed at exploring the microbial diversity associated with *Galdieria sulphuraria* samples collected at the Mefite d'Ansanto (southern Apennines, Italy). The variation in microbial community composition over time was studied using samples collected in different years. To investigate the interactions between the primary microbial groups and potential metabolic exchanges, selected samples were cultivated under photoautotrophic conditions. Following DNA extraction and sequencing, bioinformatic analyses were conducted to further examine these interactions.

References

- [1] Weber, A., Oesterhelt, C., Gross, W., Brautigam, A., Imboden, L., Krassovskaya, I., Linka, N., Truchina, J., Schneidereit, J., Voll, H., Voll, L., Zimmermann, M., Jamaï, A., Riekhof, W., Yu, B., Garavito, R., & Benning, C. (2004).
- [2] Ferris, M. J., Sheehan, K. B., Köhl, M., Cooksey, K., Wigglesworth-Cooksey, B., Harvey, R., & Henson, J. M. (2005).
- [3] Amin, S. A., Hmelo, L. R., van Tol, H. M., Durham, B. P., Carlson, L. T., Heal, K. R., Morales, R. L., Berthiaume, C. T., Parker, M. S., Djunaedi, B., Ingalls, A. E., Parsek, M. R., Moran, M. A., & Armbrust, E. V. (2015).

Cryopreservation strategies for extremophilic microalgae at ACUF: growth kinetic and photosynthetic assessment

Licata, Marzia^{1,3}; Marra, Luigi^{2,3}; Nappi, Nunzia^{1,3}; Aurino, Elena²; Oliva, Feliciano^{1,3}; Petraretti, Mariagioia⁴; De Natale, Antonino^{1,3}; Giovannelli, Donato³; Pollio, Antonino^{1,3}

¹ ACUF, Algal Collection University of Naples "Federico II", Naples, Italy; ² Department of Chemical, Materials and Industrial Production Engineering, University of Naples "Federico II", Naples, Italy; ³ Department of Biology, University of Naples "Federico II", Naples, Italy; ⁴ Dipartimento di Scienze per gli Alimenti, la Nutrizione e l'Ambiente Università degli studi di Milano Statale

Ex situ conservation allows to preserve genetic materials and functional characteristics of microalgae against environmental and ecological change that could cause the extinction of previously isolated organisms over time [1]. Microalgae conservation is also the first step for any kind of biotechnological exploitation [2]. The present work aims to illustrate the strategies of cryopreservation for extremophile and extreme tolerant strains belonging to The Algal Collection at the University of Naples Federico II (ACUF) within the framework of PNRR_IR SUS-MIRRI.IT project. *Chlorophyta* and *Rhodophyta* are among the largest groups of photosynthetic microbes maintained at ACUF [3] deserving tailored strategies of preservation [4]. We first focused on conventional protocols suitable for cryopreservation of both extreme tolerant and extremophilic microalgal strains, and also tried to find novel methods. Two strains were selected for testing: one belonging to *Stichococcus bacillaris* Nägeli (*Trebouxiophyceae*, *Chlorophyta*), an acid tolerant unicellular green alga commonly cryopreserved, and the other to *Galdieria sulphuraria* (Galdieri) Merola (*Cyanidiophyceae*, *Rhodophyta*), a thermo-acidophilic microalga isolated from an Italian sulphur spring. The extremophilic strains of *Galdieria* have proved difficult to conserve using cryogenic methods. Different cryopreservation strategies applied on two strains belonging to *Stichococcus* and *Galdieria* were compared to evaluate the adaptability of these strains to cryopreservation. A novel variable is introduced to assess viability recovery [5]. Photosynthetic rate and pigments assessment were measured to evaluate hidden metabolic cell damage.

References

- [1] doi: 10.1111/1574-6968.12034.
- [2] doi: 10.1016/j.hal.2008.11.011.
- [3] doi: 10.12775/EQ.2018.023.
- [4] Available: <http://genomics.msu.edu/galdieria>.
- [5] doi: 10.4236/nr.2024.155010.

Fermentation Technologies for the Optimization of levan Production by *Pseudomonas* strain 2ASCA

D'Amodio, Antonia¹; Poli, Annarita¹; Leone, Luigi¹; Vittoria, Maria¹; Romano, Ida¹; Finore, Ilaria¹

¹ National Research Council of Italy (CNR), Institute of Biomolecular Chemistry (ICB), Pozzuoli (NA), Italy

Pseudomonas 2ASCA is an Arctic Gram-negative psychrophilic bacterium isolated from Extremophiles Research Group of Institute of Biomolecular Chemistry of the National Research Council (CNR) in Pozzuoli (Italy) and it is stored in Extremophiles Collection (CE-ICB) (www.susmirri-catalog.di.unito.it/collections) included in the SUS-MIRRI project [1].

Strain 2ASCA produces a homopolymeric exopolysaccharide loosely bound to the cell membrane consisting of a long-chain α -(2,6)-linked fructose polymer, called levan. A crucial aspect of largescale microbial production of exopolysaccharides (EPSs) is the costs associated with production and recovery. For this reason, to optimize the growth of the strain and the hyper-production of levan, different substrates were tested as main carbon sources. Its ability to utilize many sugars as the sole carbon source has driven the search of exopolysaccharide in two different cellular fractions, in loosely and tightly bound fractions (LB and TB, respectively) and in cell-free supernatant fractions (S). Exopolymers obtained from S, TB, and LB fractions were tested for the content of extracellular polysaccharide substance (EPS), proteins, and uronic acids. Among the substrates tested, sucrose supported the best cell growth and was the most suitable carbon source for the induction of the polysaccharide pathway, resulting in the release of the exoproduct consisting of a high percentage of carbohydrates and a low percentage of proteins and uronic acids. Moreover, levan polysaccharide was produced by replacing pure sucrose with agro-industrial waste to achieve sustainability and circular economy objectives [2]. In fact, bacterial fermentation with molasses from sugar beet processing as the sole source of sucrose led to significantly amplified production of EPS. The increase in EPS yield was, most likely, due to the presence of nitrogen and mineral components in the molasses, compared to the sucrose-containing minimal growth medium.

References

- [1] Finore *et al.*, (2020) DOI: 10.3390/microorganisms8091282
- [2] Finore *et al.*, (2024) DOI: 10.1016/j.ijbiomac.2024.131664

Acknowledgment

This work was partially financed by project PRIN 2022LPPFTY, -TREASURE, by PNRR European Commission – NextGenerationEU, Project MICS (Made in Italy – Circular and Sustainable) Extended Partnership MISSIONE 4 COMPONENTE 2, INVESTIMENTO 1.3 – D.D. 1551.11-10-2022, code n. PE00000004, by the Sentinel North Program of Université Laval.

Extremophiles Collection of Institute of Biomolecular Chemistry National Research Council (CE-ICB CNR): microbial biodiversity preservation of extreme environments within the SUS-MIRRI.IT project

Vittoria, Maria¹; Romano, Ida¹; Finore, Ilaria¹; Leone, Luigi¹; Di Donato, Paola²; Nicolaus, Barbara¹; Poli, Annarita¹

¹ National Research Council of Italy (CNR), Institute of Biomolecular Chemistry (ICB), Pozzuoli (NA), Italy; ² Department of Science and Technology, University "Parthenope", Naples, Italy

Extremophiles, including thermophilic, halophilic, haloalkalophilic, psychrophilic, and polyextremophilic bacteria, are found in diverse habitats such as hydrothermal vents, salt lakes, polar regions, volcanic areas, and deserts [1]. Their adaptive strategies, involving various cellular metabolic mechanisms, led to the production of unique biomolecules with industrial applications in several biotechnology sectors [1,2]. Their ability to thrive under extreme conditions makes them valuable for astrobiology and assessing planetary habitability [2]. Extremophiles Research Group of Institute of Biomolecular Chemistry of the National Research Council (CNR) in Pozzuoli (Italy), through international collaborations and research projects, has sampled extreme niches, isolated and characterized microorganisms that are available to the scientific community and stakeholders as Extremophiles Collection (CE-ICB) (www.susmirri-catalog.di.unito.it/collections), included in the SUS-MIRRI project, which collects, characterizes, and distributes microbial strains for research and bio-industrial purposes preserving microbial biodiversity [3]. In fact, natural resource exploitation, climate change, and human activities are accelerating ecosystem destruction and microbial biodiversity loss, which means losing potential biological tools for global challenges. In addition, extremophilic biomass and its molecules (enzymes, exopolymers, lipids, etc.) can be hyper-produced by the bioprocess facility present at ICB-CNR. The availability of several sized bioreactors (www.icb.cnr.it/institute-facilities/bioprocessi) allowed the study of biotechnological potential of such biomolecules in terms of valorization of agro-industrial wastes, biodegradation, bioremediation, cultural heritage preservation, and agriculture [4,5,6]. Recently, *in silico* genome studies furnished an innovative tool for investigate the functional prediction and subsequently the potential biotechnological uses of Extremophiles [7].

References

- [1] Finore *et al.*, (2015) doi.org/10.1007/978-81-322-2595-9_13
- [2] Finore *et al.*, (2019) doi.org/10.3390/d11080128
- [3] Moretti *et al.*, (2024) doi.org/10.3390/su16093777
- [4] Finore *et al.*, (2016) doi: 10.1039/C5GC02774H
- [5] Romano *et al.*, (2020) doi.org/10.1016/j.ibiod.2020.104909
- [6] Finore *et al.*, (2024) doi.org/10.1016/j.ijbiomac.2024.131664
- [7] Yasar Yildiz *et al.*, (2022) doi: 10.3389/fmicb.2022.923038

Acknowledgment

This work was partially financed by PNRR European Commission – NextGenerationEU, Project “SUS-MIRRI.IT”, code n. IR0000005 and the MICS (Made in Italy – Circular and Sustainable) Extended Partnership MISSIONE 4 COMPONENTE 2, INVESTIMENTO 1.3 – D.D. 1551.11-10-2022, code n. PE00000004

Exopolysaccharides from psychrophilic microorganisms as biological device for plant promotion.

Pezzella, Giovanni¹; Poli, Annarita¹; Leone, Luigi¹; Santorufo, Lucia²; Caprio, Alfonso²; Vittoria, Maria¹; Romano, Ida¹; Finore, Ilaria¹

¹ National Research Council of Italy (CNR), Institute of Biomolecular Chemistry (ICB), Pozzuoli (NA), Italy; ² Department of Biology, University of Naples Federico II, Naples, Italy

Psychrophiles can grow and multiply at temperatures between 0°C and 20°C [1]. They are present in Alpine, Antarctic and Arctic environments, and have been isolated at high latitudes, in deep ocean waters, polar glaciers and snowfields. They are characterized by the presence of lipids in their cell membranes that are resistant to the rigidity caused by cold and often form antifreeze proteins to keep liquid their extracellular or intracellular space protecting the integrity of their cells even at temperatures below the freezing point of water. In addition, they secrete exopolysaccharides (EPS) that, surrounding the producer cells, represent a protective barrier against extreme low-temperature conditions. Some microbial EPS promote root colonization and retaining water form a hydrophilic biofilm. These polysaccharides form capsules or coatings around cellular surface and can aid in cell adhesion, environmental adaptation, uptake of nutrients and stress tolerance. Some psychrophiles that produce EPS are *Colwellia psychrerythraea*, *Pseudoalteromonas* sp., and *Olleya marilimosa*. An exopolysaccharide currently studied for plant growth promoting properties (PGP) determination is levan (Lev-2ASCA), produced by the psychrophilic bacterium *Pseudomonas* 2ASCA [2]. In addition, it has been proved that Lev-2ASCA efficiently chelates heavy metals, with interesting detoxifying effects on the soils in which it could be applied [3]. Therefore, experiments in pots with barley plant treated with Lev-2ASCA are ongoing, in frame of project PRIN 2022LPPFTY, -TREASURE. This strain, isolated from Extremophiles Research Group of Institute of Biomolecular Chemistry of the National Research Council (CNR) in Pozzuoli (Italy), is stored in Extremophiles Collection (CE-ICB) (www.susmirri-catalog.di.unito.it/collections) included in the SUS-MIRRI.IT project, which collect, characterize, and distribute microbial strains for research and bio-industrial purposes preserving microbial biodiversity.

References

- [1] Fendrihan, S., & Negoită, T. G. (2017). https://doi.org/10.1007/978-3-319-48327-6_7
- [2] Finore et al., (2020) <https://doi.org/10.3390/microorganisms8091282>
- [3] Finore et al., (2024) <https://doi.org/10.1016/j.ijbiomac.2024.131664>

Genome Prospecting and Wet-Lab Evidence of Plant Growth Promotion Traits in the Psychrophilic Arctic Bacterium *Pseudomonas* strain 2ASCA

Abbamondi, Gennaro Roberto¹; Papikyan, Razmik²; Leone, Luigi¹; Vittoria, Maria¹; Poli, Annarita¹; Finore, Ilaria¹

¹ CNR-ICB Institute of Biomolecular Chemistry, Italy; ² Institute of Botany After A. Takhtajyan of the National Academy of Sciences of the Republic of Armenia

The shift towards eco-sustainable techniques in agriculture is crucial to minimize the use of phytochemicals such as fertilizers and pesticides. In this context, innovative solutions like microbe-based biofertilizers are at the forefront. The PRIN project "TREASURE" explores *Pseudomonas* strain 2ASCA, an Arctic bacterium, for sustainable agriculture, focusing on plant growth promotion (PGP). *P. 2ASCA* is known for producing the polysaccharide levan, significant for its biotechnological applications [1]. The levan-based biofilm matrix could also be crucial for the successful inoculation of *P. 2ASCA* as a biofertilizer, enhancing its efficacy in promoting plant growth. *In silico* genomic analysis revealed genes (e.g., *glnA*, aconitate hydratase 2, catalase KatE) crucial for PGP, aiding nitrogen assimilation, energy production, and stress tolerance [2]. Additionally, genes for nutrient solubilization, siderophore biosynthesis, and ACC deaminase were identified, enhancing plant nutrient uptake and stress resilience. *Pseudomonas 2ASCA* also encodes enzymes for plastic degradation (e.g., LDPE, PHBV, PHA, PLA, and PCL), suggesting roles in agricultural sustainability and plastic pollution mitigation [3]. Experimental validation involved qualitative screening of direct and indirect PGP traits [4]. The bacterium displayed significant ACC deaminase activity, particularly at 15°C, with reduced activity at 5°C. IAA production was moderate at both temperatures, and organic acid production was slightly positive. Notably, *P. 2ASCA* was positive for siderophore production, indicating its biocontrol potential. Further studies are ongoing to experimentally investigate additional PGP traits and to deeply characterize the PGP features already identified, aiming to harness *P. 2ASCA*'s capabilities for biotechnological applications in sustainable agriculture. This strain belongs to Extremophiles Collection (CE-ICB) (www.susmirricatalog.di.unito.it/collections) included in the SUS-MIRRI.IT project.

References

- [1] Finore I *et al.*, (2024) International Journal of Biological Macromolecules, 268, 131664; doi: 10.1016/j.ijbiomac.2024.131664
- [2] Patz S *et al.*, (2021) bioRxiv, 2021.12.13.472471; doi: 10.1101/2021.12.13.472471
- [3] Gambarini V *et al.*, (2022) Database, baac008; doi: <https://doi.org/10.1093/database/baac008>
- [4]. Abbamondi GR *et al.*, (2016) Chem. Biol. Technol. Agric., 3:1; doi: 10.1186/s40538-015-0051-3

Acknowledgment

This work was partially financed by project PRIN 2022LPPFTY, -TREASURE, and by FOE-CNR FutuRaw Project.

Biosynthesis of Nanoparticles by Extremophilic Bacteria for Potential Biotechnological Application

Romano, Ida¹; Leone, Luigi¹; Poli, Annarita¹; Di Donato, Paola²; Finore, Ilaria¹

¹ ICB-CNR, Italy; ² University "Parthenope", Italy

Nanoscience and nanotechnology attract a great interest due to their potential impact on many areas such as energy, medicine, pharmaceutical industries, electronics, etc. This technology deals with small structures and small-sized materials of dimensions in the range of 1100 nm. Nanoparticles (NPs) show unique physical and biological properties, due to their high surface-to-volume ratio. Methods chemical and physical for the synthesis of nanoparticles, are considered unfavourable due to high capital cost, use of toxic reagents and the generation of hazardous wastes. Synthesis of nanoparticles by biological means offers cheap, non-toxic and eco-friendly alternative to the common physical and chemical methods. The use of biological samples like bacteria, plant extracts, fungi, polysaccharides and enzymes for the synthesis of NPs, provides various advantages like eco-friendliness and compatibility for biomedical and other pharmaceutical applications. In this frame, we used extremophilic bacteria to produce biogenic AgNPs and SeNPs. AgNPs were produced extracellularly by means of the thermophilic *Thermus thermophilus* strain SAMU; the haloalkaliphilic *Halomonas campaniensis* strain 5AG was used for the intracellular synthesis of SeNPs. The structural characterization was carried out by means of UV-visible spectroscopy, dynamic light scattering (DLS), transmission electron microscopy (TEM), Fourier transform infrared spectroscopy (FT-IR) and zeta potential. Both the Ag and SeNPs possessed a protein coating on their surface, they were organized in aggregates and showed antibacterial properties versus *Escherichia coli* DSM 648 and *Kokuria rhizophila* DSM 348. In addition, levan-type polysaccharide from psychrophilic *Pseudomonas* strain 2ASCA, develops particles in acidified water with heavy metal chelating capability revealing a strong affinity for Cr(III), never reported before for microbial levans. This behavior was studied using DLS and scanning electron microscopy (SEM).

Acknowledgment

This work was funded by CNR project FOE-2021 NutrAge—code DBA.AD005.225

This research was partially supported by the PNRR European Commission – NextGenerationEU, Project "Strengthening the MIRRI Italian Research Infrastructure for Sustainable Bioscience and Bioeconomy, SUS-MIRRI.IT", code n. IR0000005". This work was partially supported by PRIN 2022LPPFTY

The Haloarchaea Morphological Atlas discloses animal-like tissue structures emergence in single-celled archaea

Theopi Rados¹; Olivia Leland¹; Pedro Escudeiro²; John Mallon¹; Ido Caspy³; Elad Stolovicki⁴; Inés Patop¹; Sebastian Kadener¹; Vera Thiel⁵; Yoav Soen⁴; Tanmay Bharat³; Vikram Alva²; Alex Bisson¹

¹ Department of Biology, Brandeis University, Waltham, Massachusetts, USA; ² Department of Protein Evolution, Max Planck Institute for Biology Tübingen, Tübingen, Germany; ³ Structural Studies Division, MRC Laboratory of Molecular Biology, Cambridge, United Kingdom; ⁴ Department of Biomolecular Sciences, Weizmann Institute of Science, Israel; ⁵ Department of Microorganisms, Leibniz Institute DSMZ – German Collection of Microorganisms and Cell Cultures GmbH, Germany

With our large collection of >100 haloarchaea, the Leibniz Institute DSMZ joins a comprehensive evolutionary developmental biology project with international researchers studying the biophysical and spatio-temporal control of unusual cell shape transitions across salt-loving archaea using state-of-the-art microscopy from single-cell and single-molecule tracking, microfabrication, and genetics, building an unprecedented “Haloarchaea Morphological Atlas”. It is hypothesized that together with reading out their chemical environment, archaeal cells evolved to sense physical cues to build different cell shapes and mediate social behavior within the same and across different species, which is studied as part of the project. To explore this idea, the mechanisms of mechanosensing in Archaea, the closest single-celled prokaryote (cells without a nucleus) to animals, are investigated. Surprisingly, archaeal cells react to confinement and compression with the development of a tissue-like state, with cells physically connected, resembling the structure of an epithelial monolayer. It was shown that this multicellular development depends not on material stiffness but on axial compression. In the first hours after being compressed, the cells grow larger without cell division but with accelerated DNA and protein synthesis. After reaching a precise cell volume, cell division is once again reinstated, but now, instead of separating, cells remain attached by cell-cell junctions. In the ongoing project, it will be further addressed which specific factors respond to mechanical compression and which factors contribute to developing and maintaining the multicellular structures. This novel and unexpected discovery introduces a new way of understanding and studying tissues in a simpler and straightforward model organism.

Session 5 – Microbiomes preservation and exploitation

Wednesday 18th September, 3:00pm

Chairs: Ferrara Massimo (Italy), Aznar Rosa (Spain).

Microbiomes for the Industry

Porcar, Manuel¹

¹ *DARWIN BIOPROSPECTING EXCELLENCE, Valencia, Spain*

Our planet is home of a large biodiversity of microorganisms, which do not live alone but with a dynamic interaction with their tiny neighbours. Bioprospecting, that is, the search, characterisation and selection of microbial taxa, can lead to the use of microbiomes and simplified consortia for the food industry. In particular, simplified consortia can be very useful for improved sourdough baking; for yielding novel fermented dairy products with organoleptic and/or functional new properties; and can also be the preferred strategy for brewing alcoholic and non alcoholic beverages that combine industrialisation suitability and organoleptic value. Bioprospecting microbiomes can also lead to develop non food, environmental biosolutions, such as fostering biogas production through bioaugmentation and digestate transplantation; or plastic biodegradation and sulphate reduction in polluted wastewater. Besides culturomics, microbiomes are also a source of genes and enzymes with potential in biomedicine and bioremediation. Culture-independent, genomic mining is thus also a valuable tool to develop microbial solutions with biotechnological potential. Finally, bioprospecting lies on the interphase between industrial applications and social benefits, and the implications of such interaction in the frame of the Nagoya protocol will be discussed.

Exploring soil microbiome preservation strategies: culturable fraction, metabolic profiling and metagenomics

Sbarra, Federico¹; Visca, Andrea¹; Garello, Marco²; Sevi, Filippo¹; Tabacchioni, Silvia¹; Di Gregorio, Luciana¹; Costanzo, Manuela¹; Colantoni, Eleonora¹; Aloï, Francesco²; Varese, Cristina³; Spadaro, Davide²; Bevivino, Annamaria¹

¹ AgriFood Sustainability, Quality and Safety Laboratory. ENEA, Casaccia, Roma, Italy; ² Dept. Agricultural, Forest and Food Sciences (DISAFA), University of Torino; ³ Department of Life Sciences and System Biology (DBIOS), University of Torino

Holistic approaches are required to better understand the distribution of microbial diversity and the functional profile of communities in soil microbiomes. In the present work we investigated how different sample preservation techniques can affect the soil rhizosphere microbial communities by analysing the culturable fraction of microorganisms, the microbiota and the metabolic fingerprint. Three different storage conditions were considered: refrigeration at 4°C, freezing at -80°C, and lyophilization. Six rhizosphere samples from strawberry, grapevine and kiwifruit plants grown in two field soil types were analysed. Microbial community structures and growth strategies were investigated by means of total microbial count, enumeration of copiotrophs and oligotrophs bacteria, r-k strategists and eco-physiological (EP) index. A total of 300 bacterial strains and 315 fungi were isolated for further investigation of plant growth promoting traits. Functional-level metabolic profiling of microbial communities based on Biolog EcoPlates highlighted widespread carbon sources utilisation in samples with high population diversity, confirmed by metagenomic analysis, such as kiwifruit and strawberry, when compared to poor species abundance soils, as grapevine. Linking the distribution of microbial diversity and ecosystem functioning is essential to understand community responses to changing environment, especially in forced conditions such as samples preservation. For that reason, the same analysis will be performed on samples at 6 and 12 months and results will help better understand structural shifts during microbiome conservation and which functional features in the community can be affected during long term storage. In parallel, within isolates showing PGP traits, microbial consortia set-up will be carried out to exploit natural inhabitants of soil communities (NatComs) as biofertilizer to enhance crop productivity.

Acknowledgements

The authors thank Luca Cocolin for his valuable support. This activity has received funding from the SUS-MIRRI.IT Project "Strengthening the MIRRI Italian Research Infrastructure for Sustainable Bioscience and Bioeconomy", Project code IR0000005, CUP D13C22001390001.

Microbiome Biobanking: The missing link

Kostic, Tanja¹; CONSORTIUM MICROBE²

¹ AIT Austrian Institute of Technology GmbH; ² MICROBE project

Culture collections have a history of supporting microbiological research, primarily through the accession, preservation and supply of axenically cultured microorganisms. However, in nature, microorganisms do not exist on their own; they interact with millions of other microorganisms in complex systems called microbiomes [1]. Microbiomes provide various crucial ecosystem services and are thus essential to the well-being of plants, animals and the environment. With developments in technology, microbiome research is changing the way culture collections and biobanks need to support their user communities. Key requirements for microbiome biobanking have been proposed [2, 3], including the development of standards, and emphasising the need to deposit material and supply cultures, samples and associated data for future research. Additionally, this will provide a mechanism to protect intellectual property, and help researchers adhere to legislative and regulatory requirements, including the Nagoya Protocol of the CBD. Integral to the above is the need to preserve environmental samples and their microbiota. MICROBE project (www.microbeproject.eu) aims to deliver innovative validated technological approaches that will enable:

- maximal preservation of taxonomic and functional biodiversity in selected microbiome samples;
- optimal collection and preservation of microbiome samples for defined subsequent analyses;
- targeted isolation of microbiome members from different domains and assembly of synthetic consortia that retain (and even optimize) the functional diversity of original microbiomes. Furthermore, MICROBE will provide a comprehensive operational blueprint for the establishment of microbiome biobanking infrastructure, including technological requirements, methodological workflows, data pipelines, standards, legal and ethical guidelines, training plans and business opportunities. First insights from the project will be provided.

References

- [1] Berg, et al. (2020) Microbiome definition re-visited: old concepts and new challenges. *Microbiome* 8. <https://doi.org/10.1186/s40168-020-00875-0>
- [2] Ryan, et al. (2020a) Development of Microbiome Biobanks – Challenges and Opportunities. *Trends in Microbiology*. <https://doi.org/10.1016/j.tim.2020.06.009>
- [3] Ryan, et al. (2020b) Towards a unified data infrastructure to support European and global microbiome research: a call to action. *Environmental Microbiology*. <https://doi.org/10.1111/14622920.15323>

Acknowledgements

MICROBE project is funded by the European Union (Grant agreement ID: 101094353)

Cryopreservation and recovery of a complex hypersaline microbial mat community

Grego, Michele¹; Bolhuis, Henk²

¹ *Sorbonne University, Roscoff Culture Collection (RCC), Station Biologique de Roscoff, Pl.Georges Teissier, Roscoff;* ² *NIOZ*

Cryopreservation of microorganisms is an essential tool in industrial and food applications where conservation of microbial activity and critical beneficial traits need to be guaranteed to provide a consistent product or production process. This often refers to simple, single species or low diversity assemblages in liquid cultures that can easily be revived and regrown to perform the desired process. Cryopreservation is also of essence for scientific experimentation where many environmental samples are taken in remote sampling sites and at high costs. Biobanking, or the long term preservation and potential revival of complex, structured samples come with an additional challenge related to maintaining the structure upon revival. Here we look at cryopreserving and reviving a complex photosynthesis driven microbial mat from a hypersaline ecosystem. Amplicon sequencing of the 16S and 18S ribosomal RNA gene was used to determine the community composition of bacteria and eukaryotes respectively. The tests included the use of different cryopreservative agents and different times of cryopreservation at 150 °C. Upon revival, the cryopreservatives cannot be separated from the preserved samples without disturbing the community structure, while carryover of these compounds may influence reconstitution of the communities. Indeed, although both glycerol and Me₂SO are good cryopreservatives of microbial assemblages, carryover of these compounds had a profound negative effect on the reestablishment of a functional microbial mat. Best cryopreservation and reconstitution results were obtained in the absence of a cryopreservative agent or when methanol was used.

Direct Injection Mass Spectrometry for the Real-Time Volatilomics in Food System Microbiomes: the Potential of Providing Temporal Dimension in Multi-Omics Studies

Corvino, Antonia¹; Capozzi, Vittorio²; Khomenko, Iuliia¹; Biasioli, Franco¹

¹ *Research and Innovation Centre, Fondazione Edmund Mach, San Michele all'Adige (TN), Italy;* ² *National Research Council of Italy Institute of Sciences of Food Production (ISPA), Foggia, Italy*

Fermentation exemplifies sustainable innovation in food systems by utilizing microbial diversity to enhance product quality and safety with low environmental impact. Diverse fermentation processes also serve as model ecosystems for microbiology studies and bio-tool design. Understanding and optimizing fermentation through 'omics' technologies, especially metabolomics, is crucial. Metabolomics offers insights into the metabolic state of microorganisms and the dynamics of metabolic pathways. Microbial Volatile Organic Compounds (mVOCs) are key metabolites that influence sensory perception and consumer acceptance. Profiling VOCs during fermentation provides valuable data on microbial ecosystems, identifying markers to monitor fermentation dynamics and optimize product yield, quality, and safety. Proton Transfer Reaction-Time-of-Flight Mass Spectrometer (PTR-ToF-MS), a Direct-Injection Mass Spectrometric (DIMS) technology, enables non-destructive, real-time analysis with high automation and sensitivity. PTR-ToF-MS effectively evaluates microbial volatilome in various agro-industrial fermentation processes. We present case studies demonstrating PTR-ToF-MS's potential for volatilomics in food microbiology and microbiome research, emphasizing the significance of temporal dimension metabolomics. This approach is particularly relevant for integrating the temporal perspective in multi-omics studies of dynamic systems like food-related microbiomes.

Acknowledgements

We developed this overview in the framework of the projects i) "Interconnected Nord-Est Innovation Ecosystem (iNEST) funded by the European Union Next-Generation EU (PIANO NAZIONALE DI RIPRESA E RESILIENZA (PNRR) – MISSIONE 4 COMPONENTE 2, INVESTIMENTO 1.5 D.D. 1058 23/06/2022, ECS00000043) and ii) "ON Foods-Research and innovation network on food and nutrition Sustainability, Safety and Security– Working ON Foods" funded by NextGeneration EU [PNRR], in the framework of the Component 2 Investment 1.3 Award Number: Project code PE00000003, Vittorio Capozzi is also supported by NUTRAGE CNR project FOE2021 DBA.AD005.225.

Possible “microbial bioremediators” isolated from wasted lands

De Santis, Alessandro¹; Corbo, Maria Rosaria¹; Bevilacqua, Antonio¹; Daniela, Campaniello¹; Sinigaglia, Milena¹

¹ University of Foggia, Foggia, Italy

Human population is increasing every year and consequently the need for goods and services, and the associated waste and refuse. The current practices used for waste disposal usually cause considerable damages to environment, reducing the disponible land for agriculture and raw food production; furthermore, the constant use of pesticides, phytochemicals and antibiotics to improve productivity is negatively impacting the consumers' health. This research aims at proposing solutions which can improve the current situation of soil health with direct consequences on human health by using selected microorganisms able to reduce harmful substances.

As a first part of this project, microbiological parameters were analysed starting from ten different lands wasted by anthropic activity. The available results are as follows (average values): *Pseudomonas* spp. (9.40×10^6 CFU/g), actinobacteria (4.09×10^6 CFU/g), clostridia (1.35×10^2 CFU/g), and aerobic bacteria (3.53×10^6 CFU/g). Pseudomonadaceae were chosen and isolated because of their involvement, as demonstrated in other studies [1,2], in bioremediation activity against heavy metals and organic pollutants.

From a chemical perspective, the health profile of soils was studied, with a focus on heavy metals, for every soil sample and compared to the microbiological parameters to search clusters and correlations among the factors useful to deepen the genomic research.

Then, all the isolated strains were tested for phenotypic and technological parameters to assess the main characteristics with a focus on the resistance to heavy metals.

The information acquired in the first phase will be the background to assess the reducing capacity against heavy metals, the metagenomic research on soil samples and the 16S RNA analysis on selected strains.

References

- [1] Vélez JMB, Martínez JG, Ospina JT, Agudelo SO. Bioremediation potential of *Pseudomonas* genus isolates from residual water, capable of tolerating lead through mechanisms of exopolysaccharide production and biosorption. *Biotechnol Rep (Amst)*. 2021 Oct 29; 32:e00685. doi:10.1016/j.btre.2021.e00685.
- [2] Fanghui Hu, Panlin Wang, Yunhan Li, Jiahuan Ling, Yongqiang Ruan, Jiaojiao Yu, Lihui Zhang, Bioremediation of environmental organic pollutants by *Pseudomonas aeruginosa*: Mechanisms, methods and challenges, *Environmental Research*, Volume 239, Part 1, 2023, 117211, <https://doi.org/10.1016/j.envres.2023.117211>

Food microbiomes and packaging: past, present and future

Gökşen, Gülден¹; Cunedioğlu, Hülya²; Fragasso, Mariagiovanna²; de Chiara, Maria Lucia³; Capozzi, Vittorio³

¹ Department of Food Technology, Vocational School of Technical Sciences at Mersin Tarsus Organized Industrial Zone, Tarsus University, Turkey; ² Department of Agricultural Sciences, Food, Natural Resources and Engineering, University of Foggia, Foggia, Italy; ³ Institute of Sciences of Food Production of National Research Council (ISPA-CNR), Foggia, Italy

Microbiomes are resources of growing interest for promoting sustainable innovation in food systems. Applications in the field of packaging represent an emerging sector that is still little explored. Here, we propose a review of the different types of packaging that include microorganisms and which are used/proposed for the food industry, giving particular emphasis to edible packaging containing lactic acid bacteria and packaging solutions that enhance microbial biocontrol activities. In light of the solutions that use single resources, a discussion is proposed on the potential use of microbiomes, carrying out a comparative analysis of the strengths and weaknesses. This is an extremely dynamic sector with broad prospects, crucial for the development of sustainable solutions for the food industry.

Acknowledgements

We are exploring the potential of this driver of innovation in the framework of the ongoing projects i) INTACTBioPack INTelligent, ACTIVE MicroBIOMebased, biodegradable PACKaging for Mediterranean food – founded by the PRIMA Section 2 2023 multi-topic and ii) the NextGeneration EU [PNRR], in the framework of the Mission 4 Component 2 Investment 1.3-Award Number: Project code PE00000003, Project title: “ON Foods-Research and innovation network on food and nutrition Sustainability, Safety and Security–Working ON Foods”.

Marine microbial community sample cryopreservation: finding the best preservation strategy

Bachy, Charles¹; Grego, Michele²; Romac, Sarah¹; Gachenot, Martin²; Hervéou, Katell²; Gourvil, Priscillia¹; Joannic, Julie²; Probert, Ian²

¹ Centre National de la Recherche Scientifique (CNRS); ² Sorbonne University

Marine microbiomes comprise communities of diverse microorganisms (bacteria, archaea, protists and microalgae) and their 'theatre of activity' (i.e. metabolites, signalling molecules and surrounding environmental conditions). These microorganisms play an essential role for the oceans and our planet by participating in the balance of natural cycles (nutrients, oxygen and carbon dioxide). In order to facilitate the science required to make major advances in microbiome research, methods and technologies are needed to capture and ensure stable long-term maintenance. Since 2010, the Roscoff Culture Collection (RCC) has cryopreserved more than 3000 mono-specific phototrophic marine microbial strains, including different lineages of microalgae and cyanobacteria. The RCC is now developing methods to cryopreserve complex marine microbiomes with the aim of capturing and culturing the widest possible range of microbial diversity. We investigated the effects of different cryoprotectants on the viability of coastal marine microbiomes using various approaches to assess post-cryopreservation viability. Once samples were thawed, we tracked changes in the microbial community using RNA metabarcoding, measured growth rates using cytometry, and used a variety of traditional methods to culture bacterial and algal strains. Together, the results will be used to identify the best strategy for conserving marine microbial community samples through cryopreservation.

Fermenting metacommunities of food interest: an invaluable biodiversity asset to be maintained overtime in its wholeness and functionality Micro4ever

Ferrara, Massimo¹; Latronico, Rosanna²; Cozzi, Giuseppe¹; Verrone, Laura¹; Zotta, Teresa³; Filannino, Pasquale²

¹ *Institute of Sciences of Food Production, National Research Council (ISPA-CNR), Bari, Italy;* ² *Department of Soil, Plant and Food Science, University of Bari Aldo Moro, Bari, Italy;* ³ *Department of Agricultural, Forestry, Food and Environmental Sciences, University of Basilicata, Potenza, Italy*

Micro4ever aims to investigate the effect of cryopreservation (CrP) and freeze-drying (FD) on the functionalities of whole microbial consortia (MC) associated with different fermented foods (dairy and bakery products) after their long-term storage and revitalization. Specifically, we will use the MC of milk starters (lattoinnesto; LI), whey starters (sieroinnesto; SI) and sourdoughs (SD) as models of dynamic MC characterized by a high level of biodiversity. The main objectives of Micro4ever are: (i) to preserve intact MC with retained viability and functionality for future meta-omics studies, cultivation, and application; (ii) to preserve and exploit in situ the invaluable biotechnological potential behind the poorly cultivable microbial players of MC; (iii) to optimize the conditions of sampling and long-term storage of MC, providing the maximum protection to different kinds of cells and cellular components within the MC; (iv) to validate the post-preservation viability and functionality of preserved MC; (v) to propose new models governing the assembly, organization, functionality and stability of MC of food interest. While the advantages of fully exploiting the whole functional potential of MC are clear, the accomplishment of that intent is fraught with conceptual and technical challenges. Micro4ever will establish and optimize the cultivation and preservation methods that protect the full phenome of MC and ensure genotypic stability of the MC, which is essential for the preservation of MC biodiversity. In particular, Micro4ever will implement a multi-omics workflow, through which we will first decipher the taxon composition and the functional redundancy of MC from LI, SI, and SD at metagenomics, metatranscriptomics, meta-phenomics and metabolomics levels. Then, we will investigate in depth the effect of short- and long-term storage, and will validate the effectiveness of CrP and FD protocols and the functionality of preserved MC also in food matrices.

Acknowledgements

Project funded by the European Union Next Generation EU; Project code P2022RJYCN, CUP H53D23010720001, Project title "Fermenting metacommunities of food interest: an invaluable biodiversity asset to be maintained overtime in its wholeness and functionality Micro4ever".

Biochemical and biophysical characterization of two recombinant enzymes obtained from termite gut metagenome

Bruno Baron, Camila¹ ; Mon, Maria Laura¹; Marrero Díaz de Villegas, Rubén¹; Cattaneo, Andrea²; Di Donato, Paola²; Poli, Annarita²; Soria, Marcelo A.³; Rojo, Maria Cecilia³; Combina, Mariana³; Finore, Ilaria²; Talia, Paola Monica⁴

¹ Instituto de Agrobiotecnología y Biología Molecular (IABIMO), INTA/CONICET Buenos Aires, Argentina; ² Institute of Biomolecular Chemistry (ICB), Consiglio Nazionale delle Ricerche (CNR), Pozzuoli, Italy; ³ Estación Experimental Agropecuaria Pergamino, Instituto Nacional de Tecnología Agropecuaria (INTA), Pergamino, Buenos Aires, Argentina; ⁴ Estación Experimental Agropecuaria Mendoza, Instituto Nacional de Tecnología Agropecuaria (INTA), Mendoza, Argentina

In this study, we characterized two novel glycoside hydrolase family 10 (GH10) enzymes (Xyl10C and Xyl10E) [1] identified in the termite gut microbiome. Both enzymes showed activity on beechwood xylan, whereas Xyl10E also showed activity on barley -glucan. Purified Xyl10C showed optimum activity in the pH range 7.0-8.0 and at a temperature of 50-60°C, while Xyl10E was active at a wider pH range (5.0-10.0), and at 50°C. The residual activities of Xyl10C and Xyl10E after 8 hours of incubation at 40°C were 85% and 70%, respectively, showing thermal stability. The enzymatic activity of Xyl10C increased in the presence of 5 M NaCl (115%) and was only inhibited in the presence of 0.5% SDS and reduced with β -mercaptoethanol, whereas both the xylanase and glucanase activities of Xyl10E were inhibited only in the presence of MnSO₄, NaCl and SDS. In addition, both enzymes were able to hydrolyze pretreated Sorghum bicolor bagasse (SBB), releasing xylose and xylobiose as the main products [2]. Xylose and xylobiose production suggest their potential use as prebiotics and as other valuable industrial applications.

References

- [1] Ben Guerrero EB (2020) doi:10. 1007/ s0025302010831-5;
- [2] Finore et al., (2016) doi:10.1039/c5gc02774h

Acknowledgements

This study was supported by grants from (INTA) (PI 085, 089, 122 and 159), the Agencia Nacional de Promoción Científica y Tecnológica (ANPCyT) Proyectos de Investigación Científica y Tecnológica (PICT) 2018-#4149, 2019-#3156, 2020-#3570, Fundación Williams #014 and CONICET Project # PIP-2021-2561. The authors thank the MICS (Made in Italy—Circular and Sustainable) Extended Partnership from the European Union Next-Generation EU (PIANO NAZIONALE DI RIPRESA E RESILIENZA (PNRR) —MISSIONE 4 COMPONENTE 2, INVESTIMENTO 1.3—D.D. 1551.11-10-2022, PE00000004).

Exploring Microbiomes in PFAS Contaminated Environments: Preservation for Bioremediation Strategies.

Nawaz, Asma¹; Andreolli, Marco¹; Tontini, Beatrice¹; Lampis, Silvia¹

¹ *Department of Biotechnology & VUCC-DBT, University of Verona, Verona, Italy*

Per- and poly-fluoroalkyl substances (PFASs) are a group of xenobiotics widely used for their water and grease-resistant properties [1] that have raised significant environmental and health concerns due to their persistence and potential toxicity [2]. Their bioaccumulation led to biomagnification up the food chain, exerting their toxic effects, and increasing the likelihood of severe health outcomes over time [3]. Bioremediation may be a promising strategy to mitigate PFAS effects in contaminated environments. Understanding microbiomes in PFAS-polluted sites is crucial due to their potential for bio-transforming or bio-accumulating PFAS. Thus, our focus is to study microbiomes in PFAS-polluted environmental matrices and their potential for PFAS bioremediation. This aim is pursued in conjunction with the SUS-MIRRI.IT project regarding the study of microbiome in different matrices. The experimental approach involves preserving environmental samples to study how microbial communities can be efficiently stored preserving their bioremediation potential. To achieve our objectives, we analyzed two different matrices: soil and sediment. Soil samples (microbiomes) were preserved at -80°C (for 12 months) in glycerol (30%) as a cryoprotective agent to investigate their vitality through total microbial counts and Ecoplates assays. Additionally, we employed microbial barcoding to identify microbial diversity and the bioremediation potential of these microbiomes will be evaluated through PFAS content determination. We decided to perform these analyses over a twelve-month time span; T₀ (pre-storage), T₁ (3 months post-storage), T₂ (9 months post-storage), and T₃ (12 months post-storage). Preliminary results (T₀ and T₁) show successful preservation. However, confirmation awaits experiments at T₂ and T₃. Further analysis at T₂ and T₃ will be performed to determine the long-term effects of preservation of samples on the microbiome and their potential application to bio-transform PFAS.

References

- [1] Giesy, J. P., & Kannan, K. (2002). Perfluorochemical surfactants in the environment. *Environmental Science & Technology*, 36(7), 146A-152A.
- [2] Sunderland, E. M., Hu, X. C., Dassuncao, C., Tokranov, A. K., Wagner, C. C., & Allen, J. G. (2022). A review of the pathways of human exposure to poly and perfluoroalkyl substances (PFAS) and the health effects associated with exposure. *Environmental Science & Technology*, 56(2), 846-855.
- [3] Cheng, H., Lv, C., Li, J., Wu, D., Zhan, X., Song, Y., & Jin, H. (2022). Bioaccumulation and biomagnification of emerging poly-and perfluoroalkyl substances in marine organisms. *Science of The Total Environment*, 851, 158117.

UMCC's investigative approach for the preservation and exploitation of fermented beverage microbiomes

Arena, Mattia Pia¹; Signorello, Lara¹; Catellani, Lisa¹; Gullo, Maria¹

¹ *Department of Life Sciences, University of Modena and Reggio Emilia, Reggio Emilia, Italy*

The role of microbial culture collections is fundamental for the preservation, study and utilization of microbial resources. Although the conservation and maintenance of pure microbial strains is a well-established procedure within the management system of microbial collections, the preservation of microbiome, defined as the entire microbial community that occupies a defined habitat and their theater of metabolic activities [1], still needs to be consolidated as a routine practice. However, acquiring knowledge on microbiome conservation and a better understanding of the interactions among microbial communities and food substrates, is essential for their biotechnological valorization.

Unimore Microbial Culture Collection (UMCC) (<https://umcc.bioaware.com/>), located at the University of Modena and Reggio Emilia, cryopreserves over 3000 microbial strains isolated from different food substrates. Recently, UMCC began an in-depth study regarding microbiome preservation from fermented beverages. A total of ten microbiome samples from wine vinegar, pomegranate vinegar, apple cider vinegar and Kombucha tea were appropriately sampled, analyzed through both culture-dependent and molecular approaches, and preserved by ultrafreezing and freeze-drying. A comparative evaluation of microbiomes before and after storage is currently underway with the aim of evaluating the effectiveness of long-term storage methods and, then, the possible industrial exploitation of microbiomes.

References

[1] Berg, G., et al., Microbiome definition re-visited: old concepts and new challenges. *Microbiome*, 2020. 8(1): p. 103.

Acknowledgements

UMCC is supported by the European Commission – NextGenerationEU, Project SUS-MIRRI.IT "Strengthening the MIRRI Italian Research Infrastructure for Sustainable Bioscience and Bioeconomy", code n. IRO000005 and by and by the European Union-NextGenerationEU Grant, CN_00000033, Project "National Biodiversity Future Center-NBFC", CUP E93C22001090001.

A cryopreservation protocol of microbial consortia from Apulian table olives

Ferrara, Massimo¹; Gialluisi, Katia¹; Verrone, Laura¹; De Vero, Luciana¹; Petruzzino, Giuseppe¹; De Simone, Nicola²; Perrone, Giancarlo¹; Moretti, Antonio¹; Capozzi, Vittorio²

¹ *Institute of Sciences of Food Production, National Research Council (ISPA-CNR), Bari, Italy;* ² *Institute of Sciences of Food Production of National Research Council (ISPA-CNR), Foggia, Italy*

The field of microbiome research has evolved rapidly over the past few decades and has become atopic of great scientific, industrial, and public interest. Exploiting microbiomes as drivers of innovation in food systems and promoting reproducibility in basic, applied and industrial science shed new light on the preservation of microbial communities associated with food fermentation/specific matrices. Therefore, the challenges of optimally preserving microbiome samples are huge, with difficulties ranging from conserving microbial consortia to preserving microbiome integrity/functionality. Fermented table olives represent the most representative and diffused plant-based fermented products in Mediterranean countries, with relevant productions in Italy, in general, and in the Apulian region, in particular. Within the SUS-MIRRI.IT project, a cryopreservation protocol has been developed to preserve the microbiota from typical Apulian table olives cv Leccino, by using glycerol or DMSO as cryoprotectants, and a storage temperature of -135 °C. The microbial consortium was studied before and after mid-term storage using a culture-dependent approach, RNA-based metabarcoding analysis, and metabolic profiling evaluation. Results showed that after six months of cryopreservation, the viability of the microbial consortium slightly decreased regardless of the cryoprotectant used, and no significant changes in the metabolic profile were observed. Also, the metabarcoding analysis showed no significant differences in relative abundances after storing period. Results confirmed the effectiveness of the developed cryopreservation protocol, proper preservation of the microbial consortium, and its functionality during a mid-term storage period. They also pointed out preliminary insights on the possibility of exploiting them in fermenting processes after cryopreservation.

Acknowledgements

Project SUS-MIRRI.IT “Strengthening the MIRRI Italian Research Infrastructure for Sustainable Bioscience and Bioeconomy” Area ESFRI “Health and Food”, granted by the European Commission – NextGenerationEU Code N° IR0000005

Synthetic microbial communities as a tool to understand microbial interactions and their role in bacterial cellulose production

Lasagni, Federico¹; Giavalisco, Marilisa²; Arena, Mattia Pia¹; Zotta, Teresa²; Gullo, Maria¹

¹ *Unimore Microbial Culture Collection Laboratory, Department of Life Sciences, University of Modena and Reggio Emilia, Reggio Emilia, Italy;* ² *Department of Agricultural, Forestry, Food and Environmental Sciences, University of Basilicata, Potenza, Italy*

Genetic and physiological diversities among microorganisms result in the emergence of complex microbial communities, where interactions among cells can influence each other's behaviours and phenotypes. Microbial communities play a crucial role in the functionality of several ecosystems, such as bio-geochemical cycles, agri-food productions, and human health, boasting an enormous metabolic potential. The construction of synthetic microbial communities allows the generation of targeted systems with reduced complexity, capable of achieving multiple tasks in a more efficient way compared to single strains. Synthetic microbial communities may be exploited as models to understand microbial dynamics and regulatory pathways, as well as tools for several biotechnological applications in chemical, pharmaceutical, and agri-food fields. Within the PRIN 2022 Project "BioCellulose production from a Synthetic Microbial Community: sustainable process for food and healthy applications SynBioCell" (code: 20228Z34PF, CUP: E53D23010680006), a novel synthetic microbial community is being designed and assembled. Following a chemometric approach, different species belonging to acetic acid bacteria, lactic acid bacteria and yeasts were tested for their suitability to be combined in a synthetic microbial community, aimed at enhancing bacterial cellulose synthesis using various defined media and agri-food substrates. Each synthetic microbial community member will cooperate for the development of a functional ecosystem, where interactions and metabolite sharing are believed to promote cell growth and bacterial cellulose yield. The development of a targeted synthetic microbial community may help facing the increasing demand of bacterial cellulose and other bioproducts through a sustainable and efficient approach.

Acknowledgements

Part of this work was granted by the European Commission NextGenerationEU – within the projects PRIN 2022 – SynBioCell (code 20228Z34PF) and SUS-MIRRI.IT (code IR0000005).

Back-slopping of dried, freeze-dried and frozen Tempeh: the effect of long-term preservation techniques on the microbiome and volatilome

Reale, Anna¹; Di Renzo, Tiziana¹; Boscaino, Floriana¹; Nazzaro, Stefania¹; Marena, Pasquale¹; Ferrocino, Ilario²

¹ *Institute of Food Sciences, National Research Council (CNR-ISA), Avellino, Italy;* ² *DISAFA, University of Turin, Italy*

Tempeh is a traditional Indonesian fermented food, commonly made from soya, but often also from peas, beans, chickpeas and other legumes. It is mainly produced in Asia, particularly in Indonesia, where it is a staple food, but its production is also growing in India, Japan, United States and Europe. Tempeh is an ideal meat substitute, high in protein, and a nutritious food that may confer health benefits due to its high bioavailability of nutrients and phytochemicals, showing improved action on oxidative stress, glycaemic control and blood lipid levels [1].

Tempeh production involves several steps, including soaking, dehulling, cooking and fermentation with the starter mould *Rhizopus* spp. However, the nature of the tempeh production processes creates a unique microbiome derived not only from the microbial inoculum, but also from the raw materials and environment. Over the past decade, metagenomic studies of the microbial community during tempeh production highlighted a complex microbial consortium consisting not only of *Rhizopus* spp. (used as inoculum) but also of other microorganisms belonging to the phyla Proteobacteria, Firmicutes, and Bacteroidetes [2].

Preservation of the tempeh microbiome is essential to maintain its functional and nutritional properties. This can be achieved through proper fermentation processes, storage, and handling techniques to minimize contamination and degradation of the microbial community [3].

Our aim was to evaluate the effect of different long-term preservation methods (drying, freeze-drying and freezing) on the tempeh microbiome and the possibility of reusing the stored product as a starter for a subsequent back-slopping fermentation, a method in which a portion of the fermented product is added back to the new batch to continue fermentation. Tempeh stability over time was evaluated for microbiome and volatilome (HS-SPME-GC-MS) aspects.

References

- [1] Rizzo G. (2024) Soy-Based Tempeh as a Functional Food: Evidence for Human Health and Future Perspective. *Front. Biosci. (Elite Ed)* 16, 3.
- [2] Yulandi A. et al. (2020) Shotgun metagenomic analysis reveals new insights into bacterial community profiles in tempeh. *BMC Res Notes* 13, 562.
- [3] Subali D. et al. (2023) Soy-Based Tempeh Rich in Paraprobiotics Properties as Functional Sports Food: More Than a Protein Source. *Nutrients* 15, 2599.

Acknowledgements

This research was funded by the project SUS-MIRRI.IT "Strengthening the MIRRI Italian Research Infrastructure for Sustainable Bioscience and Bioeconomy", code n. IR0000005.

New frontiers for *Saccharomyces cerevisiae* flor strains

Sanna, Anna Maria Laura¹; Zara, Giacomo¹; Bianco, Angela¹; Coronas, Roberta¹; Budroni, Marilena¹

¹ Department of Agricultural Sciences of University of Sassari, Italy

The process of biological ageing by *Saccharomyces cerevisiae* flor strains characterizes wines of considerable value, such as Spanish Sherry wines and some traditional wines of Sardinia such as Malvasia di Bosa, Vernaccia di Oristano, and Arvisionadu del Goceano. This process is dependent on FLO11 gene expression and involves the transition from fermentative to oxidative metabolism, with the formation of the typical yeast biofilm (flor) on the wine surface at the end of fermentation. In recent years, the evolution and application of modern winemaking techniques, which are increasingly effective in avoiding the presence of even minimal oxygen concentrations during wine ageing, makes it more difficult to isolate flor yeasts from the Sardinian wines. In this context, the role of culture collections for the supply and valorization of microbial resources previously isolated and characterized, appears crucial. In Sardinia, the MBDS-UNISSCC microbial collection (certified ISO 9001:2015) belonging to the MIRRI-IT network of collections and equipped with a federated database "www.mbds.it", preserves microbial resources isolated from food and environmental matrices. Yeast strains, including flor strains, of oenological interest have been isolated from Sardinian musts and wines, characterized, and stored since 1960. Currently, these resources are cryopreserved at -80 °C and have shown a satisfactory response during viability testing. Recently, the flor formation ability of 56 strains ascribed to the genus *Saccharomyces*, isolated, characterized and stored between 1965 and 1985, has been confirmed. MBDS-UNISSCC aims to preserve this microbial biodiversity by characterizing all the stored flor strains at the molecular level and developing preservation methods of the whole biofilm.

References

- Legras, J. L., Moreno-Garcia, J., Zara, S., Zara, G., Garcia-Martinez, T., Mauricio, J. C., & Budroni, M. (2016). Flor yeast: new perspectives beyond wine aging. *Frontiers in microbiology*, 7, 503.
- Zara, G., Budroni, M., Mannazzu, I., Fancello, F., & Zara, S. (2020). Yeast biofilm in food realms: occurrence and control. *World Journal of Microbiology and Biotechnology*, 36, 1-10

Microbiome characterisation of sourdough from the MBDS-UNISSCC microbial collection

Coronas, Roberta²; Zara, Giacomo¹; Bianco, Angela¹; Sanna, Anna Maria Laura²; Budroni, Marilena¹

¹ Department of Agricultural Sciences of University of Sassari, Italy. Associated Member of the JRU MIRRI-IT, Italy; ² Department of Agricultural Sciences of University of Sassari, Italy

The MBDS-UNISSCC includes a collection of over 80 sourdoughs originated from different areas of Sardinia and various types of bread. Three of these sourdoughs, namely SD81, SD82, and SD84, have been chemically, microbiologically and technologically characterized, duplicated and stored at -80 °C to assess the stability of the microbial consortium during and after storage. Each sample underwent microbiological analyses using standardized procedures, both for sampling and DNA extraction. As a physiological test, the microbiomes from the three samples were evaluated for their ability to metabolize over ninety carbon and nitrogen sources using OmniLog® PM technology. In samples SD82 and SD81, the bacterial populations prevailed over yeasts, while in SD84, the abundances were similar. In SD81, the most represented bacterial isolates belonged to *Fructilactobacillus sanfranciscensis*, followed by *Levilactobacillus brevis* and *Lactiplantibacillus plantarum*, while the prevalent yeast species were *Kazachstania pseudohumilis*, *Pichia kudriavzevi*, and *Torulaspora delbrueckii*. In SD82, all the bacteria (24 isolates) were identified as *F. sanfranciscensis*, while the yeasts belonged to *Kazachstania humilis* and *Saccharomyces cerevisiae*. SD84 shows higher biodiversity with *Pediococcus parvulus* and *P. kudriavzevi* as the predominant bacterial and yeast species, respectively. The SD81 microbiome was capable of catabolize 43% and 88.5% of the tested carbon and nitrogen source, showing a high phenotypic diversity compared to SD84 and SD82. In comparison, SD84 and SD82 microbiomes were only able to catabolise 33.4% and 29% of nitrogen sources and 38% and 8% of carbon sources, respectively. Overall, the microbiomes of samples 81 and 84 showed greater phenotypic diversity compared to that of SD82. Long-term preservation strategies based on the extraction and storage of the microbiome from the sourdoughs are currently being implemented within the SUS-MIRRI project.

References

Cocolin, L., Bevivino, A., Alduina, R., Andreolli, M., Bianco, A., Capozzi, V., Costa, C., Costanzo, M., Curtoni, A., Di Gregorio, L., Elsayed, A., Felis, G., Ferrara, M., Ferrocino, I., Fiore, A., Franzetti, A., Gaglio, R., Gallo, G., Garelo, M., Varese, C. G. (2023). STANDARD OPERATING PROCEDURES (SOPs) FOR SAMPLING OF MICROBIOME IN DIFFERENT ECOSYSTEMS. Zenodo. <https://doi.org/10.5281/zenodo.8430600>

Culturomics and microbiome-based approach for the development of NatComs as biofertilizers and biocontrol agents for sustainable agriculture: the reinforcement of ENEA microbial culture collection.

Sbarra, Federico^{1,2}; Visca, Andrea¹; Sevi, Filippo¹; Tabacchioni, Silvia¹; Colantoni, Eleonora¹; Di Gregorio, Luciana¹; Costanzo, Manuela¹; Aracri, Benedetto¹; Del Fiore, Antonella¹; Varese, Giovanna Cristina²; Bevivino, Annamaria¹

¹ Department for Sustainability, Italian National Agency for New Technologies, Energy and Sustainable Economic Development, ENEA Casaccia Research Center, Italy; ² Department of Life Sciences and System Biology (DBIOS), University of Turin, Italy

Beneficial microorganisms can be used to increase crop yield and/or to reduce the damage caused by pathogens and pests in agricultural fields. Synthetic microbial communities (SynComs) fail to mimic the natural composition, the diversity and the ecological functions found in the original ecosystem. In the present work, a combined culturomics and metagenomics approach was explored to define the core microbiome of two composite rhizosphere samples from strawberry (*Fragaria* spp.) plants grown in field and to identify strains with plant-growth promoting (PGP) and biocontrol activity for the set-up of tailored Natural microbial communities (NatComs). The total DNA of the microbial community was extracted and used as template for PCR targeting the region ITS1-ITS4 for eukaryotic samples and V3-V4 for prokaryotic samples. The raw sequences were filtered and denoised and the ASVs were taxonomically assigned using Silva 138.1 database. Microbial community growth strategy was investigated by means of total microbial count, enumeration of copiotrophs and oligotrophs and eco-physiological (EP) index. A total of 159 bacteria, 57 putative N-fixing bacteria and 14 fungi with different morphologies were isolated. Bacterial isolates were characterised for siderophore production, using chrome azurol S (CAS) agar plates, and in-vitro antifungal activity. A total of 109 strains were identified as siderophore producers based on the halo thickness around the colonies. Only one fungal strain resulted positive to P-solubilisation assay, using Pikovskaya's medium, and nine strains resulted positive to the PCR amplification of *nifH* gene. Overall, a total of 119 out of 227 isolates with PGP activities were selected and will be identified by 16S/ITS sequencing and MALDI-TOF technology. The well-characterised strains will be included in the database of ENEA Microbial culture collection and exploited for the set-up of NatComs.

Acknowledgements

The authors thank Luca Cocolin for his valuable support, and Davide Spadaro, Marco Garelo and Francesco Aloï for providing rhizosphere samples. This activity has received funding from the SUS-MIRRI.IT Project "Strengthening the MIRRI Italian Research Infrastructure for Sustainable Bioscience and Bioeconomy", Project code IR00000005, CUP D13C22001390001.

Trichoderma asperellum and *Bacillus* spp. antifungal activity evaluation against *Fusarium* fungi

Senkovs, Maris¹; Jauga, Kristians²; Nikolajeva, Vizma³; Petriņa, Zaiga¹

Researcher¹; student²; associate professor³

The aim of the study was to investigate the antifungal activity of *Trichoderma asperellum* MSCL 309 as well as *B. subtilis* MSCL 897, *B. subtilis* MSCL 1441 and *Priestia* (*Bacillus*) *megaterium* MSCL 49 against twelve phytopathogenic *Fusarium* spp.. *Fusarium* isolates previously isolated from oats was evaluated in three independent experiments involving double culture, volatile and dissolved compound detection methods. In a fourth experiment, the effect of *T. asperellum* solutes on the growth of *Fusarium* isolates was tested. In the double culture method, *T. asperellum* inhibited *Fusarium* strains AS 181401 and AS 1809025 most actively, and AS 070701 strain least actively. The highest antifungal activity of *T. asperellum* liquids was found against *Fusarium* AS 181202S, *Fusarium* AS 181501S, *Fusarium* AS 181401S, *Fusarium* AS 70701, *Fusarium* AS 071301S and, *Fusarium* AS 70402 isolates. No significant effect of volatiles from *T. asperellum* strain on any *Fusarium* isolate was detected. *B. subtilis* strain MSCL 1441 inhibited the growth of all *Fusarium* spp. isolates, while *B. subtilis* MSCL 897 and *P. megaterium* did not inhibit the growth of any *Fusarium* isolates.

The use of microbial inoculants for the control of *Fusarium* related diseases in durum wheat

Alberoni, Daniele¹; Cappelletti, Eleonora¹; Cali, Martina¹; Amaral Carneiro, Greice¹; Rabbi, Francesca¹; Prodi, Antonio¹; Di Gioia, Diana¹

¹ Department of Agricultural and Food Science, University of Bologna, Italy

Biocontrol to counteract pathogens in agricultural crops with the use of selected microorganisms represents a sustainable alternative to pesticides. Durum wheat, a significant cereal crop globally, faces threats from diseases like *Fusarium* Foot Rot (FFR), *Fusarium* Crown Rot (FCR), and *Fusarium* Head Blight (FHB). These diseases, induced mainly by *Fusarium culmorum*, *Fusarium graminearum*, and *Fusarium pseudograminearum*, cause yield and quality losses [1]. Current farming defenses rely on agronomic practices and fungicides. However, due to resistance phenomena, environmental impact and toxicity towards consumers, greener alternatives are looked for. In this work, about 90 bacterial strains, 75 of which isolated from durum wheat, were screened in vitro against *Fusarium* strains using different microbiological assays. Promising candidates mainly belonging to the *Bacillus* and *Pseudomonas* genera and Lactic Acid Bacteria (LAB) were selected for further studies. In vitro seed priming using the selected strains of treated seeds proved effective against FCR/FFR with also positive influence on the germination capacity. In controlled environment trials, a LAB strain was able to *F. culmorum* at crown level, stimulating plant growth. Open-field trials treatments with the selected microorganisms exhibited positive effects in FHB control. Metabolomic profiling underscored the presence of antifungal compounds, showing the synthesis of surfactin and fengycin by *Bacillus* species, whereas benzoic acid and hydrocinnamic acid were identified in *Lactobacillus* strains. Overall, this biocontrol approach provides valuable insights into the applications of beneficial bacteria for managing *Fusarium*-related diseases in durum wheat.

References

- [1] Bankina, B., Bimšteine, G., Neusa-Luca, I., Kaņeps, J., Roga, A., and Fridmanis, D. (2020). Wheat stem base diseases and their causal agents. In IOP Conference Series: Earth and Environmental Science (Vol. 578, No. 1, p. 012001). IOP Publishing.

Acknowledgements

Research was funded under the National Recovery and Resilience Plan (PNRR) Prin 2022 “Filling gaps in the biological control of *Fusarium* diseases of durum wheat and of related mycotoxins (BICONTRARIUM)”, code 2022HAF42 CUP J53D23018320001

Session 6 – Microbes and Citizen Sciences: through the lens of public opinion

Thursday 19th September, 4:00pm

Chairs: Moretti Antonio (Italy), Perrone Giancarlo (Italy).

The Isala project: characterizing the female microbiome through citizen science

Wittouck, Stijn¹

¹ *Department of Bioscience Engineering, University of Antwerp, Antwerpen, Belgium*

The composition of the vaginal microbiome is known to be associated with several aspects of women's health and reproduction. Studies on the vaginal microbiome have until recently been mostly conducted in a clinical setting, which limits attainable sample sizes and which may create various biases in the sample. One way to overcome these limitations is a citizen science approach, where participants take their own samples in the comfort of their home. However, such an approach comes with its own set of challenges concerning recruitment, logistics, quality control and communication. In 2020, we launched Isala: a citizen science project aiming to study the female microbiome of women from the general population of Flanders, Belgium. For the first phase of the project, participants were asked to take swabs from their vagina and some other body sites and fill in a very extensive questionnaire. Recruitment was handled by launching a general call through social and regular media channels and by framing the project in a larger societal context surrounding women's health. A web designer and a graphic design firm were hired to create the sampling kits, promotion material and a professional website. Samples were transported through the Belgian postal office, after a pilot study showed no adverse effects on sample quality. The highlights of the results were communicated through various Belgian media, while an abstracted version of participant's personal microbial profiles were sent to them through the website. After the initial call for participants, more than 6,000 women registered in a single week, while quality control of the microbiome samples indicated a very high overall sample quality. The data has already led to various insights on the vaginal microbiome of women in Flanders, associations with lifestyle and other factors and co-occurrences between vaginal taxa. Follow-up projects are currently underway and are yielding more specific insights.

The Isala project was created by a team of researchers led by professor Sarah Lebeer at the University of Antwerp and was also made possible by some academic and non-academic partners. All are listed at <https://isala.be/en/about/>.

Mediterranean Diet and microbes: an emerging target for educational paths and citizen science

Fragasso, Mariagiovanna¹; Omri, Ghofrane¹; Diaferio, Antonietta¹; Spano, Giuseppe¹; Capozzi, Vittorio²

¹ Department of Agricultural Sciences, Food, Natural Resources and Engineering, University of Foggia, Foggia, Italy; ² Institute of Sciences of Food Production of National Research Council (ISPA-CNR), Foggia, Italy

The Mediterranean Diet represents a dietary model, but more generally, a set of good lifestyle practices, which receives particular attention for its impact on well-being in childhood, adult life and ageing. While the diversity of foods of plant origin (i.e. legumes, vegetables) are generally well correlated with the strengths of the Mediterranean Diet, little emphasis has usually been given to aspects relating to microbial resources, i.e. fermented foods, starter cultures, and dietary microbes. The intake of fermented foods, bio-molecules of microbial origin and microbes in the diet represent an area that is receiving growing interest in various fields of science. Here, we present an overview of Mediterranean fermented products, highlighting the interesting aspects of the main categories. Regarding the categories, recent works are presented that promote knowledge in food microbiology as topics to promote educational paths at school and university levels, as well as citizen science experiences. Finally, the experience carried out in schools is reported, underlining how the microbiological aspects represent a topic that allows the Mediterranean Diet to be promoted in a new light to students and citizens.

Acknowledgements
This work was completed as part of the bEat project (We all eat microbes: diet as reservoir of microorganisms that preserve the ecosystem services of the human gastrointestinal microbiota) supported within the call “Progetti di Ricerca di Rilevante Interesse Nazionale (PRIN) 2022” granted by the Italian Ministry for Universities and Research (MUR) (Prot. 2022TF9AHZ, CUP G53D2300574 0006). Vittorio Capozzi was funded by CNR project “NUTRAGE FOE2021 DBA.ADO05.225”. Mariagiovanna Fragasso is supported by the European Union NextGeneration EU funded ‘Agritech National Research Centre for Agricultural Technologies’ [PNRR, CN00000022]

Didactic Proposals in Education for Sustainable Development in Teaching in Portugal and Brazil: Circular Relationship between Climate Change, Healthy Ecosystems and Soil Microbial Diversity

Ricardo, Cruz M.¹; Lima, Nelson²

¹ CIEC Research Center for Child Studies, University of Minho, Braga, Portugal; ² MUM Micoteca da Universidade do Minho (MUM), Centre of Biological Engineering, University of Minho, and LABBELS Associate Laboratory, Braga/Guimarães, Portugal

Several educational materials, when explored in a formal or non-formal teaching milieu and mediated by a teacher or educator, become rich in meaning for the students. The co-creation of new teaching materials for Education for Sustainable Development (EDS) is, therefore, a permanent need that requires constant updating. For this study, we consider climate change, healthy ecosystems and soil microbial diversity, as societal challenges to be addressed by EDS. Additionally, anthropogenic pressure on Earth has reached an unprecedented scale, where abrupt global environmental changes can no longer be ignored. In this context, the study starts from the following research question: How to create didactic proposals for Education for Sustainable Development in children's Education in Portugal and Brazil which facilitate a circular view of the relationships between climate change, healthy ecosystems and soil microbial diversity? The methodology of this study is qualitative and comparative, based on content analysis (curriculum official programmes and textbooks) and their gaps. The first preliminary results of this analysis will be presented and show how the role of microorganisms is almost absent from the children's teaching contents. Based on the results, it subsequently, proposed that the study will be complemented with semi-structured interviews with teachers from early grade levels of education in Portugal and Brazil, to end with the co-creation of proposals for teaching activities, including manipulatives, and outdoor activities, among others, and which will be validated by the interviewed teachers. The outcome of this PhD thesis project is expected to obtain didactic proposals for children in the context of formal or non-formal education, for sustainable development.

Acknowledgements

This work has received funding from the European Union's Horizon Europe research and innovation programme under grant agreement number 101131818 (Microbes-4-Climate); by the Portuguese Foundation for Science and Technology (FCT) under the scope of the strategic funding of the CIEC (UIDB/00317/2020), CEB (UIDB/04469/2020) units and LABBELS – Associate Laboratory in Biotechnology, Bioengineering and Microelectromechanical Systems (LA/P/0029/2020).

Session 7 – Advanced approaches in taxonomy, phylogeny and functional genomics

Friday 20th September, 9:30am

Chairs: Clermont Dominique (France), Masiello Mario (Italy).

Metabarcoding with Nanopore MinION: not only 16S or ITS

Faino, Luigi¹

¹ *Department of Environmental Biology, University of Rome Sapienza*

The microbial species identified so far account for only a small portion of Earth’s overall biodiversity. To enhance our understanding, DNA barcoding serves as a powerful tool for identifying multiple species, relying on the amplification and sequencing of conserved genetic regions. However, traditional short-read sequencing offers limited taxonomic resolution due to incomplete sequences. The advent of new long-read sequencing techniques, such as Nanopore sequencing, addresses this issue. Long-read sequencing improves species identification accuracy and increases the ability to detect low-abundance species by including larger portions of the standardized marker sequence. Despite these advances, there is a shortage of bioinformatics tools specifically designed for analyzing Nanopore sequence data compared to earlier technologies. Consequently, our study focused on establishing an amplicon-Nanopore sequencing workflow and developing a pipeline capable of processing long, noisy reads. To evaluate and compare our pipeline with existing methods, we amplified and sequenced the 16S and ITS genes of various known microorganisms with differing taxonomic similarities using Nanopore technology. The resulting sequences were analyzed using both our pipeline and available software. Our results demonstrated precision and recall rates close to 1, comparable to the Emu software. Additionally, we tested the flexibility and reliability of our method by applying it to other selected genetic markers to achieve detailed identification at the pathovar/subspecies level for quarantine bacterial pathogens. Our approach proved promising for the precise identification of microorganisms at the genus-species level for broader metabarcoding studies and deeper taxonomic resolution, such as pathovar or sequence type.

Genomic overview over 100 cyanobacterial strains: taxonomy, microbiome and biosynthetic gene clusters

Luz, Rúben^{1,2}; Cordeiro, Rita²; Borges, Inês¹; Fonseca, Amélia^{1,2}; Gonçalves, Vitor^{1,2}

¹ Universidade dos Açores, Faculdade de Ciências e Tecnologia, Ponta Delgada, Portugal; ² CIBIO, Centro de Investigação em Biodiversidade e Recursos Genéticos, InBIO Laboratório Associado, BIO-POLIS Program in Genomics

In the Bank of Algae and Cyanobacteria of the Azores, there are more than 800 non-axenic strains of microalgae and cyanobacteria, and more than 400 correspond to cyanobacterial strains. In the last year, five genera and ten species have been described based on the polyphasic approach increasing the available data for genetic studies, but lacking genomic data. To better support further taxonomic studies, more than 100 non-axenic strains have been sequenced by Illumina. Most of the cyanobacterial genomes produced are of high quality with >99% completeness and <1% contamination. Although the objective of this study was the cyanobacterial genomes, many bacteria associated with the cyanobacterial cultures were also sequenced, providing high-quality genomic data and allowing the characterization of the microbiome associated with the cyanobacterial organism. This data allows for higher quality taxonomic studies of cyanobacteria in several problematic clades, such as the *Prochlorococcaceae*, and complements biased single-gene phylogenies. It can also support targeted studies by taking into account genomic analysis and the diversity of biosynthetic gene clusters (BGCs), not only of the cyanobacteria but also by the associated microbiome with novel genomic information. The genomic study of non-axenic cyanobacterial strains can provide many benefits, not only by studying the target organism but also the associated microbiome, which is likely to contain many unculturable bacteria that represent unique resources for taxonomic, conservation and biotechnological studies.

Acknowledgements

Rúben Luz was supported by a PhD. grant (M3.1.a/F/002/2020) from the Fundo Regional da Ciência e Tecnologia (FRCT).

This work was funded by FEDER funds through the Interreg-MAC 2014-2020 Program under the projects REBECA—Red de excelencia en biotecnología azul (algas) de la región macaronesia (MAC1.1a/060) and REBECA-CCT—Red de Excelencia en Biotecnología Azul de la Región Macaronésica. Consolidación, Certificación y Transferencia (MAC2/1.1b/269).

By Portuguese National Funds, through FCT—Fundação para a Ciência e a Tecnologia, the European Union, QREN, FEDER, COMPETE, by funding the CIBIO/InBIO (project UID/BIA/ 50027/2013 and POCI-01-0145-FEDER-006821).

Drivers of aquatic fungal diversity and ecological functions throughout Europe: Biodiversa+ FUNACTION and MoSTFun projects

Marchese, Pietro¹; Garzoli, Laura¹; Böhm M.² Calore R.¹²; Canteiro C.²; Costa S.^{3,4}; Denfeld B.^{5,15}; Eckert E.M.¹; Fernandes I.^{3,4,17}; Ferrari E.¹; Fontaneto D.¹; Frainer A.⁶; Ganzert L.⁷; Graça D.^{3,4}; Garcés E.⁸; Grossart H.P.^{7,14,17}; Kisand V.⁹; Mammola S.¹; Millette K.¹⁶; Panksep K.⁹; Pinnow S.⁷; Prins V.⁹; Rämä T.¹¹; Reñé A.⁸; Retter A.⁷; Rogora M.¹; Romero-Mujalli D.¹²; Sousa R.^{3,4}; Tedersoo L.¹³; Vinnere Pettersson O.¹⁰; Wang Z.⁵; Anderson J.^{5,17}; Bruder A.^{12,17}

¹ National Research Council of Italy, Water Research Institute CNR-IRSA of Verbania, Italy. ² Global Center for Species Survival, Indianapolis Zoo, USA. ³ University of Minho, Centre of Molecular and Environmental Biology, Portugal. ⁴ University of Minho, Institute of Science and Innovation for Bio-Sustainability, Portugal. ⁵ Swedish University of Agricultural Sciences, Department of Aquatic Sciences and Assessment, Sweden. ⁶ Norwegian Institute for Nature Research (NINA), Norway. ⁷ Leibniz Institute of Freshwater Ecology and Inland Fisheries (IGB), Germany. ⁸ Institut de Ciències del Mar CSIC. Dept. Biologia Marina i Oceanografia, Spain. ⁹ University of Tartu, Institute of Technology, Estonia. ¹⁰ Uppsala University, SciLifeLab, Sweden. ¹¹ Faculty of Biosciences, Fisheries and Economics, UiT The Arctic University of Norway, Norway. ¹² Institute of Microbiology, University of Applied Sciences and Arts of Southern Switzerland, Switzerland. ¹³ Mycology and Microbiology Center, University of Tartu, Estonia. ¹⁴ Potsdam University, Institute of Biochemistry and Biology, Germany. ¹⁵ Swedish Infrastructure for Ecosystem Science (SITES), Uppsala, Sweden. ¹⁶ GEOBON secretariat, Montreal (CA). ¹⁷ IUCN SSC Aquatic Fungi Specialist Group, IUCN Species Survival Commission, Gland, Switzerland

Aquatic fungi (AF) and fungal-like organisms are ecologically diverse, play key functions in freshwater and marine ecosystems and provide numerous ecosystem services and disservices. Despite their relevance, quantitative evidence about AF diversity and functions is still scarce. The panEuropean FunAqua project, led by our Estonian team, implemented eDNA metabarcoding to explore the taxonomic, phylogenetic, and functional diversity of AF at the global scale. Capitalizing on this experience, we developed two transnational projects within the Biodiversa+: FUNACTION and MoSTFun.

FUNACTION aims to fill knowledge gaps in AF ecology and develop conservation guidelines based on AF diversity recorded across Europe. FUNACTION will implement a modeling approach to map drivers of AF diversity change and identify priority areas for conservation within the EUBiodiversity Strategy for 2030. Members of the project, together with worldwide AF experts, also initiated the Aquatic Fungi Specialist Group of the IUCN Species Survival Commission (SSC), aimed at coordinating the first conservation assessment, planning and action efforts for AF. MoSTFun aims to integrate AF into practicable monitoring procedures based on Essential Biodiversity Variables and open accessibility of sequencing and environmental data in FAIR databases. MoSTFun will test novel approaches including -omics technologies, Earth Observation, GIS, and species distribution modeling leveraging on existing biomonitoring programs and networks to reanalyze samples and evaluate procedures to include AF in routine biodiversity monitoring.

Both projects, by harmonizing data and expertise on AF with existing initiatives, will set the stage for broad exchanges with relevant stakeholders. This is a fundamental step to change people's perception about this neglected, yet highly diverse group of organisms and facilitate initiatives to preserve healthy aquatic ecosystems for future generations.

ITSoneDB v1.144 and BioMaS@ITSoneWB: two ELIXIR-IT main resources for amplicon based mycobiome investigation

Defazio, Giuseppe¹; Tangaro, Marco Antonio²; Pesole, Graziano^{1,2,3}; Fosso, Bruno¹

¹ Department of Biosciences, Biotechnologies and Environment, University of Bari, Bari, Italy; ² National Research Council (CNR) – Institute of Biomembranes, Bioenergetics and Molecular Biotechnologies; ³ Consorzio Interuniversitario Biotechnologie, Trieste, Italy

DNA-metabarcoding represents the cheaper and faster way to explore and profile the composition of microbial communities colonizing several habitats, such as soil and water, as well as animal hosts niches as human skin and livestock guts. Profiling the microbiota taxonomic composition is crucial for understanding how microorganisms may influence the overall state of the investigated habitats.

To date, the exploration of the so called mycobiota (i.e. fungal components of microbiota) is a constantly growing field and ITS1 metabarcoding is one of the most reliable approach [1]. The choice of the most appropriate bioinformatic workflow is a key point to obtain reliable and interpretable results.

In this scenario, we present ITSoneDB v1.144 [2] an ITS1 sequence database and BioMaS at ITSoneWB [3] a bioinformatic resource for Illumina paired end reads sequence analysis and taxonomic classification. We present a comparative assessment of ITSoneDB versus UNITE [4] as well as of BioMaS@ITSoneWB versus QIIME2 [5]. Shortly QIIME2 relies on the inference of Amplicon Sequence Variants (ASV) by wrapping DADA2 algorithm, while BioMaS [6] is an ASV and Operational Taxonomic Unit (OTU) free approach particularly suitable for barcodes with a highly variable length as ITS1 (i.e. 100-1,000 nucleotides long).

The results of the comparative assessment, carried out on a known microbial mock community, will be presented providing evidences of the relative performances of the tested combinations of reference databases (ITSoneDB and UNITE) and classification tools (BioMaS and QIIME2).

References

- [1] Wang et al. 2015
- [2] Santamaria et al. 2018. «ITSoneDB: A Comprehensive Collection of Eukaryotic Ribosomal RNA Internal Transcribed Spacer 1 (ITS1) Sequences». NAR
- [3] Tangaro et al. 2021. «ITSoneWB: profiling global taxonomic diversity of eukaryotic communities on Galaxy». Bioinformatics
- [4] Kõljalg et al. 2005. «UNITE: A Database Providing Web-Based Methods for the Molecular Identification of Ectomycorrhizal Fungi: Methods». New Phytologist
- [5] Bolyen et al. 2019. «Reproducible, Interactive, Scalable and Extensible Microbiome Data Science Using QIIME 2». Nature Biotechnology
- [6] Fosso et al. 2015. «BioMaS: A Modular Pipeline for Bioinformatic Analysis of Metagenomic AmpliconS». BMC Bioinformatics

The new challenge in virus taxonomy: a binomial nomenclature for virus species

Rubino, Luisa¹

¹ *Consiglio Nazionale delle Ricerche, Istituto per la Protezione Sostenibile delle Piante, Bari, Italy*

The International Committee on Taxonomy of Viruses (ICTV) is responsible for developing and maintaining an internationally agreed system of hierarchical classification of viruses and naming of taxa, available to the scientific community through a website (<https://ictv.global/>) and published reports/papers. Over the years the ICTV has adapted its taxonomic framework to reflect current knowledge on the evolution of global virosphere. The ICTV now allows the classification of viruses known only from genomic data. Taxonomic ranks have been expanded beyond orders to 15 hierarchical ranks, paving the way to comprehensive studies on evolutionary connections of viruses and bringing virus taxonomy closer to other biological taxonomies. Species naming in all these taxonomies follow a Latinized binomial format (i.e., binomial nomenclature) first introduced by Carl Linnaeus in 1753, consisting of two italicized words indicating the genus ("genus name") and the species ("specific name/species epithet"), respectively. Typical examples of binomial species names are *Arabidopsis thaliana*, *Saccharomyces cerevisiae*, *Homo sapiens*, or *Escherichia coli*. On the contrary, a uniform format was not enforced for viral species names, which overlapped those of viruses, except for a requirement to be italicized and to have the first letter of the first word capitalized. To adopt a standardized format, a binomial nomenclature for virus species was ratified by the ICTV in 2021. Thus, a virus species name consists of two italicized words, the first one being the genus name and the second one being a "free-form" species epithet, composed using the standard Latin-script English alphabet containing 26 letters and/or Arabic numbers. Linnaean-style, Latinized virus species are permitted, but not mandated. Importantly, this change in nomenclature applies to virus species only. Established species names have all been converted to binomials. Common virus names are not affected and will stay unchanged.

Genomic insights into *Mrakia*: expanding horizons in microbial biotechnology

Di Cesare, Francesca¹; Andreani, Daniele²; Lazzari, Gioele³; Turchetti, Benedetta⁴; Parente, Eugenio⁵; Felis, Giovanna¹; Vitulo, Nicola³

¹ Department of Biotechnology, University of Verona & VUCC-DBT, University of Verona, Verona, Italy; ² Department of Agricultural, Food and Environmental Science, University of Perugia, Perugia, Italy; ³ Department of Biotechnology, University of Verona, Verona, Italy; ⁴ Department of Agricultural, Food and Environmental Sciences, Industrial Yeasts Collection DBVPG, University of Perugia, Perugia, Italy; ⁵ Department of Agricultural, Forestry, Food and Environmental Sciences, University of Basilicata, Potenza, Italy

In recent years, multi-omic sciences and next-generation sequencing (NGS) technologies have revolutionized microbiological research and microbial ecology, making DNA sequencing faster and more affordable [1]. This has enabled the examination of various ecosystems and the development of genetic tools for non-conventional yeasts (NCY), though the field is still in its early stages [2].

This study aims to define an efficient genomic analysis pipeline for NCY, non-model organisms, with a focus on the *Mrakia* genus, comprising 30 species belonging to the phylum Basidiomycota and of environmental origin. Interestingly, some species/strains display fermentation abilities suggesting biotechnological potential in fermented beverages, evidenced by successful tests on *Mrakia* gelida in beer prototype production [3,4].

In this context, producing high-quality genomic and transcriptomic sequences for different *Mrakia* species allows for a more comprehensive understanding of their genetic and functional bases, thus uncovering hidden potential in the fermentation sector and beyond.

Currently, aiming to sequence all *Mrakia* type strains, we obtained Oxford Nanopore long reads and Illumina short reads for 7 accessible species. To obtain high-quality genome assemblies, we tested 3 assembly and correction tools, selecting the best combination based on the assembly statistics. Additionally, repetitive sequence masking was performed. The average nucleotide identity between species genomes was calculated to assess phylogenetic relationships. Gene prediction will be conducted on the assembled genomes by means of transcriptome data to identify biotechnological fermentation markers.

Future plans involve broadening genomic and transcriptomic data to include 11 more *Mrakia* species, improving gene annotation, and enhancing taxonomic analysis. In conclusion, this study aims to deepen our understanding of unexplored *Mrakia* species, maximizing their potential in microbiological and biotechnological fields.

References

- [1] Cao et al., 2017, 10.3389/fmicb.2017.01829
- [2] Geijer et al., 2022, 10.1093/fem-syr/foab071
- [3] Turchetti et al., 2023, 10.1016/j.foodres.2023.113004
- [4] Linnakoski et al., 2023, 10.3389/fmicb.2023.1108961

Acknowledgments

Study carried out within the PRIN 2022 PNRR Project “NCYdiversity” P20229JMMH, funded by European Union Next-GenerationEU, CUP C53D23007560001 (PIANO NAZIONALE DI RIPRESA E RESILIENZA (PNRR)—MISSIONE 4 COMPONENTE 2, INVESTIMENTO1.4—D.D. 1364/2023)

New Enterobacterales species isolated from dead seals in Antarctica

Teturová, Kateřina¹; Koublová, Vendula¹; Staňková, Eva¹; Sedláček, Ivo¹

¹ Czech Collection of Microorganisms, Department of Experimental Biology, Faculty of Science, Masaryk University, Brno, Czech Republic

One of the goals of Antarctic research is to monitor and preserve microbiological diversity in animals and the environment.

Among samples collected on James Ross Island within the Czech Antarctic Research Programme, two strains isolated from dead seals, P4252 and P6696, were not identified as known species using phenotyping methods and, therefore, chosen for further molecular analysis. These two strains were gram-negative, rod-shaped, fermenting, psychrotolerant, and moderately halotolerant bacteria.

Sequencing of the 16S rRNA gene revealed that the closest hits for strain P4252 were *Rahnella inusitata* with 98.41 % sequence similarity and *Rouxiella badensis* with 98.32 % similarity. The gene sequence of strain P6696 showed the highest similarity to *Moellerella wisconsensis* with 98.58% and *Providencia burhodogranariae* with 98.08 %.

Next, whole-genome sequences were obtained, and the Digital DNA-DNA Hybridization (dDDH) analysis was performed using the TYGS pipeline by DSMZ. The dDDH values were 23.6 % for *Ewingella americana* as the closest to strain P4252, and 22.7 % for *M. wisconsensis* as the nearest to strain P6696. These values are below the accepted threshold indicating that the studied strains belong to potential new species of the order Enterobacterales. The Average Nucleotide Identity (ANI) values were 81.08 % between strain P4252 and *E. americana*, and 76.44 % for strain P6696 and *M. wisconsensis* which implies that the isolates represent novel species within one-species genera of *Ewingella* and *Moellerella*.

The analyses based on whole-genome sequences are more robust, thus the results of dDDH and ANI are more reliable than comparing the similarities of the 16S rRNA gene only.

Acknowledgments

The work was supported by the Czech Antarctic Research Programme (CARP) and the Czech Collection of Microorganisms (CCM).

Taxonomic classification of Metagenome-Assembled Genomes with kMetaShot

Defazio, Giuseppe¹; Tangaro, Marco Antonio¹; Pesole, Graziano^{1,2,3}; Fosso, Bruno¹

¹ Department of Biosciences, Biotechnologies and Environment, University of Bari, Bari, Italy; ² National Research Council (CNR) – Institute of Biomembranes, Bioenergetics and Molecular Biotechnologies; ³ Consorzio Interuniversitario Biotechnologie, Trieste, Italy

The investigation of environmental microbiome such as soil, water and air play an essential role to deepen biodiversity knowledge and compositional profiles in pollution context [1]. To unveil microbiome role, it is essential to both unveil its taxonomic composition and assess its genetic potential reconstructing the component genomes as Metagenome-Assembled Genomes (MAG) [2]. To this aim, we developed kMetaShot, an accurate tool for MAGs taxonomic classification.

kMetaShot relies on kmer/minimizer counting and consists of two modules: the reference generator counts minimizers from RefSeq Bacterial and Archaea RNAs and stores relevant minimizers (i.e. exclusively shared among strains of the same genus) in a dedicated reference matrix; the classification module processes MAGs, performs the minimizer counting and queries the reference matrix to achieve the taxonomic classification relying on a prevalence-based approach. kMetaShot has been benchmarked against two of the most cited comparable tools, i.e. GTDBtk [3] and CAMITAX [4] by classifying MAGs obtained by 20 rhizosphere in silico generated samples of the CAMI2 [5].

kMetaShot outperformed both GTDBtk and CAMITAX in term of precision at species (80%, 25%, 30%, respectively) and genus (100%, 90%, 55%, respectively) levels. Differently from other tools, kMetaShot is able to perform classification at strain level with a Balanced-Accuracy at ~63%. kMetaShot is also the fastest and the most computationally effective.

kMetaShot is an easy-to-use, easy-to-install and computationally effective tool for MAGs taxonomic classification with high accuracy classification at strain level. It takes in input MAGs FASTA files and is available and documented at <https://github.com/gdefazio/kMetaShot> and distributed as Conda package and Docker container.

References

- [1] Beale et al. 2017. «A Community Multi-Omics Approach towards the Assessment of Surface Water Quality in an Urban River System». International Journal of Environmental Research and Public Health
- [2] Quince et al. 2017. «Shotgun Metagenomics, from Sampling to Analysis». Nature Biotechnology
- [3] Chaumeil et al. 2020. «GTDB-Tk: A Toolkit to Classify Genomes with the Genome Taxonomy Database». Bioinformatics
- [4] Bremges et al. 2020. «CAMITAX: Taxon labels for microbial genomes». GigaScience
- [5] Meyer et al. 2022. «Critical Assessment of Metagenome Interpretation: The Second Round of Challenges». Nature Methods

Presumptive identification of two Gram-positive microaerophilic strains isolated from Weddell's seals in Antarctica.

Nováková, Dana¹; Sedláček, Ivo¹

¹ Department of Experimental Biology, Czech Collection of Microorganisms, Faculty of Science, Masaryk University, Czech Republic

The microbial biodiversity of bacteria inhabiting mucous membranes in the mouths of various animals in Antarctica is being investigated within the Czech Antarctic Research Programme (CARP). An analysis of palindromic PCR profiles of repetitive elements, carried out with the primer (GTG)₅, was performed initially on a group of 100 microaerophilic Gram-positive bacterial isolates. The profiles of two strains isolated from the oral cavity of Weddell's seals (*Leptonychotes weddellii*) during the Antarctic summer of 2021 showed fingerprints that differed significantly from all other profiles in the in-house database. Both strains, P13308 and P13329, were catalase-negative, microaerophilic, -hemolytic cocci belonging to Lancefield serological group C. The partial 16S rRNA gene sequence of the strains analyzed showed the highest similarity to the *Streptococcus pacificus* type strain CSL7591T. As the calculated similarity of the 16S rRNA sequences (98.5 %) was below the recommended threshold for species delimitation, there is a strong indication that the strains studied represent a novel species within the genus *Streptococcus*.

Acknowledgements

This work was supported by the CARP project and in part by the Czech Collection of Microorganisms.

New protocol development of bioinformatics analysis for species verification of the genus *Pseudomonas* from Polish Collection of Microorganisms resources.

Kostrzewa, Maja¹; Korzeniowska-Kowal, Agnieszka¹; Cieniuch, Gabriela¹; Grzywa, Renata²; Gamian, Andrzej¹

¹ Polish Collection of Microorganisms, Hirsfeld Institute of Immunology and Experimental Therapy, Wrocław, Poland; ² Faculty of Chemistry, Wrocław University of Science and Technology, Wrocław, Poland

In the case of bacteria from the genus *Pseudomonas*, verification methods considered as the goldstandard, such as 16S rRNA gene sequence analysis and MALDI-TOF MS analysis, prove to be insufficient. The 16S rRNA sequence does not exhibit a significant degree of nucleotide polymorphism, making it difficult to accurately differentiate species within the genus. Additionally, MALDI-TOF MS analysis is inadequate due to the high similarity of protein profiles and the phenotypic variability of some species caused by environmental factors. Thus, more advanced methods are necessary to precisely identify bacteria from the genus *Pseudomonas*. Multi-Locus Sequence Analysis (MLSA) method relies on the analysis of multiple housekeeping genes, providing higher resolution of phylogenetic relationships of species within the genus.

The main aim of this study was to develop a protocol utilizing molecular biology methods and bioinformatics tools for the effective species identification of bacteria from the genus *Pseudomonas* deposited in the Polish Collection of Microorganisms (PCM).

Materials for the study included 11 bacterial strains from PCM: 4 reference strains and 7 strains for verification.

The methods used for initial identification included MALDI Biotyper and 16S rRNA sequencing. Further identification employed MLSA based on the housekeeping genes: *gyrB*, *rpoD*, *rpoB*, and *andrecA*. Phylogenetic analysis was conducted using the Maximum Likelihood method.

The new protocol involving MLSA combined with phylogenetic analysis confirmed the identity of strains: PCM 1411 *P. syringae*, PCM 2856 *P. silesiensis* and PCM 2904 *P. laurysulfatovorans*. Strains PCM 1994 *P. fluorescens*, PCM 2124 *P. putida*, and PCM 2219 *Pseudomonas* sp. are proposed for reclassification to new species. PCM 2123 *P. fluorescens* was not clearly classified, so Whole Genome Sequencing is necessary in this case.

The obtained results proved that proposed protocol allows for more accurate identification of *Pseudomonas* species.

Novel insights into heterocyte patterning and proheterocyte germination in *Kaarinaea lacus* PCC 9237T gen. nov. sp. nov.

Saraf, Aniket¹; Blondet, Eddy¹; Boullie, Anne¹; Criscuolo, Alexis²; Gugger, Muriel¹

¹ Collection of Cyanobacteria, Département de Microbiologie, Institut Pasteur, Université Paris Cité, Paris, France; ² GIPhy Genome Informatics and Phylogenetics, CRBIP, Institut Pasteur, Université Paris Cité, Paris, France

Cyanobacteria are oxygenic photosynthetic prokaryotes and are broadly grouped into unicellular and filamentous forms. Certain filamentous cyanobacteria develop specialized cells called heterocyte that perform nitrogen fixation. These cells are terminally differentiated that not only differ functionally but also morphologically and physiologically from the other vegetative cells. Heterocyte differentiation is a complex and highly regulated process involving initiation, commitment, patterning, morphogenesis and maturation stages [1,2]. Among the heterocytous cyanobacteria, the genus *Nostoc* is one of the most widely studied genera. In this study, we conducted an in-depth characterization of a toxic *Nostoc* sp. 9237T isolated in 1986 from a freshwater bloom in Finland. Multilocus and 16S rRNA gene phylogenetic trees both confirmed that PCC 9237T represents a novel *Nostoc* genus for which we propose the name *Kaarinaea lacus* gen. nov., sp. nov. Remarkably, PCC 9237T showed interesting morphological attributes such as unusual heterocyte patterning leading to the development of heterocytes in series followed by the fragmentation of trichomes at this site. This cyanobacterium displayed occasional proheterocyte germination, a morphological feature last reported more than five decades before [3,4]. We conducted a detailed genomic and morphological examination to gain a better understanding of these phenomena. Furthermore, the genomic investigation for biosynthetic gene clusters (BGCs) revealed a significantly different heterocyte glycolipid (HG) BGC in PCC 9237T compared to the well-studied HG BGC of the model organism PCC 7120 [5]. In conclusion, the results obtained indicate *Kaarinaea lacus* PCC 9237T as a potential novel model organism to study proheterocyte germination and gain further insights into heterocyte patterning and differentiation.

References

- [1] Muro-Pastor, A.M., Hess, W.R., 2012. Heterocyst differentiation: From single mutants to global approaches. *Trends Microbiol* 20(11), 548-557.
- [2] Zeng, X., Zhang, C.C., 2022. The making of a heterocyst in Cyanobacteria. *Annu Rev Microbiol* 76, 597-618.
- [3] Geitler, L., 1921. Versuch einer Lösung des Heterocysten-Problems. *Sitz. - Ber. K. Akad. Wiss., math. -naturwiss. Kl.* 130, 223-246.
- [4] Wilcox, M., Mitchison, G.J., Smith, R.J., 1973a. Pattern formation in the blue-green alga *Anabaena*. II. Controlled proheterocyst regression. *J Cell Sci* 13, 637-649.
- [5] Fan, Q., et al., 2005. Clustered genes required for synthesis and deposition of envelope glycolipids in *Anabaena* sp. strain PCC 7120. *Mol Microbiol* 58, 227-243

Exploring *Fusarium oxysporum* biodiversity in Beauvericin gene-cluster

Gatto, Angela¹; Moretti, Antonio¹; Susca, Antonia¹

¹ Institute of Sciences of Food Production, National Research Council (CNR-ISPA), Bari, Italy

Fusarium oxysporum is classified as the fifth most important fungal plant pathogen in the world, is considered a species complex (FOSC), that includes over 106 formae speciales, known to infect more than 100 different hosts where it causes vascular wilt. It can also produce a wide range of active metabolites, among which the mycotoxin beauvericin [1,2].

The biosynthetic gene cluster for beauvericin (Bea) consists of 4 genes: Bea1, encoding NRPS22, the non-ribosomal peptide synthase responsible for the synthesis of the beauvericin backbone, Bea2, Bea3 and Bea4, encoding proteins with transport and regulatory functions. A special role was discovered for the ABC transporter (Bea3): deletion of the coding gene led to a significant upregulation of Bea1 and Bea2 and to drastically increase product yields [3]. Understanding the genetic basis of beauvericin biosynthesis may provide insights into the regulation of biosynthesis, the potential for genetic manipulations and the development of strategies to control the production of beauvericin in food and feed crops.

In this study, we analysed the biosynthetic gene cluster in *F. fujikuroi* and in 4 isolates of *F. oxysporum* f.sp. melonis that produce or do not produce beauvericin and originate from two different areas: specifically, California in USA and Sicily in Italy. Bioinformatic comparison between *F. fujikuroi* and *F. oxysporum* in the BEA gene cluster revealed some differences in coding conserved regions, possibly non-affecting beauvericin biosynthesis. Specifically, in Bea3 an extra 56-bp region in exon 6 of *F. fujikuroi* is predicted as intron in *F. oxysporum*, generating an extra exon. Moreover, we found a difference in the Bea3 coding region, leading to a change in the amino acid sequence, apparently linked to different geographic origins of isolates.

Further studies are needed to understand whether this difference could be a marker to distinguish the origin of *F. oxysporum* isolates and to verify if it affects beauvericin production.

References

[1] Dean R, Van Kan JA, Pretorius ZA, Hammond-Kosack KE, Di Pietro A, Spanu PD, Rudd JJ, Dickman M, Kahmann R, Ellis J. The Top 10 fungal pathogens in molecular plant pathology. Mol. Plant Pathol. 2012; 13: 414–430. doi: 10.1111/j.1364-3703.2011.00783.x.

[2] Edel-Hermann V, Lecomte C. Current status of *Fusarium oxysporum* formae speciales and races. Phytopathology. 2019; 109: 512–530. doi: 10.1094/PHYTO-08-18-0320-RVW

[3] Niehaus EM, Studt L, von Bargen KW, Kummer W, Humpf HU, Reuter G, Tudzynski B. Sound of silence: the beauvericin cluster in *Fusarium fujikuroi* is controlled by cluster-specific and global regulators mediated by H3K27 modification. Environ Microbiol. 2016; 18 (11): 4282–4302. doi: 10.1111/1462-2920.13576. PMID: 27750383

Exploring cyanobacteria genomes for antithrombotic secondary metabolites

Borges, Inês¹; Cordeiro, Rita²; Luz, Rúben²; Gonçalves, Vítor²; Fonseca, Amélia²

¹ Faculdade de Ciências e Tecnologia, Universidade dos Açores, Ponta Delgada, Portugal; ² CIBIO, Centro de Investigação em Biodiversidade e Recursos Genéticos, InBIO Laboratório Associado, BIOPOLIS Program in Genomics

Cyanobacteria are photosynthetic microorganisms capable of producing secondary metabolites with relevant bioactivities to human health, such as fungicidal activities, iron chelating, and protease inhibition. Anabaenopeptins, Microviridins, and Micropeptin K139 are some of the compounds produced with antithrombotic effect, being the anabaenopeptins capable of inhibiting Thrombin Activatable Fibrinolysis Inhibitor (TAFIa). Therefore, using resources from the Bank of Algae and Cyanobacteria of the Azores (BACA), the main goal of this work was to search for compounds with anticoagulant activities in 50 strains of cyanobacteria, using an OMIC approach. As a result, the genome of 50 cyanobacteria strains was obtained through Illumina sequencing. Genomic and phylogenomic analysis was performed and the search for biosynthesis gene clusters (gene mining) was conducted. To date, 22 of the strains under study, belonging to the genera *Azorothrix*, *Dulcicalothrix*, and *Nostoc*, among others, have shown genes capable of producing Anabaenopeptins, Microviridins, and Micropeptin K139, indicating potential for anticoagulant activities. The confirmation of the production of these compounds will be performed through metabolomic analysis, using HRLC-MS/MS and a bioassay to determine antithrombotic activity. Given that diseases such as thrombosis are one of the leading causes of death, these results may become promising for the development of new anticoagulant drugs.

Acknowledgements

This work was funded by FEDER funds through the Interreg-MAC 2014-2020 Program under the projects REBECA—Red de excelencia en biotecnología azul (algas) de la región macaronesia (MAC1.1a/060) and REBECA-CCT—Red de Excelencia en Biotecnología Azul de la Región Macaronésica. Consolidación, Certificación y Transferencia (MAC2/1.1b/269).

By Portuguese National Funds, through FCT—Fundação para a Ciência e a Tecnologia, the European Union, QREN, FEDER, COMPETE, by funding the CIBIO/InBIO (project UID/BIA/ 50027/2013 and POCI-01-0145-FEDER-006821).

GEN-ERA toolbox: reproducible bioinformatics workflows for genomics and taxonomy in culture collection.

Lupo, Valérian¹; Baurain, Denis²; Wilmotte, Annick³; Declerck, Stéphane¹; Daniel, Heide-Marie¹; Cornet, Luc³

¹ BCCM/MUCL, Université catholique de Louvain, Louvain-la-Neuve, France; ² Eukaryotic Phylogenomics, Université de Liège, Liège, Belgio; ³ BCCM/ULC, Université de Liège, Liège, Belgio

Microbial culture collections play a key role in taxonomy, as they facilitate the exploration of strain diversity and provide well-characterized biological material for both fundamental and applied research. To enhance species delineation, microbial resource centers are now adopting advanced methods such as whole-genome sequencing and phylogenomics. In line with this, the genomic needs of the Belgian Coordinated Collections of Microorganisms (BCCM) were studied, leading to the development of the GEN-ERA toolbox. This unified cluster of bioinformatic workflows is dedicated to both bacteria and small eukaryotes (e.g., yeasts). The GEN-ERA toolbox is publicly accessible and tailored for researchers lacking specialized bioinformatics training, operable with a single command line. Hence, it facilitates all steps from genome downloading and quality assessment, including genomic contamination evaluation, to phylogenetic tree reconstruction. Additionally, it supports workflows for average nucleotide identity comparisons and metabolic modeling. To enhance reproducibility, all workflows are containerized using Singularity and managed with Nextflow. The utility of the GEN-ERA toolbox is demonstrated through two studies focusing on bacteria and yeast. The first study establishes the presence of five species within the recently described cyanobacterial genus *Laspinema*, three of which are strains held in the BCCM/ULC collection. The second study involves phylogenetic analyses of the *Starmerella* genus, where we sequenced 21 genomes from the BCCM/MUCL collection, doubling the number of publicly available *Starmerella* genomes, including two high-quality long-read PacBio sequences. The results of the two studies highlight the importance of phylogenomics and GEN-ERA in interpreting evolutionary relationships and taxonomic delineation, particularly through the use of Average Nucleotide Identity.

Comparison study between molecular fingerprinting and FTIR BIOTYPER

Soares, Célia¹; Vale Dias, Teresa¹; Venâncio, Armando¹; Lima, Nelson¹

¹ MUM Micoteca da Universidade do Minho (MUM), Centre of Biological Engineering, University of Minho, and LABBELS Associate Laboratory, Braga/Guimarães, Portugal

Within a microbial species there are different individuals carrying their own characteristics known as strains. Using the correct strain ensures consistency in biotechnological processes, leading to reliable and reproducible results. Identification of strains within a species can be done by molecular fingerprinting where it involves analysing the genome using different techniques such as RAPD and AFLP. Results can be obtained through gel electrophoresis or sequencing and creates a unique genetic fingerprint of the strain. Another technique is the Bruker's Fourier Transform InfraRed (FTIR) spectroscopy system, the IR Biotyper. FTIR spectroscopy measures the absorption of infrared light by chemical bonds and the resulting spectrum represents the molecular composition which can be used to distinguish strains based on their biochemical signatures. It is reliable and fast; however, IR Biotyper protocols were developed for bacteria. Because the reading of the plate is done with the incidence of infrared light it is essential to have a homogeneous solution and since filamentous fungi have complex, filamentous structures, ensuring a representative and homogeneous sample can be difficult.

In this sense, a filamentous fungi protocol to be used in the IR Biotyper was developed and compared with the results obtained in the molecular fingerprinting. For this, 16 strains of *Penicillium crustosum* with different origins (country and commodity) were used. Molecular fingerprinting was obtained by (GACA)₄ and M₁₃ primers amplification and the recommended protocol for bacteria was modified. Molecular fingerprinting showed that some strains created clusters according to their origins. The preliminary tests showed that the modified protocol is suitable for this species and the results of the IR Biotyper seem to be consistent with the fingerprinting results. However, additional strains are going to be analysed as proof of concept.

Acknowledgements

This work was supported by the Portuguese Foundation for Science and Technology (FCT) under the scope of the strategic funding of UIDB/04469/2020 unit and LABBELS – Associate Laboratory in Biotechnology, Bioengineering and Microelectromechanical Systems (LA/P/0029/2020). “MIRRI-PT (Polo Norte)” project (PINFRA04/84445/2020) funded by the European Regional Development Fund under Norte2020-Programa Operacional Regional do Norte allows the establishment of the molecular biology and proteomic platforms. Teresa Vale Dias acknowledges FCT for the PhD grant 2020.05849.BD.

Advancing research in synthetic biology through DTU Bioengineering's Fermentation and HT-automation facility

Iosca, Giovanna¹; Munoz Fernández, Gloria¹; Torres-Montero, Pablo¹; Durmusoglu, Deniz¹; Johansen, Tina¹; Booth, Nicky Sebastian¹; Ruiz Martinez, José Luis¹

¹ DTU Bioengineering, Denmark

Synthetic biology has emerged as a revolutionary field with vast potential to address global challenges ranging from healthcare to environmental sustainability. Central to this endeavor is the development of efficient fermentation platforms capable of producing a variety of bio-based products. We herein present insights into the advancements achieved through a state-of-the-art fermentation platform facility dedicated to synthetic biology research and upstream bioprocess design.

The facility serves as a nexus for interdisciplinary collaboration, bringing together experts from biology, chemistry, engineering, and computational sciences. Through meticulous design and optimization, the facility enables custom design and optimization over a wide array of microbial fermentation processes, facilitating the production of complex biomolecules with tailored functionalities.

Key highlights include the utilization of novel genetic constructs and metabolic engineering strategies to enhance product yields and diversify output. Moreover, the integration of high-throughput screening techniques expedites the identification and optimization of microbial strains for specific applications.

Case studies illustrate our facility's versatility, ranging from the sustainable production of microbial based biofuels and pharmaceuticals to the synthesis of biodegradable polymers and nutraceuticals. Overall, this work underscores the pivotal role of the DTU fermentation core facility in advancing synthetic biology, paving the way for developing innovative solutions to address societal and environmental challenges.

Identification and characterization by a polyphasic approach of the microbial agent responsible of spoilage in Grana Padanocheese

Zago, Miriam¹; Rossetti, Lia¹; Bonvini, Barbara¹; Tidona, Flavio¹; Giraffa, Giorgio¹

¹ CREA Research Centre for Animal Production and Aquaculture, Italy

A peculiar alteration of microbial origin in Grana Padano (PDO) cheese was investigated. The defect appeared after 6–7 months of ripening as whitish spots located in the innermost parts of the cheese wheel and evolved with a degradation of the cheese matrix, followed by a formation of spongy cavities. Portions from both spoiled (S) and non-spoiled (NS) cheese areas were sampled and analyzed. Butyric clostridia spores (MPN g⁻¹) accounted 2.5 log in S areas compared to <-0.5 in NS ones. Metataxonomic analysis showed *Clostridium spp.* as the most abundant taxon in all the samples, with two distinct *Clostridium spp.* taxa dominating in NS and S areas, respectively. From the positive MPN tubes of both S and NS samples, 24 strains were isolated, identified as *Clostridium (C.) sporogenes* by species-specific PCR, and grouped into four distinct genotypes after RAPD-PCR fingerprinting analysis. One of the four genotypes included only isolates from NS-areas whereas the other three included isolates from the S-areas. 16S RNA sequencing of the dominant genotype of the S areas showed a 5 bp sequence polymorphism within the first 500 bp compared to the sole genotype of the NS areas. High-Resolution Melting applied to two strains representing these two genotypes showed two distinct profiles, which further suggests a taxonomic heterogeneity between them. Taken together, results confirmed the involvement of *C. sporogenes* in the onset of the defect of the cheeses studied. Genotypic and phenotypic characterization suggested the presence of two different prevailing biotypes of *C. sporogenes*, corresponding to two separate genetic lines, with notable morphological and taxonomic differences, isolated respectively from spoiled and defect-free zones of the cheeses. Further chemotaxonomic analysis is underway to clarify the taxonomic affiliation of the two distinct *Cl. sporogenes* genotypes isolated in this study.

The importance of the *Legionella* surveillance approach implemented in the company's water distribution system to discover a novel *Legionella* species: *Legionella petroniana* sp. nov.

Girolamini, Luna¹; Pascale, Maria Rosaria¹; Caligaris, Laura¹; Derelitto, Carlo¹; Marino, Federica¹; Spiteri, Simona¹; Cristino, Sandra¹

¹ Department of Biological, Geological, and Environmental Sciences, University of Bologna, Bologna, Italy

The study presents the description and characterization of a novel *Legionella* species isolated in Italy during a *Legionella* surveillance program developed in a company, where a risk assessment plan has been implemented since 2015. Sampling was conducted on cold and hot water distribution systems (WDS), following ISO19458:2006, and *Legionella* isolation was performed according to ISO11731:2017. *Legionella* isolates were identified using serological test and MALDI-TOF MS technique. Moreover, *mip* and *rpoB* gene sequencing, other than whole genome sequencing (WGS), were performed for *Legionella* species identification. In 2018, a *Legionella* contamination at a concentration of 400 +CFU/L was detected in a single cold point of WDS, located on the terrace of the building, used for air handling unit (AHU) cleaning and maintenance procedures. The colonies showed differences in terms of morphology with respect to the common *Legionella* features, although growth was only on BCYE with L-cysteine(Cys+). The isolates characterization process provided the following results: cells were Gram and Ziehl Neelsen-stain-negative, rod-shaped, and motile. The strains grew in a range of 32–37 °C on Cys+, GVPC, and MWY agar medium, with a positive reaction for oxidase and gelatinase. The dominant fatty acids were summed features 3 (C16:17c/C16:16c) (28.9%), C16:0 iso (18.4%), and C15:0 anteiso (15.4%), and Q13 as the major ubiquinone. The *mip* and *rpoB* gene sequences showed a similarity of 98.2% and 95.1%, with *L. feeleii* (ATCC 35072T). The GC content was 41.37 mol%, and the digital DNA-DNA hybridization (dDDH) analysis returned a value of 54.5% DNA-DNA relatedness with *L. feeleii* (NCTC 11978T). The Average Nucleotide Identity (ANI) between the strains and *L. feeleii* (NCTC 11978 T) was 94.02%, confirming that the isolates represent a novel species of the genus *Legionella*. The name proposed for this species is *Legionella petroniana* sp. nov., with 31f133T (=DSM 114357T=CCUG 76442T) as the type strain.

Session 8 – Networking, services, quality and data management of Microbial culture collections

Friday 20th September, 11.45:00am

Chairs: Lima Nelson (Portugal), Verkley Gerard (Netherlands).

MIRRI-ERIC as a microbial, genetics and data resources hub to advance scientific discovery

Melo Portugal, Ana¹

¹ *Executive Director at MIRRI-ERIC*

The Microbial Resource Research Infrastructure (MIRRI-ERIC), recognized as an ESFRI Landmark, serves as a linchpin for advancing bioscience and bioindustry endeavors. It offers a centralized platform for accessing diverse, high-quality bioresources and associated data, fully compliant with European-level data policies. Supported by a global network of microbial research centers and experts, MIRRI-ERIC fosters collaboration aimed at safeguarding microbial biodiversity, driving scientific inquiry, and nurturing professional development within the field. Presently, MIRRI-ERIC has six members (Belgium, France, Greece, Latvia, Portugal, and Spain) and one observer (Romania). Several other countries have shown interest in joining MIRRI-ERIC and contributing to strengthening its role as a hub for microbial, genetic, and data resources to advance scientific discovery. MIRRI-ERIC coordinates the Microbes4Climate Horizon Europe (HE) project, with the participation of two other research infrastructures (RIs), AnaEE-ERIC and LifeWatch-ERIC, along with twenty-eight research and innovation organizations. This initiative aims to provide researchers worldwide, including those focused on biodiversity, with streamlined access to a suite of integrated services, in harmony with the European Green Deal and the Digital Single Market Strategy. Additionally, MIRRI-ERIC, together with EMBRC-ERIC, ELIXIR, INSTRUCT-ERIC, four companies, and sixteen research and innovation organizations, was recently awarded the MALDIBANK project. This project aims to build a global cloud-based multi-domain open MALDI spectra bank for the identification of microorganisms. MIRRI-ERIC is reshaping its services and is deeply committed to making its culture collections FAIR—findable, accessible, interoperable, and reusable. These efforts align with the European Union's Digital Agenda and the European Strategy for Data, fostering innovation and knowledge exchange while safeguarding biodiversity for future generations.

Acknowledgements

The author is greatly indebted to MIRRI-ERIC and MIRRI-Nodes People, and to Microbes4Climate and MALDIBANK teams. Microbes-4-Climate – Project No. 101131818 is funded by the European Union's Horizon Europe program. MALDIBANK will be funded by the European Union's Horizon Europe program.

How the MIRRI-PT has been developed in its smart specialisation and cutting-edge technologies to offer better services

Lima, Nelson¹

¹ *MUM Micoteca da Universidade do Minho (MUM), Centre of Biological Engineering, University of Minho, and LABBELS Associate Laboratory, Braga/Guimarães, Portugal*

MIRRI-PT brings together 11 microbial culture collections from different regions. MIRRI-PT is well aligned with the domains of the “Regional Smart Specialisation Strategies”, namely: i) “Agroenvironmental Systems and Food”; ii) “Life Sciences and Health; iii) “Sea Resources and Economy” and iv) “Human Capital and Specialized Services”. In domain (1) we find the most relevant goods produced and exported include fermented alcoholic beverages, dairy products, and other agricultural products. MIRRI-PT has among its members that having a proven track record of collaboration with startups, SMEs, large companies, and regional farmer organisations in the referred sectors. In domain (2) we find a considerable amount of consolidated companies in medical devices, which have been mostly active in segments with a not very high incorporation of knowledge/technology, along with the youngest, smaller companies that target high-tech, high added-value segments. MIRRI-PT intends to strengthen its research capabilities and broaden the reach of services it provides. In the domain (3) MIRRI-PT is engaged in biotechnological activities associated with these microbial resources that will be the source of new products with applications in industry: antifouling compounds, biomaterials with application in food and health, and colourants for food and textile industries. Microbial resources may also be important for the development of microalgae and cyanobacteria-based biodiesels and biofuel as well as for the development of feed used in aquaculture. In domain (4) the approach has been the development and launch of an online platform featuring an e-catalogue of microbial resources, associated data and specialised services, to boost and facilitate access to the different user communities. This is an example of the efforts to promote digital transformation and the dematerialisation of processes. Finally, acquiring new equipment allows MIRRI-PT to offer new cutting-edge technologies and services.

Acknowledgements

This study partially supported by the Portuguese Foundation for Science and Technology (FCT) under the scope of the strategic funding of UIDB/04469/2020 unit and LABBELS – Associate Laboratory in Biotechnology, Bioengineering and Microelectromechanical Systems (LA/P/0029/2020); and by “MIRRI-PT (Pólo Norte)” project (PINFRA04/84445/2020) funded by the European Regional Development Fund under Norte2020-Programa Operacional Regional do Norte

PLAVIT, the Italian Plant Virus Collection in 2024

Forgia, Marco¹; Prigigallo, Maria Isabella²; Marra, Monica¹; Mussano, Paolo¹; Ciuffo, Marina¹; Vaira, Anna Maria¹; Vallino, Marta¹; Miozzi, Laura¹; Rubino, Luisa²; Accotto, Gian Paolo¹

¹ *Institute for Sustainable Plant Protection of National Research Council of Italy (CNR-IPSP), Torino, Italy;* ² *Institute for Sustainable Plant Protection of National Research Council of Italy (CNR-IPSP), Bari, Italy*

Viruses still generate major concern in public opinion, because mainly considered as human pathogens. However, they are more and more recognized as essential within ecosystems to maintain an ecological balance among different microorganisms and living beings in general. Therefore, viruses are key to environmental and human health, to science as well as to bioeconomy and biotechnology applications. Hence the need to develop and improve collections and biobanks not only of human viruses, but also of plant, microbial, and environmental viruses.

Plant virology started in Italy in the late 60's when specific expertise and equipment became available and new diseases of unknown etiology were studied. At that time virus identification was limited to bioassays on test plants, electron microscopy and basic serological tests. Dried leaf samples were collected and stored, mainly in laboratories of the National Research Council of Italy (CNR) located in Torino and Bari. The Plant Virus Italy (PLAVIT) collection, which includes more than 1,000 isolates, began to be redeveloped and quality-checked starting from 2020 thanks to dedicated European and national funds. The process, consisting of several steps (revitalization, authentication, sequencing, preservation, storage, distribution) is implemented in the frame of a Quality Management System and of international guidelines specifically developed for viruses that define mandatory information necessary for an isolate to be included into the PLAVIT catalogue. Up to now approximately half of the virus and viroid isolates present in the collection underwent the quality check and can be distributed upon request to academic and private research centers. In the framework of a continuous improvement scenario, more isolates will be processed, new ones will be acquired, and the categories in the catalogue will be expanded to include mycoviruses, bacteriophages and phytoplasmas, as well as nucleic acids and detection tools.

Acknowledgements

We wish to thank: the EVA-Global project, that received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No. 871029; the SUSMIRRI.IT project, granted by the European Commission—NextGenerationEU, code n. IR0000005; the BioMemory project, funded by the National Research Council of Italy.

The italian network of microbial culture collections: an overview on management and sustainability of the future research infrastructure MIRRI-IT

Moretti, Marino¹; Borlatto, Edoardo²; Varese, Giovanna Cristina¹

¹ *Department of Life Sciences and Systems Biology, University of Turin, Turin, Italy;* ² *Department of Management "Valter Cantino", University of Turin, Turin, Italy*

Majority of Italian microbial Culture Collections (mCCs) belong to the Joint Research Unit (JRU) MIRRI-IT, a network aiming at improving quality management system and addressing needs of stakeholders operating in the biotechnology transfer. Several institutions are currently undergoing an intensive “restyling” due to the NextGenerationEU-funded project SUS-MIRRI.IT, which focuses on the implementation of mCC management and offer of services. The creation of the National Node of a Research Infrastructure (RI) that will coordinate the network of Italian mCCs, under the umbrella of MIRRI-ERIC, the Microbial Resource Research Infrastructure-ERIC, is recognized as valuable upgrade to get recognition and accountability at international level, to garner support from stakeholders, to attract funding, and to foster continued and fruitful collaborations. However, beside these advantages, memberships in a RI requires compliance with several obligations and adoption of rules. The big challenges to overcome are the implementation of the quality management system according to International Standards; the compliance with legal frameworks; the digital information and technology access guided by the FAIR (Findable, Accessible, Interoperable, and Reusable) principle; the continuous amended capacity building to stay at the forefront of science and maintain stakeholder expectations, the top-level training of the personnel, the collaboration with policy-makers and industry; and, last but not least, the guaranteeing of a future financial sustainability. The situation of the Italian mCCs will be presented proposing a possible governance of the Italian RI MIRRI-IT after the end of the SUS-MIRRI.IT project. Moreover, a focus on the environmental, scientific, and socio-economic impact of this RI will be highlighted presenting stakeholder engagement, value proposition, collaborations with private sector, partnerships with national and local authorities, and funding strategy.

References

- [1] Moretti M. et al. (2024) Treasures of Italian Microbial Culture Collections: An Overview of Preserved Biological Resources, Offered Services and Know-How, and Management. *Sustainability* 2024, 16(9), 3777.

Acknowledgements

This work was granted by the European Commission – NextGenerationEU, Project SUS-MIRRI.IT "Strengthening the MIRRI Italian Research Infrastructure for Sustainable Bioscience and Bioeconomy", code n. IR0000005.

Managing change within a culture collection: A case study on the challenges within quality, data management and infrastructure within CABI's collection

Kermode, Anthony¹; Cobb, Emma¹; Buddie, Alan¹; Ryan, Matthew¹; Ananda, Amrutha¹; Adamin, Tomasz¹

¹ CABI, Egham, UK

Established in 1947, the CABI Genetic Resource Collection (GRC) is one of the most significant global collections of microorganisms, containing >28,000 fungi and approximately 2,000 bacteria. The collection is maintained through a combination of core funding from 48 member countries, commercial services and research projects. Until recently, commercial services were the focus, with some microbial services operating with ISO17025:2017 accreditation. A recent science-focused CABI review has resulted in the GRC's positive move to better supporting CABI's mission, member countries and project research activities, with an associated managed reduction in third-party audited commercial services. Quality is being maintained by refocussing resources on our other regulatory responsibilities such as plant health licensing, biological safety and the Nagoya protocol/CBD, contributing to recent works by the FAO on DSI regulation. Additional resources have been allocated to biological maintenance activities associated with supply of cultures and data management related to managing workflows within the collection. We have also been exploring ISO20387:2018 as a more relevant quality standard and to this end we maintain a presence on the steering group for the Biobank standard in the United Kingdom Accreditation Service (UKAS). Any of our remaining commercial activities either have a direct benefit to GRC operations or are performed with a consultation style approach that fits better with our mission and our expanding research portfolio. CABI's UK science centre is relocating to Imperial College's Silwood Park campus in 2025 with associated infrastructure upgrades for the GRC. In preparation, a programme of digitising all historic records relating to our strains has begun. Whilst the diversity of the archives provided challenges, it has positively impacted our ability to examine the provenance of strains when performing assessments on Nagoya compliance.

References

- [1] Smith D, Kermode A, Cafà G, Buddie AG, Caine TS & Ryan MJ (2020) Strengthening mycology research through coordinated access to microbial culture collection strains. *CABI Agriculture and Bioscience* 1: 2 <https://doi.org/10.1186/s43170-020-00004-9>

Acknowledgements

CABI is an international intergovernmental organisation, and we gratefully acknowledge the core financial support from our member countries and lead agencies. The EU (for EU Microbe and EU Microbes4Climate) and BBSRC-UKRI (for the UK Crop Microbiome Cryobank) are acknowledged for funding.

Establishment of the ITEM Culture Collection backup and development of the related disaster recovery plan

De Simone, Nicola¹; Ferrara, Massimo²; De Vero, Luciana²; Gialluisi, Katia²; Anelli, Pamela²; Petruzzino, Giuseppe²; Cefola, Maria¹; Palumbo, Michela¹; Cilenti, Lucrezia¹; de Chiara, Maria Lucia¹; Pelosi, Sergio¹; Capozzi, Vittorio¹; Moretti, Antonio²; Perrone, Giancarlo²

¹ *Institute of Sciences of Food Production of National Research Council (ISPA-CNR), Foggia, Italy;* ² *Institute of Sciences of Food Production, National Research Council (ISPA-CNR), Bari, Italy*

The development of microbial and microbiome-based solutions is decisive in delivering progress and sustainable innovations in agro-food systems. In this contest, Culture Collections (CCs) have a key role in ensuring scientific progress, promoting bioeconomy. The ITEM Microbial Culture Collection of ISPA (Institute of Sciences of Food Production, CNR, Bari) includes more than 13,000 strains belonging to various agri-food microorganisms (filamentous fungi, yeasts, and bacteria). In the framework of the European-funded SUS-MIRRI.IT project, which aims to improve the management and sustainability of Italian microbial collections, ITEM collection is implementing a disaster recovery plan to prevent accidental loss that includes an additional storage as “safe backup” in Foggia, about 130 kilometers from the primary territorial unit (ISPA-CNR, Bari). This safe storage facility includes a dedicated room with new devices for cryopreservation at -150°C and -80°C. As part of the project activities, a new molecular biology laboratory has been set up, including a capillary sequencer of nucleic acids, a spectrophotometer for nucleic acid and protein measurements, and an imaging system for molecular biology. A laboratory for preparatory procedures completes this infrastructural effort. All these facilities are also functional to restore the correct operation of ITEM collection in the occurrence of an emergency or unexpected negative event in the headquarters. Strategies are being implemented for the preservation of digital information (microbiological passport data). Tailored training activities are ongoing to ensure the appropriate recovery, transfer and cryopreservation of the microorganisms in the safe storage facility. Furthermore, we highlight how the different research lines active in the secondary territorial unit of Foggia will represent the driver of innovation to increment and exploit the micro-biodiversity preserved in the ITEM culture collection.

Acknowledgements

Project SUS-MIRRI.IT “Strengthening the MIRRI Italian Research Infrastructure for Sustainable Bioscience and Bioeconomy” Area ESFRI “Health and Food”, granted by the European Commission – NextGenerationEU Code N° IR0000005.

A cloud-based, open-access platform for comprehensive metagenomic data analysis

Contaldo, Sandro Gepiro¹; Tortarolo, Dora¹; Venice, Francesco²; Pantini, Francesco¹; Bosio, Lorenzo¹; Ferrocino, Ilario³; Cordero, Francesca¹; Varese, Giovanna Cristina²; d'Acierno, Antonio⁴; Beccuti, Marco¹

¹ Department of Computer Science, University of Torino, Turin, Italy; ² Department of Life Sciences and Systems Biology, Mycotheca Universitatis Taurinensis, University of Torino, Turin, Italy; ³ Department of Agricultural, Forestry and Food Sciences, University of Torino, Grugliasco, Italy; ⁴ Institute of Food Sciences, CNR-ISA, Avellino, Italy

The Italian Node of the Microbial Resource Research Infrastructure, overseen by the Joint Research Unit MIRRI-IT, is essential for advancing microbial research in Italy. In 2022, the Italian government allocated €17 million from the NextGeneration EU-funded PNRR to support the SUS-MIRRI.IT project. Since microbial resources include extensive metadata and encompass high-throughput sequencing datasets, within the SUS-MIRRI project, we have worked to create metagenomic analysis workflows and to implement them as an online service orchestrating various software within an HPC architecture. To ensure accessibility, we implemented a web user interface using Next.js. The workflow implementation utilises StreamFlow [1], a Workflow Management System based on the Common Workflow Language standard, to be executed within the HPC system using SLURM [2], to allow multiple workflows to run simultaneously.

To show the effectiveness of our approach, we present here two workflows. The first one is designed for microbial genome assembly: it supports long-read assembly by harnessing various assembler tools [3,4,5], then polishes and merges the results to improve the outcome. The second one performs metagenomic analysis, including taxonomic annotation and microbial population characterisation through: i) reading alignment for host removal and taxonomic classification using Kraken2 [6] and MetaPhlAn4 [7], ii) post-processing for normalisation, decontamination, filtering, abundance estimation, and diversity calculation, and iii) metadata analysis and regression model computation. These two workflows demonstrate our service's scalability by varying the number of computational resources.

Our platform empowers scientists with remarkable computing power, unavailable to most laboratories, and supports their research endeavours with an interactive and clever environment. By leveraging interdisciplinary knowledge, our platform enables the exploration of results from multiple perspectives.

References

- [1] Colonnelli et al, DOI: 10.1109/TETC.2020.3019202 (2020)
- [2] Georgiou et al, DOI: 10.1007/978-3-642-35867-8_8 (2012)
- [3] Koren et al, DOI: 10.1101/gr.215087.116 (2017)
- [4] Kolmogorov et al, DOI: 10.1038/s41587-019-0072-8 (2019)
- [5] Ruan et al, DOI: 10.1038/s41592-019-0669-3 (2019)
- [6] Wood et al, DOI: 10.1186/s13059-019-1891-0 (2019)
- [7] Blanco-Míguez et al, DOI: 10.1038/s41587-023-01688-w (2023).

Acknowledgements

This work was supported by the European Commission NextGenerationEU, Project "Strengthening the MIRRI Italian Research Infrastructure for Sustainable Bioscience and Bioeconomy", code n. IRO000005 (within the Italy's National Recovery and Resilience Plan).

Strengthening the CoSMi culture collection: a contribution to preserving the biodiversity of microalgae

Cerino, Federica¹; Di Poi, Elena¹; Natali, Vanessa¹; Bordiga, Manuela¹; Beran, Alfred¹

¹ *Istituto Nazionale di Oceanografia e di Geofisica Sperimentale (OGS), Sgonico (TS), Italy*

Microalgae are unicellular autotrophic organisms that produce half of the oxygen we breathe and remove carbon dioxide from the atmosphere through photosynthesis. They are the main primary producers of biomass for aquatic systems, and thus support life on earth. In addition to this key ecological role, their potential importance for biotechnology has emerged in last decades. Microalgae are indeed a diverse group, containing a large amount of proteins, lipids, carbohydrates, pigments, -3 fatty acids and other chemicals that could serve as feedstock for various products in different fields such as pharmaceuticals, cosmetics, food and feed, but also as an alternative source of renewable fuels. Against this background, culture collections play a key role in protecting and preserving this invaluable microalgae biodiversity. However, these collections are often the result of the research needs and efforts of individual researchers, so they do not meet the relevant international standards.

The Collection of Sea Microorganisms (CoSMi) at the National Institute of Oceanography and Applied Geophysics (OGS) is one of the few official collections specialised in marine microalgae. It is part of the WFCC and is linked to national and international research infrastructures such as MIRRI.IT, LifeWatch, ECCSEL and EMBRC ERIC, as well as being an integral part of the BioMarine Lab and the Gulf of Trieste Observatory System. In the framework of the Italian PNRR project SUS-MIRRI.IT, promoted by JRU MIRRI.IT, the CoSMi collection will be strengthened by: 1) improving technological equipment and scientific instruments; 2) better characterization of microbial biodiversity through morphological and molecular analyses; 3) developing and implementing standard procedures for sampling, characterization and conservation; 4) improving online accessibility of information by implementing and publishing the local database and harmonising it with the national database and by upgrading the website.

The SUS-MIRRI.IT project: research, services, and training for sustainable bioscience and bioeconomy by the MIRRI Italian Research Infrastructure

Arena, Mattia Pia¹ ON BEHALF OF THE SUS-MIRRI.IT PARTNERS

¹ *Department of Life Sciences, University of Modena and Reggio Emilia, Reggio Emilia, Italy*

SUS-MIRRI.IT is a project funded by the NextGenerationEU programme, with a total budget of about € 17,000,000. The project involves 24 Italian Operative Units.

The mission of SUS-MIRRI.IT is to strengthen the Italian network of Culture Collections (CCs) holding different microbial resources, including bacteria, filamentous fungi, yeasts, microalgae and viruses. The allocation of funds is aimed at the acquisition of cutting-edge equipment, infrastructural improvements, recruitment of high-qualified staff, strengthening of the preserved microbial resources with associated data, and top-level services. The project is organized into 6 Work Packages (WPs): WP1 drives management, governance, and sustainability; WP2 is involved in microbial CCs consolidation; WP3 focuses on digital activities for the creation of the single-entry point to the National Database of microbial resources and services; WP4 aims to study the biotechnological valorisation of microbiomes; WP5 deals with the implementation of offered services and training courses; WP6 is dedicated to communication and dissemination actions.

To date, over 40 researchers, PhD students, and technicians have been hired; about 70% of the funds for innovative technologies have been utilized; the online platform including the catalogue of Italian microbial resources has been launched; 19 advanced training courses have been delivered, with a participation rate of over 900 attendees; 30 scientific papers have been published. All project updates are communicated by the dedicated website and related social media.

The strategic impact of the project is to promote partnerships on the territory and encouraging the aggregation of skills, structures, and bioresources. Targeted initiatives and stakeholder-oriented actions will ensure that results are disseminated and, when possible, transferred to industry, promoting the transition towards the green economy, and contributing to the sustainability and safety of the environment.

The Usage and Fate of Patent Deposits

Bajerski, Felizitas¹; Felsch, Kathrin¹

¹ *Leibniz Institute DSMZ-German Collection of Microorganisms and Cell Cultures GmbH, Germany*

Since 1981, the DSMZ has been recognised as an International Depositary Authority (IDA) under the Budapest Treaty. The tasks of an IDA are clearly defined: Deposit of samples for patent purposes and provision of biological material to authorised parties. The DSMZ holds more than 9,000 patent deposits, that are mentioned in more than 24,000 patents (Scopus, using "DSMZ" as a search parameter). However, many patent deposits never make it into a patent, either because the deposited material does no longer fall within the scope of the depositor or the Budapest Treaty deposit is "misused" for other purposes. Many patent deposits are also mentioned in scientific publications, although this form of deposit may not be in line with the need for public availability, especially when type strains are involved. Interestingly, DSMZ patent deposits also appear in EFSA publications. This is presumably because EFSA requires that "organisms to be evaluated should be deposited in an internationally recognised culture collection that has acquired the status of an international depository under the Budapest Treaty" [1], which refers to the organisation of the culture collection but not the form of deposit. In this case, a safe deposit would be recommended. Considering that patent deposits can be used for other purposes than intended, we would like to point out that it is not clear what should happen to these deposits after the mandatory storage period of at least 30 years after the date of deposit (Rule 9.1 Budapest Treaty), with options ranging from destruction to open availability of the material. We would therefore like to continue the discussion on what should happen to these deposits after the described storage period, considering various aspects such as the disclosure of patents, the open availability of the material and the feasibility for IDA collections. As well as asking whether other defined forms of deposit are needed to fulfil stakeholder requirements.

References

- [1] Additives, E. Panel o., Feed, P. o. S. u. i. A., Rychen, G., Aquilina, G., Azimonti, G., Bampidis, V., Bastos, M. d. L., Bories, G., Chesson, A., Cocconcelli, P. S., Flachowsky, G., Gropp, J., Kolar, B., Kouba, M., López-Alonso, M., López Puente, S., Mantovani, A., Mayo, B., Ramos, F., Saarela, M., Villa, R. E., Wallace, R. J., Wester, P., Glandorf, B., Herman, L., Kärenlampi, S., Aguilera, J., Anguita, M., Brozzi, R., & Galobart, J. (2018). Guidance on the characterisation of microorganisms used as feed additives or as production organisms. *EFSA Journal*, 16(3), e05206. <https://doi.org/https://doi.org/10.2903/j.efsa.2018.5206>

Performing an upgraded and interoperable data management system for ITEM Microbial Culture Collection

Petruzzino, Giuseppe¹; De Vero, Luciana¹; Cozzi, Giuseppe¹; Gialluisi, Katia¹; Anelli, Pamela¹; Perrone, Giancarlo¹; Moretti, Antonio¹; Susca, Antonia¹

¹ *Institute of Sciences of Food Production, National Research Council, Bari, Italy*

ITEM is an internationally recognized and ISO 9001-certified culture collection, hosted at the CNR Institute of Sciences of Food Production (ISPA) in Bari (Italy). ITEM holds more than 14.000 strains of agro-food interest including filamentous fungi, yeasts and bacteria. It is a partner of the Italian network of microbial resources collections "JRU MIRRI-IT" (<https://mirri-it.it/>) and has been involved in the MIRRI-ERIC Infrastructure constitution (<https://mirri.org>). In addition to the ITEM database, strains-related data are included in the i) World Data Centre for Microorganisms (<https://wdcm.org/>), ii) MIRRI-ERIC Catalogue and iii) CNR Biomemory Project network (<https://biomemory.cnr.it/>).

Thanks to the support of the ongoing project "SUS-MIRRI.IT", ITEM is upgrading and improving both the collection database and infrastructure, to integrate all the information associated with the strains preserved, improve quality control management and setting up a disaster recovery plan by creating a physical backup at the CNR-ISPA institute in Foggia (Italy). Generally, collections databases should not be standalone systems, but they must interact with other bioinformatics systems to achieve a global view of data and passport information available on strains [1]. Accordingly, the ITEM upgraded database management relies on interconnections among different platforms storing biological data and microbial culture data: BioMICS (BioAware) for on-line catalogue, base statistics and images DB (<https://item.bio-aware.com/>); EasyTrack2D (Twin Helix) for internal physical storage reference; Bionumerics (Applied Maths) for DNA/RNA sequences and passport data. All information is locally managed by R scripts (R Core Team) to conform different standards.

The final goal is to standardize all information under FAIR principles (<https://go-fair.org/>), in a unique platform with Application Programming Interface capabilities to intercommunicate with other external components.

References

- [1] De Vero L, Boniotti MB, Budroni M, Buzzini P, Cassanelli S, Comunian R, Gullo M, Logrieco AF, Mannazzu I, Musumeci R, Perugini I, Perrone G, Pulvirenti A, Romano P, Turchetti B, Varese GC. Preservation, Characterization and Exploitation of Microbial Biodiversity: The Perspective of the Italian Network of Culture Collections. *Microorganisms*. 2019 Dec 12;7(12):685. doi: 10.3390/microorganisms7120685. PMID: 31842279; PMCID: PMC6956255.

Acknowledgements

"Strengthening the MIRRI Italian Research Infrastructure for Sustainable Bioscience and Bioeconomy" research project (acronym: SUS-MIRRI.IT; European Commission – NextGenerationEU code n. IR0000005

Bank of Algae and Cyanobacteria of the Azores – BACA culture collection

Luz, Rúben^{1,2}; Costa, Vanessa²; Cordeiro, Rita²; Fonseca, Amélia^{1,2}; Gonçalves, Vitor^{1,2}

¹ Universidade dos Açores, Faculdade de Ciências e Tecnologia, Ponta Delgada, Portugal; ² CIBIO, Centro de Investigação em Biodiversidade e Recursos Genéticos, InBIO Laboratório Associado, BIO-POLIS Program in Genomics

The Bank of Algae and Cyanobacteria of the Azores (BACA) culture collection was created in 2018 reuniting several microalgae and cyanobacteria culture collections from the University of the Azores (Portugal) in one unique collection, guaranteeing quality and long-term maintenance of these important genetic resources. Since then, BACA has reached 800 cultured strains of cyanobacteria and microalgae. Is a member of ECCO and Pt-mBRCN since 2021, promoting access to its cultures through a public catalog (baca.uac.pt). The BACA culture collection is a hub of biodiversity allowing for several studies on taxonomy, with several novel genera and species published, supporting the conservation of these unique genetic resources isolated from the Azores archipelago. Several works have been published regarding the toxicity of cyanobacteria but also taking into account biotechnological applications, valorizing the BACA culture collection as a source of important compounds. Recently, more than 100 genomes of cyanobacteria from the BACA culture collection have been sequenced by Illumina, creating high-quality draft genomes, which further support genetic and biotechnological studies.

Acknowledgements

Rúben Luz was supported by a Ph.D. grant (M3.1.a/F/002/2020) from the Fundo Regional da Ciência e Tecnologia (FRCT).

This work was funded by FEDER funds through the Interreg-MAC 2014-2020 Program under the projects REBECA—Red de excelencia en biotecnología azul (algas) de la región macaronesia (MAC1.1a/060) and REBECA-CCT—Red de Excelencia en Biotecnología Azul de la Región Macaronésica. Consolidación, Certificación y Transferencia (MAC2/1.1b/269).

By Portuguese National Funds, through FCT—Fundação para a Ciência e a Tecnologia, the European Union, QREN, FEDER, COMPETE, by funding the CIBIO/InBIO (project UID/BIA/ 50027/2013 and POCI-01-0145-FEDER-006821).

A ranking tool for microbial biobanks

Boutroux, Martin¹; Chiarelli, Adriana¹; Ferrari, Mariana L.¹; Chesneau, Olivier²; Clermont, Dominique²; Betsou, Fay¹

¹ Institut Pasteur, Université Paris Cité, Biological Resource Center of Institut Pasteur – Project Management Office, Paris, France; ² Institut Pasteur, Université Paris Cité, Biological Resource Center of Institut Pasteur – Collection de l'Institut Pasteur, Paris, France

Microbial biobanks preserve and provide microbial bioresources for research, training, and quality control purposes. They ensure the conservation of biodiversity, contribute to taxonomical research, and support scientific advancements. Microbial biobanks can cover a wide range of phylogenetic and metabolic diversity, or focus on specific taxonomic, thematic, or disease areas. The strategic decisions about strain selection for certain applications or for the biobank culling necessitate a method to support prioritization and selection.

Here, we propose an unbiased scoring approach based on objective parameters to assess, categorize, and assign priorities among samples in stock in a microbial biobank. We describe the concept of this ranking tool and its application to identify high priority strains for whole genome sequencing with two main goals: (i) genomic characterization of quality control, reference, and type strains;

(ii) genome mining for the discovery of natural products, bioactive and antimicrobial molecules, with focus on human diseases. The general concept of the tool can be useful to any biobank and for any ranking or culling needs.

Acknowledgements

The work was financed by H2020-INFRADEV-03-2019 Project IS_MIRRI21 (Grant Agreement No. 871129).

A shared quality management system for Italian Culture Collections: The challenge of SUS-MIRRI.IT project

Perugini, Iolanda¹; De Vero, Luciana²; Turchetti, Benedetta³; Prigione, Valeria¹; Accotto, Gian Paolo⁴; Varese, Giovanna Cristina¹; Perrone, Giancarlo²

¹ Department of Life Sciences and Systems Biology, University of Turin, Turin, Italy; ² Institute of Sciences of Food Production, National Research Council, Bari, Italy; ³ Department of Agricultural, Food and Environmental Sciences, University of Perugia, Perugia, Italy; ⁴ Institute for Sustainable Plant Protection, National Research Council, Turin, Italy

Quality management system (QMS) implementation is one of the main goals of the European Microbial Resource Research Infrastructure (MIRRI-ERIC), which operates actively on the design of standard guidelines. Culture collections (CCs) must manage the quality of their bioresources and guarantee the deposited microbial strains' viability, purity, and identity according to internal standard operation procedures (SOPs). In this respect, the SUS-MIRRI.IT project aims to harmonize and improve the QMS of the Italian network of CCs. Within the activities of the project, three working groups dealing with (i) Common Collections Procedures, (ii) Compliance with the Nagoya protocol, and (iii) Code of Conduct on Biosafety and Biosecurity, are working to define a general and harmonized management system for the Italian CCs. The description of the common procedures is still in progress with the definition of SOPs for accepting microbial cultures, their quality check, and protocols for microorganisms' preservation and distribution. Regarding the Legal compliance with the Nagoya protocol, starting from the Material Transfer Agreement (MTA) and Material Deposit Agreement (MDA) models recently published [1], a common document implementing MTA and MDA for the Italian CCs has been developed and proposed with the support of an external expert in law. The QMS should also address the capability of collections to meet all relevant national and international rules concerning the safe distribution of different bioresources. Accordingly, the working group on biosafety and biosecurity, which includes experts on human, animal, and plant diseases, is implementing guidelines for handling microorganisms based on their risks. The adoption of common rules for quality management is a challenging goal of primary importance for the CCs involved in SUS-MIRRI.IT to build a valid national network capable of responding to the needs of public and private actors in the fields of research and bioeconomy.

References

[1] Verkley, G. et al. FEMS Microbiology Letters, 367, 2020 (<https://doi.org/10.1093/femsle/fnaa044>)

Acknowledgments

"Strengthening the MIRRI Italian Research Infrastructure for Sustainable Bioscience and Bioeconomy" research project (acronym: SUSMIRRI.IT; European Commission – NextGenerationEU code n. IRO000005)

The ENEA Microbial Culture Collection as valuable resource for developing new sustainable biotechnologies and bioeconomy

Bevivino, Annamaria¹; Ambrico, Alfredo¹; Molino, Antonio¹; De Rosa, Luigi¹; Colantoni, Eleonora¹; Casella, Patrizia¹; Tabacchioni, Silvia¹; Alisi, Chiara¹; Costanzo, Manuela¹; Lico, Chiara¹; Bianco, Linda²; Cuna, Daniela¹ and the ENEA's team of SUS-MIRRI.IT project

¹ Department for Sustainability, Italian National Agency for New Technologies, Energy and Sustainable Economic Development, Roma, Italy; ² Department of Energy Technologies and Renewable Sources, Italian National Agency for New Technologies, Energy and Sustainable Economic Development, Roma, Italy

The ENEA microbial culture collection (EMCC) has been created over the course of 30 years by an interdepartmental team of researchers that have collected microorganisms and microbial consortia from different environments, like contaminated sites, hypogea and archaeological sites, food, lake sediments, sea, soil, rhizosphere, water [1]. The microbial collection has several important applications ranging from the health and quality of soil and cultivated plants to the degradation of environmental contaminants, from the production of bio-based molecules for industrial, energy and food uses to new products for the restoration of artistic heritage, from the development of functionalized packaging to the support of food quality and safety [2]. EMCC's mission is to isolate, identify, characterise, collect, store and distribute the microbial resources and related information, to improve their characterization and optimise their management, thus unlocking their genomic and metabolic potential. EMCC collection actually spans over 1400 microbial strains including mainly bacteria, but also fungi, yeasts, two microalgae and a plant virus. The most abundant genera are *Burkholderia*, *Pseudomonas*, *Bacillus*, *Arthrobacter*, *Streptomyces*, *Paenibacillus*, *Staphylococcus*, *Stenotrophomonas*, and *Microbacterium* for bacteria, *Aspergillus*, *Penicillium*, *Engyodontium*, *Cladosporium*, and *Trichoderma* for fungi, and *Saccharomyces*, *Rhodotorula* for yeasts. All the information i.e., the taxonomic assignment, origin, geographical geolocation, growth conditions and fields of application, is stored in a relational database, accessible through a web application and easily consultable using search keys that allow you to get the desired data. The preparation of bacterial starter in 2-5 L bioreactors and the scaling-up of the process in 15, 50 and 500-liter represent one of the important services that the EMCC will provide over the years to third parties in the agricultural and novel food sectors.

References

- [1] Bevivino A. La collezione microbica ENEA e l'infrastruttura MIRRI, Microbial Resources Research Infrastructure. EAI, 50/2020
- [2] Moretti et al. Treasures of Italian Microbial Culture Collections: An Overview of Preserved Biological Resources, Offered Services and Know-How, and Management. Sustainability 2024, 16, 3777. <https://doi.org/10.3390/su16093777>

Acknowledgements

This work was granted by the European Commission NextGenerationEU within the framework of the National Recovery and Resilience Plan (PNRR) (Mission 4 "Education and Research", Component 2, Investment 3.1), Project "Strengthening the MIRRI Italian Research Infrastructure for Sustainable Bioscience and Bioeconomy" (SUS-MIRRI.IT), code IR0000005.

Speaking the same language: Changing the way to share microbial strain data

Whitte, Julius Fredrich¹; Reimer, Lorenz Christian¹; Yurkov, Andrey¹; Bunk, Boyke¹; López-Coronado, José Miguel²; Alokoy, Susan²; Zuzuarregui, Aurora²; Aznar, Rosa²; de Vries, Ronald P.³; Robert, Vincent²; Overmann, Jörg¹

¹ *Leibniz Institute DSMZ -German Collection of Microorganisms and Cell Cultures GmbH, Braunschweig, Germany;* ² *Spanish Type Culture Collection-University of Valencia (CECT-UVEG), Paterna, Spain;* ³ *Westerdijk Fungal Biodiversity Institute, Utrecht, The Netherlands*

Microbial strain data is often highly unstructured or the structure is tailored to only fit one system. To solve this problem we looked at the existing and historical standards for strain data and came up with a modular, flexible and technical up to date solution, integrating existing schemas. In addition to this linked data we will also develop a technical infrastructure providing online tools to support and use this format.

Acknowledgements

This work was created by WP6 from the EU-Project Bioindustry 4.0 Grant Agreement #101094287

Alphabetical Index of Authors

*Presenting Authors in **bold**

Abbamondi, Gennaro Roberto	Bevilacqua, Antonio	Carrasco-Acosta, Marina
Abouloifa, Houssam	Bevivino, Annamaria	Carriero, Francesco
Accettulli, Alessandra	Bianco, Angela	Caruso, Tiziano
Accotto, Gian Paolo	Bianco, Linda	Cascone, Martina
Adamin, Tomasz	Biasioli, Franco	Casella, Patrizia
Akyildiz, Gürçay Kivanç	Bindo, Arianna	Cataldi, Mauro
Alberoni, Daniele	Blancke, Gillian	Catellani, Lisa
Alessandro, Bernetti	Blondet, Eddy	Cattaneo, Andrea
Alex Bisson	Böhm M. Calore R.	Cecchi, Grazia
Aliano, Anita	Bolhuis, Henk	Ceci, Andrea
Alisi, Chiara	Bonnin, J. Miguel	Cefola, Maria
Allemand, Denis	Bonvini, Barbara	Cerino, Federica
Aloi, Francesco	Booth, Nicky Sebastian	Chaillou, Stéphane
Alokozay, Susan	Bordeau, Thierry	Chambellon, Émilie
Altieri, Clelia	Bordiga, Manuela	Chesneau, Olivier
Amaral Carneiro, Greice	Borges, Inês	Chessa, Luigi
Ambrico, Alfredo	Borlatto, Edoardo	Chiarelli, Adriana
Ananda, Amrutha	Boroduske, Anete	Chiellini, Carolina
Anderson J. ,	Borruso, Luigimaria	Cieniuch, Gabriela
Andreani, Daniele	Boscaino, Floriana	Ciers, Maarten
Andreolli, Marco	Bosio, Lorenzo	Cifarelli, Valentina
Andries, Vanessa,,	Botticella, Lucia	Cilenti, Lucrezia
Anelli, Pamela	Boullie, Anne	Ciro, Sannino
Aracri, Benedetto	Boutroux, Martin	Ciuffo, Marina
Arena, Mattia Pia	Bracco, Perangiola	Civolani, Eleonora
Asehraou, Abdeslam	Brasca, Milena	Clermont, Dominique
Ashley, Dias	Briandet, Romain	Cobb, Emma
Assandri, Davide	Bruder A.,	Colantoni, Eleonora
Astolfi, Maria Luisa	Bruno Baron, Camila	Combina, Mariana
Aurino, Elena	Bubici, Giovanni	Comunian, Roberta
Aznar, Rosa	Buddie, Alan	Contaldo, Sandro Gepiro
Bachy, Charles	Budroni, Marilena	Conversa, Lucia
Bagiyan, Valeri	Bunk, Boyke	Copetti V., Marina
Bajerski, Felizitas	Buzzini, Pietro	Corbo, Maria Rosaria
Balode, Laura	Cabassi, Giovanni	Cordeiro, Rita
Bang-Berthelsen, Claus Heiner	Cali, Martina	Cordero, Francesca
Barth, Dorothee	Caligaris, Laura	Cordone, Angelina
Baruzzi, Federico	Caliro, Stefano	Cornet, Luc
Baurain, Denis	Calpe, Jorge	Coronas, Roberta
Beccuti, Marco	Canonica, Laura	Corso, Davide
Bellino, Mariarosa	Canteiro C.	Corvino, Antonia
Beran, Alfred	Capozzi, Nunzio	Cosentino, Sofia
Bernini, Roberta	Capozzi, Vittorio	Cossu, Carla
Berri, Cécile	Cappelletti, Eleonora	Costa S.
Berx, Geert	Caprio, Alfonso	Costa, Vanessa
Betsou, Fay	Carella, Daria	Costantini, Antonella
Betta, Emanuela	Carimi, Francesco	Costanzo, Manuela

Alphabetical Index of Authors

*Presenting Authors in **bold**

- Cozzi, Giuseppe
Crespi, Matteo
Criscuolo, Alexis
Cristino, Sandra
Crucitti, Dalila
Cufino, Gaia
Cuna, Daniela
Cunedioğlu, Hülya
d'Acierno, Antonio
d'Amelio, Annalisa
D'Amodio, Antonia
Daga, Elisabetta
Dall'Asta, Chiara
Daniel, Heide-Marie
Daniela, Campaniello
Daniele, Andreani
De Simone, Nicola
De Bellis, Palmira
De Caro, Salvatore
De Chiara, Maria Lucia
De Natale, Antonino
De Noni, Ivano
De Rosa, Luigi
De Santis, Alessandro
De Santis, Letizia
De Santis, Michele
De Simone, Nicola
De Vero, Luciana
de Vries, Ronald P.
Declerck, Stéphane
Defazio, Giuseppe
Del Fiore, Antonella
del Solar, Gloria
Dell'Olmo, Eliana
Della Mura, Brigida
Denfeld B.,
Deplano, Maura
Derelitto, Carlo
Di Biase, Mariaelena
Di Cesare, Francesca
Di Donato, Paola
Di Gioia, Diana
Di Gregorio, Luciana
Di Piazza, Simone
Di Poi, Elena
Di Renzo, Tiziana
Diaferio, Antonietta
Diependaele, Louise
Dindo, Maria Luisa
Dodhia, Maya
Donati, Enrica
Duclos, Michel
Durmusoglu, Deniz
Eckert E.M.
Elad Stolovicki
Elewaut, Dirk
Epifani, Filomena
Escriva, Laura
Faino, Luigi
Fanelli, Francesca
Felis, Giovanna
Felsch, Kathrin
Fernandes I.
Ferrara, Massimo
Ferrari E.
Ferrari, Mariana L.
Ferrier-Pages, Christine
Ferrocino, Ilario
Filannino, Pasquale
Finore, Ilaria
Fiorentino, Sofia
Fiorin, Anita
Flagella, Zina
Florio Furno, Matteo
Fonseca, Amélia
Fontaneto D.
Forgia, Marco
Fosso, Bruno
Fragasso, Mariagiovanna
Frainer A.
Francati, Santolo
Gachenot, Martin
Gagliardi, Roberto
Gamian, Andrzej
Ganzert L.
Garcés E.
Garello, Marco
Garzoli, Laura
Gatto, Angela
Germain, Karine
Gialluisi, Katia
Giancaspro, Angelica
Giavalisco, Marilisa
Giorgetti, Lucia
Giovannelli, Donato
Giovannini, Roberto
Giraffa, Giorgio
Girolamini, Luna
Goginyan, Vigen
Gökşen, Gülden
Gonçalves, Vitor
Gourvil, Priscillia
Graça D.,
Grego, Michele
Grossart H.P.
Grzywa, Renata
Gudra, Dita
Guerrieri, Angela
Gugger, Muriel
Guglielmetti, Simone
Gullo, Maria
Gumiel, Marta
Haidukowski, Miriam
Harutyunyan, Seda
Helloin, Emmanuelle
Hernández-Alcántara, Annel
Hervéou, Katell
Hölttä, Matti
Hondelatte, Alice
Hryniewicz, Katarzyna
Ida Helene Steen
Ido Caspy
Inés Patop
Iori, Valentina
Iosca, Giovanna
Jaberi, Mehrdad
Jans, Maude
Jauga, Kristians
Joannic, Julie
Johansen, Tina
John Mallon
Joubert, Flavien
Kaczorowska, Anna-Karina
Kaczorowski, Tadeusz
Karganyan, Arthur
Kasenömm, Priit
Kermode, Anthony
Kęsy, Jacek
Khachatryan, Gayane
Khomenko, Iuliia
Kisand V.

Alphabetical Index of Authors

*Presenting Authors in **bold**

Koivuranta, Kari	Mazzonna, Marco	Parente, Eugenio
Kolata, Magdalena	Meca, Giuseppe	Pascale, Maria Rosaria
Korzeniowska-Kowal,	Melo Portugal, Ana	Pasquale, Marena
Agnieszka	Migliaccio, Flavia	Pedro Escudeiro
Kostic, Tanja	Millette K.	Pelosi, Sergio
Kostrzewa, Maja	Mimmo, Tanja	Perrone, Giancarlo
Koublová, Vendula	Miozzi, Laura	Perrotta, Gaetano
Laera, Angelica	Modesto, Monica	Persiani, Anna Maria
Lampis, Silvia	Molinaro, Marcella	Perugini, Iolanda
Lange, Lene	Molino, Antonio	Pesole, Graziano
Larocca, Vincenzo	Mon, Maria Laura	Petraretti, Mariagioia
Lasagni, Federico	Montemurro, Marco	Petrina, Zaiga
Latronico, Rosanna	Morandi, Stefano	Petruzzino, Giuseppe
Lavanga, Emanuela	Moreno, Ana	Petta, Ioanna
Lazzari, Gioele	Moretti, Antonio	Pezzella, Giovanni
Leone, Luigi	Mugnai, Gianmarco	Picciotti, Ugo
Licata, Marzia	Muhle, Estelle	Pinnow S.
Lico, Chiara	Mulè, Giuseppina	Pinzari, Flavia
Lima, Nelson	Munoz Fernández, Gloria	Pirra, Sara
Longo, Angela	Mussano, Paolo	Pisano, Maria Barbara
Lonigro, Stella Lisa	Muzzini, Valerio Giorgio	Poli, Anna
Lopez Ocaña, Laura	Nakurte, Ilva	Poli, Annarita
López-Coronado, José Miguel	Napoletano, Giovanna	Pollio, Antonino
Lotti, Concetta	Nappi, Nunzia	Porcar, Manuel
Luigimaria, Borruso	Natali, Vanessa	Posth, Nicole R.
Łukasz, Dzięwit	Navarro-Torre, Salvadora	Povolo, Milena
Lupo, Valérian	Nawaz, Asma	Presutto, Ester
Luz, Carlos	Nazzaro, Stefania	Prigigallo, Maria Isabella
Luz, Rúben	Němcová, Karolína	Prigione, Valeria
Magarelli, Rosaria Alessandra	Nicolaus, Barbara	Prins V.
Magistà, Donato	Nikolajeva, Vizma	Prioux, Camille
Mammola S.	Nolfi, Lorenzo	Probert, Ian
Mamone, Gianfranco	Nováková, Dana	Prodi, Antonio
Mändar, Reet	Novotný, David	Przemysław, Decewicz
Manikas, Ioannis	Nygren, Heli	Pulcini, Laura
Marangi, Matteo	Oliva, Feliciano	Quiles, Juan Manuel
Marchese, Pietro	Olivia Leland	Rabbi, Francesca
Marchitelli, Andrea Lara	Omri, Ghofrane	Racioppo, Angela
Marena, Pasquale	Overmann, Jörg	Rahi, Praveen
Marino, Federica	Paba, Antonio	Raio, Aida
Marra, Luigi	Pacifico, Davide	Rämä T.
Marra, Monica	Palazzo, Salvatore	Ravon, Laure
Marrero Díaz de Villegas, Rubén	Palojärvi, Ansa	Reale, Anna
Martino, Maria	Palumbo, Michela	Reginato, Alessia
Masiello, Mario	Panksep K.	Reimer, Lorenz Christian
Mattarelli, Paola	Pantini, Francesco	Remaut, Han
Mazzeo, Maria Fiorella	Papikyan, Razmik	Reñé A.

Alphabetical Index of Authors

*Presenting Authors in **bold**

Retter A.	Sinigaglia, Milena	Vale Dias, Teresa
Cruz, Ricardo M.	Słowakiewicz, Mirosław	Valerio, Francesca
Ricciardi, Annamaria	Soares, Célia	Vallino, Marta
Riga, Alessandra	Somma, Stefania	van Loo, Geert
Rimsa, Antonija	Soria, Marcelo A.	Varese, Giovanna Cristina
Riso, Patrizia	Sorrentino, Alida	Vaudano, Enrico
Rizzello, Carlo Giuseppe	Sousa R.,	Vazquez, Rosa
Robert, Vincent	Spada, Valentina	Venâncio, Armando
Rogora M.	Spadaro, Da vide	Venice, Francesco
Rojo, Maria Cecilia	Spagnoletta, Anna	Vera Thiel
Romac, Sarah	Spano, Giuseppe	Vereecke, Lars
Romano, Ida	Speranza, Barbara	Verrone, Laura
Romero-Mujalli D.	Spiezio, Caterina	Viale, Silvia
Rööp, Tiit	Spina, Federica	Vidgren, Virve
Rossetti, Lia	Spinelli, Veronica	Vikram Alva
Rubino, Luisa	Spiteri, Simona	Villano, Marianna
Rué, Olivier	Staňková, Eva	Vinnere Pettersson
Ruiz Martinez, José Luis	Stepanyan, Tamara	Visca, Andrea
Ruiz-Masó, José Ángel	Stewart, Helen	Vittoria, Maria
Runar, Stokke	Stilo, Giulia	Vitulo Nicola
Russo, Pasquale	Štšepetova, Jelena	Wang Z.
Ryan, Matthew	Stuknyte, Milda	Whitte, Julius Fredrich
Ryan, Matthew J.	Susca, Antonia	Wilmotte, Annick
Saar, Merili	Szymańska, Sonia	Wittouck, Stijn
Salveti, Elisa	Tabacchioni, Silvia	Wojciech, Rypniewski
Sandri, Camillo	Tagliasacchi, Ezher	Xiao, Hang
Sanna, Anna Maria Laura	Talia, Paola Monica	Yoav Soen
Santorufu, Lucia	Tangaro, Marco Antonio	Yurkov, Andrey
Saraf, Aniket	Tanmay Bharat	Zaccardelli, Massimo
Sbarra, Federico	Tedersoo L.	Zago, Miriam
Sbrana, Cristiana	Teturová, Kateřina	Zagorec, Monique
Scaffidi, Simona	Theopi Rados	Zanetti, Marco
Scarafile, Donatella	Thiran, Alexandra	Zara, Giacomo
Scarpetta, Davide	Thorp, Marie	Zotta, Teresa
Scauro, Angela	Tidona, Flavio	Zotti, Mirca
Sebastian Kadener	Tignat-Perrier, Romie	Zottini, Michela
Sedláček, Ivo	Todorov, Sanne Kjærulf	Zuzuarregui, Aurora
Selci, Matteo	Toniatti, Luca	
Senape, Veronica	Tontini, Beatrice	
Senkovs, Maris	Torres-Montero, Pablo	
Sepp, Epp	Tortarolo, Dora	
Serratore, Giovanna	Touati, Rim	
Sevi, Filippo	Trupo, Mario	
Siciliano, Rosa Anna	Tsolakis, Christos	
Sigillo, Loredana	Turchetti, Benedetta	
Signorello, Lara	Vaikjärv, Risto	
Silveti, Tiziana	Vaira, Anna Maria	

Sponsors

Platinum:



ThermoFisher
SCIENTIFIC

Thermo Fisher Scientific

Silver:



Bruker

The LABORMED logo consists of the word "LABORMED" in a blue, sans-serif font. The letters "O" and "R" are stylized with a grid-like pattern.


Haier Biomedical
Intelligent Protection of Life Science

Labormed


Others:

The 5Pascal logo features the word "5Pascal" in a blue, italicized, sans-serif font.

Levanchimica



Discover our latest solutions for proteomics

Product	Description
Stellar MS 	<p>Accelerate the transition from scientific research to clinical research at an unprecedented scale with the new Thermo Scientific™ Stellar™ mass spectrometer establishing a new paradigm to transition biomarker candidates from discovery to validation with –</p> <ul style="list-style-type: none"> • Extended scale • Higher sensitivity • Greater specificity • Massive throughput • Unrivaled productivity <p>thermofisher.com/Stellar</p>
Orbitrap Ascend Editions Tribid MS 	<p>The new Thermo Scientific™ Orbitrap™ Ascend Editions Tribid™ MS are application-specific editions with tailored features making them a fit for purpose, customized MS solution. Whether you're focused on single cell and multiplexed quantitative proteomics, metabolomics, lipidomics or glycomics, or deciphering higher-order protein structures, or small molecule-based pharmaceuticals or biotherapeutics, these customized features make the instrument uniquely suited to your needs</p> <p>thermofisher.com/OrbitrapAscend</p>
Ardia Platform 	<p>The Thermo Scientific™ Ardia™ Platform just got even more connected! It is now a scalable software solution for proteomics, metabolomics, biopharma, pharma and small molecules. New connections include –</p> <ul style="list-style-type: none"> • Thermo Scientific™ Proteome Discoverer™ software 3.2 for enhanced interpretation of complex datasets • CHIMERYS™ 4.0 for Proteome Discoverer 3.2 and Ardia Platform giving reduced analysis time for CHIMERYS and consensus workflows • Thermo Scientific™ Compound Discoverer™ software 3.4 for metabolomics, biopharma, pharma and lipidomics • Third party connections including direct data access from Skyline and the Ardia Platform's Data Sync 1.1 for automated bi-directional data transfer for other software <p>thermofisher.com/Ardia</p> <p><small>CHIMERYS is a trademark of MSAID, Skyline is from MacCoss Lab Software</small></p>
Vanquish Neo UHPLC System 	<p>New for the Thermo Scientific™ Vanquish™ Neo HPLC system is the Thermo Scientific™ PepMap™ Neo Nano/Capillary UHPLC columns for single and Tandem Direct Injection workflows. With 75 µm and 150 µm ID, these new columns facilitate up to 3 µL/min flow rates.</p> <p>thermofisher.com/lowflowhplc thermofisher.com/vanquishneo</p>

Thermo Scientific laboratory products

ThermoFisher
SCIENTIFIC

Laboratory Equipment (LE)		Laboratory Plastic Essentials (LPE)									
     	   	        									



Thermo
SCIENTIFIC

applied
biosystems

invitrogen

gibco

Together, we make science
exponentially better

-150°C Cryogenic Freezer



DW-150W209

-86°C ULT Freezers

NEW
Salvum
TwinCool



DW-86L578BPST
DW-86L728BPST
DW-86L828BPST

Salvum



DW-86L579BPT/729BPT
DW-86L829BPT/959BPT

Touch Screen Series



DW-86L419/579
DW-86L729/829

LED Display Series



DW-86L338J/388J



DW-86L490J



DW-86L578J
DW-86L728J
DW-86L828J

-86°C ULT Freezers



DW-86L486E
DW-86L628E



DW-86L51J



DW-86L100J

Chest



DW-86W100J



DW-86W420J

Water Cooling Freezer



DW-86L828W/959W

Mini ULT Freezer



DW-80WZ15

CO₂ Incubators



HCP-80
HCP-168
HCP-258



HCP-80E
HCP-168E
HCP-258E

Biological Safety Cabinets

Classical Series



HR30-IIA2
HR40-IIA2

Type B2



HR40-IIB2



HR700-IIA2



HR900-IIA2-D/S
HR1200-IIA2-D/S



HR1200-IIA2
HR1500-IIA2

Smart Series



HR1200-IIA2-X
HR1500-IIA2-X
HR1800-IIA2-X

LN₂ Cryogenic Containers

Low Temperature Transport Trolley



YDD/YDS/YDH Series



YDC-3000H



CryoBio 13

LN₂ Cryogenic Containers



CryoBio 43



CryoBio 6S



CryoBio 11Z
CryoBio 20Z
CryoBio 34Z



Bio-2T
Bio-Air

Centrifuges



LX-75L2400R



LX-155T500R



LX-165T2R



LX-185T100R



LX-165T2-J



LX-60T100-J



LX-60T500-J



Microbial Analysis by MALDI-TOF and Infrared (FT-IR) spectroscopy system

The visionary dream that changed microbiology

Over a decade ago, Bruker changed microbiology with the development of the MALDI Biotyper®, the market-leading MALDI-TOF system for microbiology, now in its 4th generation with the MALDI Biotyper® sirius System.



Speed and efficiency: The MALDI Biotyper® accelerates identification, providing rapid results that save valuable time in critical scenarios.

Cost-effective: Embrace accuracy yet say goodbye to expensive sequencing. Our method offers cost-efficient, reliable identification.

Open microbiology concept: Seamlessly integrate the MALDI Biotyper® into any laboratory environment. It's adaptable, user-friendly, and versatile.

Applications beyond clinical microbiology: From water quality assessment to pharmaceutical and food testing, the MALDI Biotyper® leads the way.

Other innovations include the MBT Sepsityper® solution for direct identification from positive blood culture, and resistance test assays. The complementary IR Biotyper®, based on FT-IR spectroscopy, enables quick and easy strain typing for hospital hygiene and epidemiological studies. Bruker also offers MICRONAUT and UMIC portfolios, which are dedicated to antimicrobial susceptibility testing.



Microbial strain typing for source tracking – be it in the case of infected patients or a contamination in a production chain - is an essential step in hygiene management. Bruker's Fourier Transform - Infrared (FT-IR) spectroscopy system, the IR Biotyper®, offers an easy-to-use method for fast and effective microorganism strain typing, with a simple workflow that needs minimal operator training.

5Pascal

VACUUM & CRYOGENIC WORLD

5Pascal provides vacuum technology solutions and has expanded its offerings to include advanced cryogenic products, coating systems, and surface analysis equipment, reflecting the latest developments in material science.



EDWARDS

GAMMA
VACUUM

VACGEN

Atlas
TECHNOLOGIES
ESS

AR

SCIENTIFIC
INSTRUMENTS

CeramTec

TSL TORR SCIENTIFIC LTD

NOIVION

LK
TECHNOLOGIES

Testbourne Ltd
Products for Industry, Research & Development

TEO TRIBOtechnic

MeiVac
INC.



Our broad product range allows us to meet our customers' diverse and unique needs with precision and expertise.



Are you looking for the right vacuum solution?

**CHALLENGE US
WITH YOUR REQUEST!**



Our partners are at the forefront of their respective sectors, and we have in-house capabilities to design top-notch systems under our brand, including freeze dryers, vacuum ovens, and TVACs.



www.5pascal.it



info@5pascal.it

